




1.2 Reacting Masses & Volumes

Mark Schemes



| | |
|------------|---------------------------------|
| Course | DP IB Chemistry |
| Section | 1. Stoichiometric Relationships |
| Topic | 1.2 Reacting Masses & Volumes |
| Difficulty | Medium |

Exam Papers Practice

To be used by all students preparing for DP IB Chemistry SL
Students of other boards may also find this useful



1

The correct answer is **C** because:

| | Steps | Calculation |
|---|--|--------------------------------------|
| 1 | Calculate the moles of the gas using the formula: Amount of gas (mol) = $\frac{\text{Volume of gas (dm}^3\text{)}}{22.7\text{dm}^3}$ | $\frac{2.27}{22.7} = 0.1\text{ mol}$ |
| 2 | Only 20% of the nitrogen was used | $0.1 \times 0.20 = 0.020\text{ mol}$ |
| 3 | Mass of Ammonia Moles = $\frac{\text{Mass}}{\text{Mr}}$ Re-arrange for mass Mass = moles \times M_r | $0.020 \times 17.04 = 0.34\text{ g}$ |
| 4 | Nitrogen is a diatomic molecule so for each molecule of nitrogen 2 ammonia molecules will be produced $\text{N}_2 \Rightarrow 2\text{NH}_3$ | $0.34 \times 2 = 0.68\text{ g}$ |



2

The correct answer is **B** because:

| | Steps | Calculation |
|---|---|--|
| 1 | Balanced symbol equation for the cold reaction | $2\text{NaOH}(\text{aq}) + \text{Cl}_2(\text{g}) \rightarrow \text{NaOCl}(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{NaCl}(\text{aq})$ |
| 2 | Calculate the relative formula mass of the compound M_r | $22.99 + 16.00 + 35.45 = 74.44$ |
| 3 | Find the moles of the product | The balanced equation shows the ratio of the Cl_2 to NaOCl is 1:1, so 0.10 mol of Cl_2 gives 0.10 mol of the NaOCl |
| 4 | Mass of Compound Moles = mass / M_r Re-arrange for mass Mass = moles x M_r | $0.10 \times 74.44 = 7.44 \text{ g}$ |

3

 The correct answer is **A** because:

| | Steps | Calculation |
|---|--|--|
| 1 | Calculate the relative formula mass (M_r) of compound $\text{Mg}(\text{NO}_3)_2$ | $24.31 + (14.01 \times 2) + (16.00 \times 6) = 148.33$ |
| 2 | Moles of $\text{Mg}(\text{NO}_3)_2$ Moles = $\frac{\text{Mass}}{M_r}$ | Moles = $\frac{1.48}{148.33}$ = 0.010 mol |
| 3 | Ratio of moles in equation: 2 moles of $\text{Mg}(\text{NO}_3)_2$ produces 4 moles of NO_2 | $0.010 \times 2 = 0.020$ mol |
| 4 | Mass of NO_2 Moles = $\frac{\text{Mass}}{M_r}$ Re-arrange for mass Mass = moles $\times M_r$ | $0.02 \times 46.01 = 0.92$ g |



4

The correct answer is **D** because:

| | Steps | Calculation |
|---|--|--|
| 1 | Find the volume of carbon dioxide that 1 mole of X would produce | 0.1 mol of X \Rightarrow 9.08dm ³ of CO ₂ 1.0 mol of X \Rightarrow 90.8dm ³ of CO ₂ |
| 2 | Find the moles of carbon dioxide using the formula, moles = $\frac{\text{volume in dm}^3}{22.7\text{dm}^3}$ | $\frac{90.8}{22.7\text{dm}^3} = 4.0 \text{ mol}$ |
| 3 | Use the number of moles of carbon dioxide to deduce the moles of carbon | 1.0 mol C \Rightarrow 1.0 of CO ₂ 4.0 mol C \Rightarrow 4.0 of CO ₂ |
| 4 | Deduce the number of carbon atoms in X | X must contain 4 carbon atoms |

Exam Papers Practice

5

 The correct answer is **B** because:

| Steps | | Calculation |
|-------|--|--|
| 1 | The molar mass of an element would be found using the formula, $M_r = \frac{\text{mass}}{\text{moles}}$ | $M_r = \frac{0.65}{\text{moles}}$ |
| 2 | The moles of a gas at stp would be found using the formula, moles = $\frac{\text{volume in dm}^3}{22.7 \text{ dm}^3}$ | $\frac{606.9 / 1000}{22.7}$ $\frac{606.9}{22.7 \times 1000}$ |
| 3 | The question tells you it is a Group II metal so the reacting ratio is 1:1 $M(s) + O_2(g) \rightarrow 2MO(s)$ From this you can deduce the number of moles | Moles of metal = moles of oxygen = $\frac{606.9}{22.7 \times 1000}$ |
| 4 | Putting these two terms together in $M_r = \frac{\text{mass}}{\text{moles}}$ | $0.65 \div \frac{606.9}{22.7 \times 1000}$ OR $\frac{22.7 \times 1000 \times 0.65}{606.9}$ |



6

The correct answer is **C** because:

| | Steps | Calculation |
|---|--|---|
| 1 | Mass of nitrogen in 14 g of fertiliser- you are told it contains 20% nitrogen | $14 \times \frac{20}{100} = 2.8 \text{ g of N}$ |
| 2 | Moles of nitrogen $\text{Moles} = \frac{\text{Mass}}{M_r}$ | $\frac{2.8}{14.01} = 0.20 \text{ mol}$ |
| 3 | Concentration of nitrogen $\frac{\text{Concentration (mol /dm}^3\text{)}}{\text{Amount of substance (mol)}} = \frac{\text{Volume of solution (dm}^3\text{)}}{\text{Volume of solution (dm}^3\text{)}}$ | $\frac{0.20}{5} = 0.04 \text{ mol dm}^{-3}$ |

Exam Papers Practice



7

The correct answer is **A** because:

- The ideal gas equation is $pV = nRT$
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)
- The closer the conditions are for ideal behaviour, the more accurate calculations are
- Ideal behaviour is identical particles with no internal structure, no volume, random motion, no intermolecular forces and no loss of momentum in collisions
- At low pressure, a gas minimises contact with other particles (intermolecular force tends towards zero as distance tends towards infinity)
- At high temperatures intermolecular forces are massively outweighed by kinetic energy and any collision between particles is less likely to be inelastic (i.e. lose momentum)

B is incorrect as at low temperatures gases deviate from ideal behaviour

C is incorrect as at high pressures gases deviate from ideal behaviour

D is incorrect as at high pressures and low temperatures gases deviate from ideal behaviour



7

The correct answer is **A** because:

- The ideal gas equation is $pV = nRT$
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)
- The closer the conditions are for ideal behaviour, the more accurate calculations are
- Ideal behaviour is identical particles with no internal structure, no volume, random motion, no intermolecular forces and no loss of momentum in collisions
- At low pressure, a gas minimises contact with other particles (intermolecular force tends towards zero as distance tends towards infinity)
- At high temperatures intermolecular forces are massively outweighed by kinetic energy and any collision between particles is less likely to be inelastic (i.e. lose momentum)

B is incorrect as at low temperatures gases deviate from ideal behaviour

C is incorrect as at high pressures gases deviate from ideal behaviour

D is incorrect as at high pressures and low temperatures gases deviate from ideal behaviour

8

The correct answer is **B** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)

- **Rearranged for pressure:**

$$p = \frac{nRT}{V}$$

- **Number of moles of hydrogen:**

$$\text{moles } (n) = \frac{\text{mass}(m)}{\text{molar mass } (M_r)} = \frac{0.96}{2.02}$$

- **Known quantities:**

- Number of moles; $\frac{0.96}{2.02}$
- Volume; $7.0 \times 10^{-3} \text{ m}^3$
- Temperature; 303 K

- **Substitution into the equation:**

$$p = \frac{nRT}{V} = \frac{0.96 \times 8.314 \times 303}{2.02 \times 7.0 \times 10^{-3}} \text{ pa}$$

Exam Papers Practice



9

The correct answer is **B** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)
- **Rearranged for number of moles:** $n = \frac{pV}{RT}$
- **Known quantities:**
 - Pressure; 103000 Pa
 - Volume; $5730 \text{ cm}^3 = 0.00573 \text{ m}^3$
 - Temperature; $60^\circ\text{C} = 333 \text{ K}$

- **Substitution into the equation:**

$$n = \frac{103000 \times 0.00573}{8.314 \times 333}$$

- **Mass of oxygen, O_2 ($M_r = 32$):**

$$\text{mass (m)} = \text{moles (n)} \times \text{molar mass (M}_r\text{)}$$

$$\text{Exam Papers Practice} = \frac{103000 \times 0.00573}{8.314 \times 333} \times 32.00$$

10

The correct answer is **D** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)
- **Rearranged for temperature:**

$$T = \frac{pV}{nR}$$

- **Known quantities:**
 - Pressure; $1.00 \times 10^5 \text{ Pa}$
 - Volume; $1.247 \times 10^{-3} \text{ m}^3$
 - moles (n) = $\frac{\text{mass}(m)}{\text{molar mass}(M_r)} = \frac{5.35}{70.90}$
- **Substitution into the equation:**

$$T = \frac{70.90 \times (1.0 \times 10^5) \times (1.247 \times 10^{-3})}{(5.35 \times 8.314)}$$

Exam Papers Practice

11

The correct answer is **A** because:

- The temperature and number of moles remain constant throughout
- **Boyle's Law:** $P_1V_1 = P_2V_2$
 - Where: P_1 is the initial pressure, V_1 is the initial volume, P_2 is the final pressure, V_2 is the final volume
- **Therefore:**

$$V_2 = \frac{P_1}{P_2} \times V_1$$

- **Known quantities:**
 - Pressure (P_1); 101,000 Pa
 - Pressure (P_2); 2,020,000 Pa
 - Volume (V_1); $200\text{cm}^3 = 0.0002\text{m}^3$
- **Substitution into the equation**

$$V_2 = \frac{101000}{2020000} \times 0.0002 = 0.00001 \text{ m}^3 \text{ or } 10 \text{ cm}^3$$

Exam Papers Practice

12

The correct answer is **B** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)

- **Rearranged for number of moles:**

$$n = \frac{pV}{RT}$$

- **Known quantities:**

- Pressure; 100,000 Pa
- Volume; $960 \text{ cm}^3 = 0.00096 \text{ m}^3$
- Temperature; $20^\circ\text{C} = 293 \text{ K}$

- **Substitution into the equation:**

$$n = \frac{100000 \times 0.00096}{8.314 \times 293}$$

- 2 moles of sodium produce 1 mole of H_2 gas therefore double the moles of sodium was used $\left(\frac{100000 \times 0.00096 \times 2}{8.314 \times 293} \right)$

- **Mass of Sodium, Na ($M_r = 22.99$):**

$$\text{mass (m)} = (n) \times \text{molar mass (M}_r\text{)}$$

$$= \frac{100000 \times 0.00096 \times 2 \times 22.99}{8.314 \times 293}$$



13

The correct answer is **B** because:

- Ideal gas behaviour is seen with identical particles with no internal structure, no volume, random motion, no intermolecular forces and no loss of momentum in collisions
- As the pressure increases the volume decreases, showing the pressure of the gas is **inversely proportional** demonstrated by **Boyle's Law**:

$$P_1 V_1 = P_2 V_2$$

- Where: P_1 is the initial pressure, V_1 is the initial volume, P_2 is the final pressure, V_2 is the final volume
- As the pressure increases the chances of inelastic collisions between the particles increases, so the gas behaves non-ideally

A is incorrect as the increase in pressure is not going to make the real gas behave more like an ideal gas

C is incorrect as the particles will not be absorbed by the vessel wall, but they are more likely to collide inelastically

D is incorrect as the gas is behaving less like an ideal gas if it is partially liquified

Exam Papers Practice

14

The correct answer is **D** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is the volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), and T is the temperature (K)
- **Rearranged for number of moles:**

$$n = \frac{pV}{RT}$$

- **Known quantities:**
 - Pressure; = 100kPa = 100,000 Pa
 - Volume; $3 \text{ dm}^3 = 0.003 \text{ m}^3$
 - Temperature; $25^\circ\text{C} = 298 \text{ K}$
- **Substitution into the equation:**

$$n = \frac{100000 \times 0.003}{8.314 \times 298}$$

- **Mass of methane gas:**

$$\begin{aligned} \text{mass}(m) &= \text{moles}(n) \times \text{Molarmass}(M_r) \\ &= \frac{100000 \times 0.003 \times 16.05}{8.314 \times 298} \end{aligned}$$

- So the answer is option **D**

A is incorrect as the volume has been left in dm^3 instead of converting to m^3

B is incorrect as the pressure has been left in kPa instead of converting to Pa

C is incorrect as the temperature has been left in $^\circ\text{C}$ instead of converting to Kelvin

Exam tip: make sure you know how the units are converted; pressure Pa, volume m^3 , temperature K

15

The correct answer is **C** because:

- **The ideal gas equation is $pV = nRT$**
 - Where p is pressure (Pa), V is volume (m^3), n is the number of moles, R is the gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$) and T is temperature (K)
- **Rearranged for pressure:**

$$p = \frac{nRT}{V}$$

- **Known quantities:**
 - Volume; $2 \text{ dm}^3 = 0.002 \text{ m}^3$
 - Temperature; $275^\circ\text{C} + 273 = 548 \text{ K}$
 - Number of moles (n) = $1.5 \times 10^{-2} \text{ mol}$
- **Substitution into the equation:**

$$p = \frac{nRT}{V} = \frac{(1.5 \times 10^{-2}) \times 8.314 \times (275 + 273)}{2.0 \times 10^{-3}}$$

Exam tip: make sure the units are converted; pressure Pa, volume m^3 , temperature K

The conversions are

$$1 \text{ m}^3 = 1000 \text{ dm}^3 = 1,000,000 \text{ cm}^3$$

$$\text{temp in } ^\circ\text{C} + 273 = \text{temp in Kelvin}$$