


## 1.2 Sequences & Series Mark Schemes



**Exam Papers Practice** 

To be used by all students preparing for DP IB Maths Al SL Students of other boards may also find this useful



a) For a geometric sequence the common ratio, r, is given by

$$C = \frac{U_2}{U_1} = \frac{U_3}{U_2} = \frac{U_4}{U_3} \cdots$$

$$u_2 = 44$$
  $u_3 = 55$ 

sub uz and uz into r tormula

$$C = \frac{55}{44}$$

$$C = \frac{55}{44}$$

$$C = \frac{5}{4}$$

$$C = \frac{11}{4}$$

# Exams Papers Practice

rearrange for un

$$U_1 = \frac{44}{\left(\frac{5}{4}\right)}$$



$$S_n = \underbrace{u_1(r^n-1)}_{r-1}$$
 (in formula booklet)

sub ui, r and n=5

$$S_{5} = \frac{35.2((\frac{5}{4})^{5}-1)}{(\frac{5}{4})-1}$$

$$S_{5} = 288.8875$$

$$S_{5} = 289 (3sf)$$

Question 2 (a) For an anthmetic sequence the sum of the first n terms is given by

Example 20 a 
$$u = \frac{n}{2} (2u + (n-1)d)$$
 (in formula booklet)

Example 20 a  $u = 275$  Practice

sub in Si6, u and n=16 and solve for d' 16-1 and 16

Alternative GDC methods

- · Plot y=920 and y=8(55+152) and find intersection.
- · Use the algebraic solver.

For more help visit our website www.exampaperspractice.co.uk



b) 
$$S_n = \frac{n}{2}(2u_1 + (n-1)d)$$
 (in formula booklet)  
 $S_{16} = 920$   $d = 11$   
Sub in  $S_{16}$ ,  $d$  and  $n = 16$  and  $S_{0}$  for  $U_1$   
 $920 = \frac{16}{2}(2u_1 + (16-1)(11))$   
 $920 = 8(2u_1 + 165)$   
 $115 = 2u_1 + 165$   
 $-50 = 2u_1$   
Alternative GDC methods  
Plot  $y = 920$  and  $y = 8(2x + 165)$  and find intersection.



a) For a geometric sequence the sum of the first in terms is given by

$$S_n = \underbrace{u_1(r^n-1)}_{r-1}$$
 (in formula booklet)

sub in Ss, us and n=5 and solve for r using your GDC



Alternative GDC methods

· Plot y=461.12 and y=200(x=1) and find intersection.

EXAMPLE the algebraic solver. Practice



b) 
$$S_n = u_1 (r^n - 1)$$
 (in formula booklet)

 $S_5 = 461.12$   $r = -2$ 

Sub in  $S_5$ ,  $r$  and  $r = 5$  and solve for  $u_1$  using your  $GDC$ 
 $461.12 = u_1 ((-2)^5 - 1)$ 
 $(-2) - 1$ 

Alternative  $GDC$  methods

Plot  $y = 461.12$  and  $y = 2(1.2)^5 - 1$  and find intersection.

Use the algebraic solver.



Question 4 a) For an arithmetic sequence the common difference, 
$$d$$
, is given by  $d = u_2 - u_1 = u_3 - u_2 = u_4 - u_3 \dots$ 
 $a_2 = 12$   $a_3 = 30$ 

sub in  $u_2$  and  $u_3$  into  $d$  formula

 $d = 30 - 12$ 
 $d = 18$ 

Use  $d = 18$  to find  $a_1$  and  $a_4$ 
 $18 = 12 - a_1$   $18 = a_4 - 30$ 
 $a_1 = -6$ 
 $a_4 = 48$ 



b) For a geometric sequence the common ratio, 
$$r_1$$
 is given by

$$\begin{aligned}
& (= \frac{U_2}{U_1} = \frac{U_3}{U_2} = \frac{U_4}{U_3} \\
& b_2 = 12 \\
& b_3 = 30 \\
& sub in b_2 and b_3 into r formula
\\
& (= \frac{30}{12}) \\
& (= 2.5) \\
& use r = 2.5 to find b_1 and b_4 \\
& 2.5 = \frac{12}{b_1} \\
& b_1 = 4.8
\end{aligned}$$

The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_2 = 12 \\
& b_3 = 30 \\
& b_4 = 75
\end{aligned}$$

The common ratio,  $r_1$  is given by

$$\begin{aligned}
& b_2 = 12 \\
& b_3 = 30 \\
& b_4 = 30
\end{aligned}$$

The common ratio,  $r_1$  is given by

$$\begin{aligned}
& b_2 = 12 \\
& b_3 = 30 \\
& b_4 = 30
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_2 = 12 \\
& b_3 = 30 \\
& b_4 = 30
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_4 = 75 \\
& b_4 = 75
\end{aligned}$$
The common ratio  $r_1$  is given by

$$\begin{aligned}
& b_2 = 12 \\
& b_3 = 30 \\
& b_4 = 75
\end{aligned}$$



The nth term tormula for an arithmetic sequence is given by

$$u_n = u_1 + (n-1)d$$
 (in formula booklet)

 $c_1 = 80$   $c_4 = 10$ 

sub  $c_1$  and  $c_4$  into the nth term formula to find  $d$ .

 $10 = 80 + (4-1)d$ 
 $3d = 10 - 80$ 
 $3d = -70$ 

Use the nth term formula to find  $c_2$  and  $c_3$ 
 $c_2 = 80 + (2-1)(-\frac{70}{3})$   $c_3 = 80 + (3-1)(-\frac{70}{3})$ 
 $c_4 = \frac{170}{3}$ 
 $c_5 = \frac{100}{3}$ 



d) The nth term formula for a geometric sequence is given by

$$U_{\Lambda} = U_{1} r^{\Lambda-1}$$
 (in formula booklet)

 $d_{1} = 80$   $d_{1} = 10$ 

sub di and dy into the nth term formula to find  $r$ 
 $10 = 80 r^{4-1}$ 
 $r = \frac{10}{80}$ 
 $r = \left(\frac{1}{8}\right)^{\frac{1}{3}}$ 

Use the nth term formula to find d2 and d3

 $d_{2} = 80\left(\frac{1}{2}\right)^{2-1}$ 
 $d_{3} = 80\left(\frac{1}{2}\right)^{3-1}$ 

Example 12 and 13 and 14 and 15 an





Alternative GDC methods

· Plot y = 26.5 and y = 4 + (x - 1)(1.5) and find intersection. · Use the algebraic solver.

c) 
$$S_n = \frac{n}{2} (2u_1 + (n-1)d)$$
 (in formula booklet)

 $u_1 = 4$   $d = 1.5$   $S_n = 220$ 

sub in  $u_1$ ,  $d$  and  $S_n$  into the formula

 $220 = \frac{n}{2} (2(4) + (n-1)(1.5))$ 
 $440 = n (8 + (n-1)(1.5))$ 

Put the equation into the algebraic solver in your 60c.

 $n = 15.0968$ 
 $S_{15} = 220$ 

Marie will complete a total of 120 km during the 16th week.

Alternative GDC method

Plot y = 220 and  $y = \frac{2}{2}(8 + (x - 1)(1.5))$  and find intersection.



Question 7 a) i) 
$$U_n = U_1 + (n-1)$$
 (in formula booklet)

 $U_8 = 18$   $d = 2$ 

Sub in  $u_8$  and  $d$  into the formula to find  $u_1$ 
 $18 = U_1 + (8-1)(2)$ 
 $18 = U_1 + 14$ 
 $u_1 = 4$ 

ii) Sub in  $u_1$  and  $d$  into the formula to find  $u_{17}$ 
 $u_{17} = 4 + (17-1)(2)$ 
 $u_{17} = 4 + 32$ 
 $u_{17} = 36$ 



b) i) beometric sequence

$$U_3 = 4$$
 $U_1 = U_1 \cap 1$ 

Sub in  $U_2$  and  $U_3$  into formula

 $U_4 = U_1 \cap 2$ 
 $U_5 = 36$ 
 $U_6 = U_1 \cap 3$ 
 $U_7 = 1$ 
 $U_8 = 36$ 
 $U_8 = U_1 \cap 3$ 
 $U_8 = U_1 \cap 3$ 

Alternative GDE methods

Plot  $O$  and  $O$  and find intersection.

Use the simultaneous equation solver.

Question 8

 $U_8 = U_1 \cap 3$ 
 $U_8 = U_1 \cap 3$ 





a) Using sigma notation
$$a_1 + a_2 + a_3 + ... + a_{12} = \sum_{k=1}^{12} a_k$$

$$\sum_{k=1}^{12} (32 - 7k)$$

$$S_n = \frac{n}{2} (2u_1 + (n-1)d) \text{ (in formula booklet)}$$

$$a_1 = 25 \qquad d = -7 \qquad n = 12$$

$$Sub \text{ in } a_1, d \text{ and } n$$

$$S_{12} = \frac{12}{2} (2(25) + (12-1)(-7))$$

$$S_{12} = -162$$
Alternative GOC method using sigma notation.



b) Using sigma notation
$$a_4 + a_5 + a_6 + ... + a_{15} = \sum_{k=4}^{15} a_k$$

$$\sum_{k=4}^{15} (32 - 7k)$$

$$S_n = \frac{n}{2} (2u_1 + (n-1)d) \text{ (in formula booklet)}$$

$$a_1 = 4 \qquad d = -7 \qquad n = 12$$
Sub in  $a_1$ ,  $d$  and  $n$ 

$$S_{12} = \frac{12}{2} (2(4) + (12-1)(-7))$$

$$S_{12} = -414$$
Alternative CDC method using sigma notation.

Alternative GDC method using sigma notation.



Question 10

a) Using sigma notation
$$g_1 + g_2 + g_3 + ... + g_{10} = \sum_{k=1}^{10} g_k$$

$$\sum_{k=1}^{10} (4 \times 3^{k-1})$$

$$S_n = \frac{u_1(r^n - 1)}{r^{-1}} \qquad \text{(in formula booklet)}$$

$$g_1 = 4 \qquad r = 3 \qquad n = 10$$

$$Sub \quad \text{in } g_1, r \quad \text{and } n$$

$$S_{10} = \frac{u_1(3^{10} - 1)}{3^{-1}}$$

$$S_{10} = \frac{u_1(3^{10} - 1)}{3^{-1}}$$

Sto = 118 000 (3sf)

EXAMELER FOR method using sigma notation.



b) Using sigma notation

$$g_8 + g_9 + g_{10} + ... + g_{18} = \sum_{k=8}^{18} g_k$$
 $\sum_{k=8}^{18} (4 \times 3^{k-1})$ 
 $k_{=8}$ 
 $S_1 = \frac{u_1(r^n-1)}{r^n-1}$  (in formula booklet)

 $g_1 = 8748$   $r = 3$   $n = 11$ 
 $sub_1 in_1 g_1, r_1 and_1 r_2$ 
 $s_{11} = 8748(3^{11}-1)$ 
 $s_{11} = 774 836 604$ 

Sin = 775 000 000 (3sf) actice

Alternative GDC method using sigma notation.



a) Identity the geometric sequence.

The common ratio, r, will be equal to the percentage of the remaining population every year (as a decimal).

Ropulation decrease is 2% (0.02) every year. Therefore the remaining population every year is

100% - 2% = 98%.

1 - 0.02 = 0.98

Hence r = 0.98

Ui = 68 000

C = 0.98

Be sure to select the correct value for n.

Ui: 2021, U2: 2022, U3: 2023... U10: 2030

Un = U1 r n-1

U10 = 68000 (0.98) 10-1

U10 = 56 694.84782

The expected population of kiwis Example 20303 5 56 700. Practice



b) 
$$u_n = u_1 r^{n-1}$$
 (in formula booklet)  $u_1 = 68000 r = 0.98$   $u_n < 50000$  sub in  $u_1$ ,  $r$  and  $u_n$  into the formula  $50000 > 68000 (0.98)^{n-1}$  solve the equation for  $n$  using your GDC swapping the nequality (>) to an equal sign (=)  $50000 = 68000 (0.98)^{n-1}$   $n = 16.22$ 

.:  $u_{16} > 50000 u_{17} < 50000$ 
 $u_{17} < 50000$ 
 $u_{18} : 2036$ 

The population of kiwis will fall below  $50000 = 10000$ 

### Question 12 a) Identify the geometric sequence Ce

Un = U, r n-1 (in formula booklet)

U1 = 240 r= 1.125 n= 5

sub in ui, r and n

Us = 240 (1.125)4

U5 = 384 km (85f)



b) 
$$S_{n} = \underbrace{u_{1}(r^{n}-1)}_{r-1}$$
 (in formula booklet)  
 $u_{1} = 240$   $r = 1.125$   $n = 10$   
Sub in  $u_{1}$ ,  $r$  and  $r$   
 $s_{10} = \frac{240(1.125^{10}-1)}{1.125-1}$   
 $s_{10} = 4310 \text{ km} (3sf)$ 

Question 13 a) i) 
$$U_n = U_1 \cap I_2 \cap I_3 \cap I_4 \cap I_4$$

### Examus pers Practice

Sn = 
$$\frac{u_1(r^n-1)}{r-1}$$
 (in formula booklet)

 $u_1 = 0.5$   $r = 3$   $n = 5$ 

Sub in  $u_1, r$  and  $n$ 
 $s = 0.5 (3^5-1)$ 
 $s = 0.5 (3^5-1)$ 



b) 
$$U_n = U_1 + (n-1)d$$
 (in formula booklet)

 $S_n = \frac{1}{2}(2u_1 + (n-1)d)$  (in formula booklet)

 $U_4 = 13.5$   $S_5 = 60.5$ 
 $13.5 = U_1 + 3d$  (i)  $60.5 = \frac{5}{2}(2u_1 + 4d)$  (2)

Input (1) and (2) into your GDC to solve for  $u_1$  and  $d_2$ .

 $u_1 = 9.3$   $d = 1.4$ 

Alternative GDC methods

1 · Plot (1) and (2) and find intersection.

• Input (1) and (2) into the simultaneous equation solver.