




# 1.2 Uncertainties & Errors

## Question Paper



Course	DP IB Physics
Section	1. Measurement & Uncertainties
Topic	1.2 Uncertainties & Errors
Difficulty	Medium

# Exam Papers Practice

To be used by all students preparing for DP IB Physics HL  
Students of other boards may also find this useful

### Question 1

Systematic and random errors can be compared.

What are the properties that apply to random errors?

- 1 The error is consistently too high or too low and by different amounts
- 2 The error is constantly too high or low and by the same amount each time
- 3 The error can be fully eliminated
- 4 The error cannot be fully eliminated
- 5 The error can be reduced by averaging repeated measurements
- 6 The error cannot be reduced by averaging repeated measurements

- A. 1, 3 and 6
- B. 2, 4 and 6
- C. 2, 3 and 6
- D. 1, 4 and 5

[1 mark]

### Question 2

The measurement of a physical quantity may be subject to random errors and systematic errors.

Which statement is correct?

- A. random errors can be reduced by taking the average of several measurements
- B. random errors are always caused by the person taking the measurement
- C. a systematic error cannot be reduced by adjusting the apparatus
- D. a systematic error results in a different reading each time the measurement is taken

[1 mark]

### Question 3

A stone falls from rest to the bottom of a water well of depth  $d$ . The time  $t$  taken to fall is  $3.0 \pm 0.3$  s. The depth of the well is calculated to be 30 m using  $d = \frac{1}{2}at^2$ . The uncertainty in  $a$  is negligible.

What is the absolute uncertainty in  $d$ ?

- A.  $\pm 0.6$  m
- B.  $\pm 3$  m
- C.  $\pm 24$  m
- D.  $\pm 6$  m


[1 mark]



# Exam Papers Practice

### Question 4

The diagram shows an experiment to measure the speed of a small ball falling at constant speed through a clear liquid in a glass tube.



There are two marks on the tube. The top mark is positioned at  $15 \pm 1$  mm on the adjacent rule and the lower mark at  $285 \pm 1$  mm. The ball passes the top mark at  $2.50 \pm 0.02$  s and passes the lower mark at  $4.50 \pm 0.02$  s.

The constant speed of the ball is calculated to be  $135 \text{ mm s}^{-1}$ .

Which expression calculates the fractional uncertainty in the value of this speed?

- A.  $\frac{2}{270} + \frac{0.04}{2.00}$
- B.  $\frac{1}{270} + \frac{0.02}{2.00}$
- C.  $\frac{1}{15} + \frac{0.02}{2.50}$
- D.  $\frac{1}{285} + \frac{0.02}{4.50}$

[1 mark]

### Question 5

The strain energy  $W$  of a spring is determined from its spring constant  $k$  and extension  $x$ . The spring obeys Hooke's law and the value of  $W$  is calculated using the equation shown.

$$W = \frac{1}{2} kx^2$$

The spring constant  $k$  is  $200 \pm 1 \text{ N m}^{-1}$  and the extension  $x$  is  $0.040 \pm 0.004 \text{ m}$ .

What is the percentage uncertainty in the calculated value of  $W$ ?

- A. 20%
- B. 20.5%
- C. 10%
- D. 10.5%

[1 mark]

### Question 6

In an experiment investigating the electrolysis of copper, a student sets out to find the electrochemical equivalent,  $Z$ .

The electrochemical equivalent of a substance is the amount of substance deposited on a cathode per Coulomb of charge.

This can be determined using the equation:

$$Z = \frac{m_1 - m_2}{It}$$

Where:

- Mass of cathode before passing current,  $m_1 = (54.39 \pm 0.01) \times 10^{-3} \text{ kg}$
- Mass of cathode after passing current,  $m_2 = (52.06 \pm 0.01) \times 10^{-3} \text{ kg}$
- Current,  $I = 3.00 \pm 1 \text{ A}$
- Time,  $t = 4800 \pm 100 \text{ s}$

What is the largest possible value of  $Z$  from these readings?

- A.  $\frac{233}{940} \times 10^{-6} \text{ kg C}^{-1}$
- B.  $\frac{231}{940} \times 10^{-6} \text{ kg C}^{-1}$
- C.  $\frac{235}{940} \times 10^{-6} \text{ kg C}^{-1}$
- D.  $\frac{253}{720} \times 10^{-6} \text{ kg C}^{-1}$

[1 mark]

### Question 7

The sides of a square are measured to be  $8.0 \pm 0.2$  cm.

Which of the following gives the area of the square and its uncertainty?

- A.  $64.0 \pm 0.2$  cm<sup>2</sup>
- B.  $64.0 \pm 0.4$  cm<sup>2</sup>
- C.  $64.0 \pm 3.2$  cm<sup>2</sup>
- D.  $64.0 \pm 1.6$  cm<sup>2</sup>

[1 mark]

### Question 8

In an experiment, a radio-controlled car takes  $1.50 \pm 0.05$  s to travel  $30.0 \pm 0.1$  m.



What is the car's average speed and the uncertainty in this value?

- A.  $20.0 \pm 0.732$  ms<sup>-1</sup>
- B.  $20.0 \pm 0.0366$  ms<sup>-1</sup>
- C.  $20.0 \pm 0.066$  ms<sup>-1</sup>
- D.  $20.0 \pm 9.91$  ms<sup>-1</sup>

Exam Papers Practice [1 mark]

### Question 9

The diagram shows a thermometer reading of a liquid's temperature, before and after heating.



What is the best estimate for the temperature increase of the liquid?

- A.  $(54.0 \pm 0.5)$  degrees
- B.  $(54 \pm 1.0)$  degrees
- C.  $(54 \pm 1)$  degrees
- D.  $(54.0 \pm 2.0)$  degrees



[1 mark]


# Exam Papers Practice

### Question 10

A student collects values of the time period,  $T$ , of a pendulum at different lengths of string,  $L$ . They want to investigate the relationship:

$$T = 2\pi\sqrt{\frac{L}{g}}$$

They plot the values on a graph along with the error bars associated with each point, as shown in the diagram.



What is the percentage uncertainty in the experimental value of  $g$ ?

- A.  $\frac{43}{32}\%$
- B.  $\frac{43}{1600}\%$
- C.  $\frac{43}{800}\%$
- D.  $\frac{215}{8000}\%$

[1 mark]