



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions


Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

1.5 Binomial T heorem



IB Maths - Revision Notes

AA SL

1.5.1 Binomial Theorem

Binomial Theorem

What is the Binomial Theorem?

- The **binomial theorem** (sometimes known as the binomial expansion) gives a method for expanding a **two-term** expression in a bracket raised to a power
 - A **binomial expression** is in fact any two terms inside the bracket, however in IB the expression will usually be linear
- To expand a bracket with a two-term expression in:
 - First choose the most appropriate parts of the expression to assign to a and b
 - Then use the formula for the binomial theorem:

$$(a + b)^n = a^n + {}^n C_1 a^{n-1} b + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n$$

- where ${}^n C_r = \frac{n!}{r!(n-r)!}$
 - See below for more information on ${}^n C_r$
 - You may also see ${}^n C_r$ written as $\binom{n}{r}$ or ${}_n C_r$

- You will usually be asked to find the first three or four terms of an expansion
- Look out for whether you should give your answer in **ascending** or **descending** powers of x
 - For **ascending** powers start with the constant term, a^n
 - For **descending** powers start with the term with x in
- You may wish to swap a and b over so that you can follow the general formula given in the formula book
- If you are not writing the full expansion you can either
 - show that the sequence continues by putting an ellipsis (...) after your final term
 - or show that the terms you have found are an approximation of the full sequence by using the sign for approximately equals to (\approx)

How do I find the coefficient of a single term?

- Most of the time you will be asked to find the coefficient of a term, rather than carry out the whole expansion
- Use the formula for the general term

Copyright

© 2024 Exam Papers Practice

$${}^n C_r a^{n-r} b^r$$

- The question will give you the power of x of the term you are looking for
 - Use this to choose which value of r you will need to use in the formula
 - This will depend on where the x is in the bracket
 - The laws of indices can help you decide which value of r to use:
 - For $(a + bx)^n$ to find the coefficient of x^r use $a^{n-r} (bx)^r$
 - For $(a + bx^2)^n$ to find the coefficient of x^r use $a^{\frac{n-r}{2}} (bx^2)^{\frac{r}{2}}$
 - For $(a + \frac{b}{x})^n$ look at how the powers will cancel out to decide which value of r to use
 - So for $(3x + \frac{2}{x})^8$ to find the coefficient of x^2 use the term with $r = 3$ and to find the constant term use the term with $r = 4$
 - There are a lot of variations of this so it is usually easier to see this by inspection of the exponents
- You may also be given the coefficient of a particular term and asked to find an unknown in the brackets
 - Use the laws of indices to choose the correct term and then use the binomial theorem formula to form and solve an equation

 **Exam Tip**

- Binomial expansion questions can get messy, use separate lines to keep your working clear and always put terms in brackets

Copyright

© 2024 Exam Papers Practice



Worked example

Find the first three terms, in ascending powers of x , in the expansion of $(3 - 2x)^5$.

$$a = 3 \quad b = -2x \quad n = 5$$

Substitute values into the formula for $(a+b)^n$

$$(a+b)^n = a^n + {}^n C_1 a^{n-1} b + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n$$

Question asks for ascending powers of x , so start with the constant term, a^n .

$$(3 - 2x)^5 = 3^5 + 5C_1 (3)^{5-1} (-2x) + 5C_2 (3)^{5-2} (-2x)^2 + \dots$$

Watch out for the negative

$$\approx 243 + 5 \times 81 \times -2x + 10 \times 27 \times 4x^2$$

$$\approx 243 - 810x + 1080x^2$$

$$(3 - 2x)^5 \approx 243 - 810x + 1080x^2$$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

The Binomial Coefficient nCr

What is ${}^n C_r$?

- If we want to find the number of ways to **choose** r items out of n different objects we can use the formula for ${}^n C_r$
 - The formula for r combinations of n items is ${}^n C_r = \frac{n!}{r!(n-r)!}$
 - This formula is given in the formula booklet along with the formula for the binomial theorem
 - The function ${}^n C_r$ can be written $\binom{n}{r}$ or ${}_n C_r$ and is often read as 'n chooser'
 - Make sure you can find and use the button on your GDC

How does ${}^n C_r$ relate to the binomial theorem?

- The formula ${}^n C_r = \frac{n!}{r!(n-r)!}$ is also known as a **binomial** coefficient
- For a binomial expansion $(a + b)^n$ the coefficients of each term will be ${}^n C_0, {}^n C_1$ and so on up to ${}^n C_n$

- The coefficient of the r^{th} term will be ${}^n C_r$

- ${}^n C_n = {}^n C_0 = 1$

- The binomial coefficients are symmetrical, so ${}^n C_r = {}^n C_{n-r}$

- This can be seen by considering the formula for ${}^n C_r$

- $${}^n C_{n-r} = \frac{n!}{(n-r)!(n-(n-r))!} = \frac{n!}{r!(n-r)!} = {}^n C_r$$

Exam Tip

- You will most likely need to use the formula for nCr at some point in your exam
 - Practice using it and don't always rely on your GDC
 - Make sure you can find it easily in the formula booklet



Worked example

Without using a calculator, find the coefficient of the term in x^3 in the expansion of $(1 + x)^9$.

$$n = 9, \quad a = 1, \quad b = x$$

Substitute values into the formula for the binomial theorem:

$$(a+b)^n = a^n + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n$$

$$\text{where } {}^n C_r = \frac{n!}{r!(n-r)!}$$

$$(1+x)^9 = \sum_{r=0}^9 {}^9 C_r (1)^{9-r} (x)^r$$

← Coefficient of x^3 occurs when $r=3$.

$$r = 3 \text{ gives } {}^9 C_3 \times (1)^{9-3} (x)^3$$

Non-calculator, so work out ${}^n C_r$ separately:

$$\begin{aligned} {}^9 C_3 &= \frac{9!}{3!(9-3)!} = \frac{9 \times 8 \times 7 \times \cancel{6} \times \cancel{5} \times \cancel{4} \times 3 \times 2}{(3 \times 2) (\cancel{6} \times \cancel{5} \times 4 \times 3 \times 2)} \\ &= \frac{9 \times 8 \times 7}{6} = 84 \end{aligned}$$

$$\begin{aligned} \text{so the term when } r=3 \text{ is } & 84 \times (1)^6 \times x^3 \\ & = 84x^3 \end{aligned}$$

$\text{Coefficient of } x^3 = 84$

Copyright

© 2024 Exam Papers Practice




Pascal's Triangle

What is Pascal's Triangle?

- Pascal's triangle is a way of arranging the binomial coefficients and neatly shows how they are formed
 - Each term is formed by adding the two terms above it
 - The first row has just the number 1
 - Each row begins and ends with a number 1
 - From the third row the terms in between the 1s are the sum of the two terms above it

PASCAL'S TRIANGLE



Exam Papers Practice

Copyright © 2024 Exam Papers Practice

How does Pascal's Triangle relate to the binomial theorem?

- Pascal's triangle is an alternative way of finding the binomial coefficients, ${}^n C_r$
 - It can be useful for finding for smaller values of n without a calculator
 - However for larger values of n it is slow and prone to arithmetic errors
- Taking the first row as zero, (${}^0 C_0 = 1$), each row corresponds to the n^{th} row and the term within that row corresponds to the r^{th} term

Exam Tip


- In the non-calculator exam Pascal's triangle can be helpful if you need to get the coefficients of an expansion quickly, provided the value of n is not too big



Worked example

Write out the 7th row of Pascal's triangle and use it to find the value of 6C_4 .

7th row of Pascal's Triangle:



7th row of Pascal's Triangle: 1, 6, 15, 20, 15, 6, 1
 ${}^6C_4 = 15$

Copyright

© 2024 Exam Papers Practice