



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions


Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

1.8 Eigenvalues & Eigenvectors



IB Maths - Revision Notes

AI HL



1.8.1 Eigenvalues & Eigenvectors

Characteristic Polynomials

Eigenvalues and **eigenvectors** are properties of square matrices and are used in a lot of real-life applications including geometrical transformations and probability scenarios. In order to find these eigenvalues and eigenvectors, the **characteristic polynomial** for a matrix must be found and solved.

What is a characteristic polynomial?

- For a matrix \mathbf{A} , if $\mathbf{Ax} = \lambda\mathbf{x}$ when \mathbf{x} is a non-zero vector and λ a **constant**, then λ is an **eigenvalue** of the matrix \mathbf{A} and \mathbf{x} is its corresponding **eigenvector**
- If $\mathbf{Ax} = \lambda\mathbf{x} \Rightarrow (\lambda\mathbf{I} - \mathbf{A})\mathbf{x} = 0$ or $(\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = 0$ and for \mathbf{x} to be a non-zero vector, $\det(\lambda\mathbf{I} - \mathbf{A}) = 0$
- The characteristic polynomial of an $n \times n$ matrix is:
$$p(\lambda) = \det(\lambda\mathbf{I} - \mathbf{A})$$
- In this course you will only be expected to find the characteristic equation for a 2×2 matrix and this will always be a **quadratic**

How do I find the characteristic polynomial?

- STEP 1
Write $\lambda\mathbf{I} - \mathbf{A}$, remembering that the identity matrix must be of the same order as \mathbf{A}
- STEP 2

Find the determinant of $\lambda\mathbf{I} - \mathbf{A}$ using the formula given to you in the formula booklet

$$\det \mathbf{A} = |\mathbf{A}| = ad - bc$$

Copyright

© 2024 Exam Papers Practice
Re-write as a polynomial

Exam Tip

- You need to remember the **characteristic equation** as it is **not** given in the formula booklet



 **Worked example**

Find the characteristic polynomial of the following matrix

$$A = \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix}$$

$$\begin{aligned} p(\lambda) &= \det(\lambda I - A) \\ &= \det\left(\lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix}\right) \\ &= \det\left(\begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix}\right) \\ &= \det\begin{pmatrix} \lambda - 5 & -4 \\ -3 & \lambda - 1 \end{pmatrix} \end{aligned}$$

Determinant of a 2×2 matrix	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow \det A = A = ad - bc$
--------------------------------------	---

$$\begin{aligned} &= (\lambda - 5)(\lambda - 1) - (-4)(-3) \\ &= \lambda^2 - 5\lambda - \lambda + 5 - 12 \end{aligned}$$

$$p(\lambda) = \lambda^2 - 6\lambda - 7$$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

Eigenvalues & Eigenvectors

How do you find the eigenvalues of a matrix?

- The eigenvalues of matrix **A** are found by solving the **characteristic polynomial** of the matrix
- For this course, as the characteristic polynomial will always be a **quadratic**, the polynomial will always generate one of the following:
 - **two real and distinct** eigenvalues,
 - **one real repeated** eigenvalue or
 - **complex** eigenvalues

How do you find the eigenvectors of a matrix?

- A value for **x** that satisfies the equation is an **eigenvector** of matrix **A**
- Any scalar multiple of **x** will also satisfy the equation and therefore there are an **infinite number** of eigenvectors that correspond to a particular eigenvalue
- STEP 1

Write $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$

- STEP 2
Substitute the eigenvalues into the equation $(\lambda I - \mathbf{A})\mathbf{x} = \mathbf{0}$, and form two equations in terms of **x** and **y**
- STEP 3
There will be an infinite number of solutions to the equations, so choose one by letting one of the variables be equal to **1** and using that to find the other variable

Exam Tip

- You can do a quick check to on your calculated eigenvalues as the values along the **leading diagonal** of the matrix you are analysing should **sum** to the **total of the eigenvalues** for the matrix

Copyright

© 2024 Exam Papers Practice

**Worked example**

Find the eigenvalues and associated eigenvectors for the following matrices.

a)
$$\mathbf{A} = \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix}.$$

Solve the characteristic polynomial to find the eigenvalues

$$p(\lambda) = \lambda^2 - 6\lambda - 7 \quad \leftarrow \text{From worked example above in Characteristic Polynomials}$$

$$(\lambda - 7)(\lambda + 1)$$

$$\Rightarrow \lambda = 7 \quad \lambda = -1$$

Use the eigenvalues in the equation $(\lambda I - A)x = 0$ to find the eigenvectors

$$\text{For } \lambda = 7 \Rightarrow \left(7 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{pmatrix} 7 & 0 \\ 0 & 7 \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -4 \\ -3 & 6 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left. \begin{array}{l} 2x - 4y = 0 \\ -3x + 6y = 0 \end{array} \right\} 2y = x$$

The eigenvector associated with $\lambda = 7$ is any multiple of $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$

Copyright

© 2024 Exam Papers Practice

$$\text{For } \lambda = -1 \Rightarrow \left(-1 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} - \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -6 & -4 \\ -3 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left. \begin{array}{l} -6x - 4y = 0 \\ -3x - 2y = 0 \end{array} \right\} 2y = -3x$$

The eigenvector associated with $\lambda = -1$ is any multiple of $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$

b)
$$\mathbf{B} = \begin{pmatrix} 1 & -5 \\ 2 & 3 \end{pmatrix}.$$



Find the characteristic polynomial

$$\begin{aligned}
 p(\lambda) &= \det \begin{pmatrix} \lambda-1 & 5 \\ -2 & \lambda-3 \end{pmatrix} \\
 &= (\lambda-1)(\lambda-3) - (5)(-2) \\
 &= \lambda^2 - 3\lambda - \lambda + 3 + 10
 \end{aligned}$$

$$p(\lambda) = \lambda^2 - 4\lambda + 13$$

Solve the characteristic polynomial to find the eigenvalues by hand or using the GDC

$$\begin{aligned}
 p(\lambda) &= \lambda^2 - 4\lambda + 13 = 0 \\
 (\lambda - 2)^2 - 4 + 13 &= 0
 \end{aligned}$$

$$(\lambda - 2)^2 = -9$$

$$\lambda = 2 \pm \sqrt{-9}$$

$$\lambda = 2 \pm 3i$$

Use the eigenvalues in the equation $(\lambda I - A)x = 0$ to find the eigenvectors

$$\text{For } \lambda = 2+3i \Rightarrow \left(\begin{pmatrix} 2+3i & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & -5 \\ 2 & 3 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{pmatrix} 2+3i & 0 \\ 0 & 2+3i \end{pmatrix} - \begin{pmatrix} 1 & -5 \\ 2 & 3 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1+3i & 5 \\ -2 & -1+3i \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned}
 (1+3i)x + 5y &= 0 \\
 -2x + (-1+3i)y &= 0
 \end{aligned}
 \left. \vphantom{\begin{aligned} (1+3i)x + 5y \\ -2x + (-1+3i)y \end{aligned}} \right\} 2x = (-1+3i)y$$

Both equations can be simplified to the same thing

The eigenvector associated with $\lambda = 2 + 3i$ is any multiple of $\begin{pmatrix} -1+3i \\ 2 \end{pmatrix}$

$$\text{For } \lambda = 2-3i \Rightarrow \left(\begin{pmatrix} 2-3i & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & -5 \\ 2 & 3 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\left(\begin{pmatrix} 2-3i & 0 \\ 0 & 2-3i \end{pmatrix} - \begin{pmatrix} 1 & -5 \\ 2 & 3 \end{pmatrix} \right) \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1-3i & 5 \\ -2 & -1-3i \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned}
 (1-3i)x + 5y &= 0 \\
 -2x + (-1-3i)y &= 0
 \end{aligned}
 \left. \vphantom{\begin{aligned} (1-3i)x + 5y \\ -2x + (-1-3i)y \end{aligned}} \right\} 2x = (-1-3i)y$$

Both equations can be simplified to the same thing

The eigenvector associated with $\lambda = 2 - 3i$ is any multiple of $\begin{pmatrix} -1-3i \\ 2 \end{pmatrix}$



1.8.2 Applications of Matrices

Diagonalisation

What is matrix diagonalisation?

- A **non-zero, square** matrix is considered to be **diagonal** if all elements **not** along its leading diagonal are **zero**
- A matrix **P** can be said to diagonalise matrix **M** , if **D** is a diagonal matrix where **$D = P^{-1}MP$**
- If matrix **M** has **eigenvalues** λ_1, λ_2 and **eigenvectors** $\mathbf{x}_1, \mathbf{x}_2$ and is diagonalisable by **P** , then
 - **$P = (\mathbf{x}_1 \ \mathbf{x}_2)$** , where the first column is the eigenvector \mathbf{x}_1 and the second column is the eigenvector \mathbf{x}_2
 - **$D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$**
- You will only need to be able to diagonalise 2×2 matrices
- You will only need to consider matrices with **real, distinct eigenvalues**
 - If there is only one eigenvalue, the matrix is either already diagonalised or cannot be diagonalised
 - Diagonalisation of matrices with complex or imaginary eigenvalues is outside the scope of the course

Exam Tip

- Remember to use the formula booklet for the **determinant** and **inverse** of a matrix

Copyright

© 2024 Exam Papers Practice



Worked example

The matrix $M = \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix}$ has the eigenvalues $\lambda_1 = 7$ and $\lambda_2 = -1$ with eigenvectors

$\mathbf{x}_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and $\mathbf{x}_2 = \begin{pmatrix} 2 \\ -3 \end{pmatrix}$ respectively.

Show that $P_1 = (\mathbf{x}_1 \ \mathbf{x}_2)$ and $P_2 = (\mathbf{x}_2 \ \mathbf{x}_1)$ both diagonalise M .

Show that $P^{-1}MP$ produces a diagonal matrix

Inverse of a 2×2 matrix	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow A^{-1} = \frac{1}{\det A} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}, ad \neq bc$
----------------------------------	---

$$P_1 = \begin{pmatrix} 2 & 2 \\ 1 & -3 \end{pmatrix} \Rightarrow P_1^{-1} = -\frac{1}{8} \begin{pmatrix} -3 & -2 \\ -1 & 2 \end{pmatrix} = \frac{1}{8} \begin{pmatrix} 3 & 2 \\ 1 & -2 \end{pmatrix}$$

$$\begin{aligned} D_1 &= P_1^{-1}MP_1 = \frac{1}{8} \begin{pmatrix} 3 & 2 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 2 \\ 1 & -3 \end{pmatrix} \\ &= \frac{1}{8} \begin{pmatrix} 3 & 2 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} 14 & -2 \\ 7 & 3 \end{pmatrix} \\ &= \frac{1}{8} \begin{pmatrix} 56 & 0 \\ 0 & -8 \end{pmatrix} \end{aligned}$$

$$D = \begin{pmatrix} 7 & 0 \\ 0 & -1 \end{pmatrix} \quad \leftarrow D = \text{Diagonal matrix of eigenvalues}$$

$$P_2 = \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \Rightarrow P_2^{-1} = \frac{1}{8} \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix}$$

$$\begin{aligned} D_2 &= P_2^{-1}MP_2 = \frac{1}{8} \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} 2 & 2 \\ -3 & 1 \end{pmatrix} \\ &= \frac{1}{8} \begin{pmatrix} 1 & -2 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} -2 & 14 \\ 3 & 7 \end{pmatrix} \\ &= \frac{1}{8} \begin{pmatrix} -8 & 0 \\ 0 & 56 \end{pmatrix} \end{aligned}$$

$$D = \begin{pmatrix} -1 & 0 \\ 0 & 7 \end{pmatrix} \quad \leftarrow D = \text{Diagonal matrix of eigenvalues}$$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

Matrix Powers

One of the main applications of diagonalising a matrix is to make it easy to find **powers** of the matrix, which is useful when modelling transient situations such as the movement of populations between two towns.

How can the diagonalised matrix be used to find higher powers of the original matrix?

- The equation to find the diagonalised matrix can be re-arranged for **M** :

$$D = P^{-1}MP \Rightarrow M = PDP^{-1}$$

- Finding higher powers of a matrix when it is diagonalised is straight forward:

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}^n = \begin{pmatrix} a^n & 0 \\ 0 & b^n \end{pmatrix}$$

- Therefore, we can easily find higher powers of the matrix using the **power formula** for a matrix found in the formula booklet:

$$M^n = PD^nP^{-1}$$

Exam Tip

- If you are asked to show this by hand, don't forget to use your GDC to **check** your answer afterwards!

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice



Worked example

The matrix $M = \begin{pmatrix} 3 & -2 \\ -4 & 1 \end{pmatrix}$ has the eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 5$ with eigenvectors $\mathbf{x}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $\mathbf{x}_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ respectively.

a) Show that M^n can be expressed as

$$M^n = -\frac{1}{3} \begin{pmatrix} (-(-1)^n - 2(5)^n) & (-(-1)^n + (5)^n) \\ (-2(-1)^n + 2(5)^n) & (-2(-1)^n - (5)^n) \end{pmatrix}$$

Find D , P and P^{-1}

$$D = \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} \quad P = \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \Rightarrow P^{-1} = -\frac{1}{3} \begin{pmatrix} -1 & -1 \\ -2 & 1 \end{pmatrix}$$

Use the matrix power formula from the formula booklet

Power formula for a matrix	$M^n = PD^nP^{-1}$	P is the matrix of eigenvectors and D is the diagonal matrix of eigenvalues
----------------------------	--------------------	---

$$M^n = -\frac{1}{3} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix}^n \begin{pmatrix} -1 & -1 \\ -2 & 1 \end{pmatrix}$$

$$= -\frac{1}{3} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} -1^n & 0 \\ 0 & 5^n \end{pmatrix} \begin{pmatrix} -1 & -1 \\ -2 & 1 \end{pmatrix}$$

When multiplying these expressions, be careful!
 $-2 \times 5^n = -2(5^n)$
NOT -10^n

$$= -\frac{1}{3} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} -(-1)^n & -(-1)^n \\ -2(5)^n & (5)^n \end{pmatrix}$$

$$M^n = -\frac{1}{3} \begin{pmatrix} (-(-1)^n - 2(5)^n) & (-(-1)^n + (5)^n) \\ (-2(-1)^n + 2(5)^n) & (-2(-1)^n - (5)^n) \end{pmatrix}$$

b) Hence find M^5 .

Substitute $n = 5$

$$M^5 = -\frac{1}{3} \begin{pmatrix} (-(-1)^5 - 2(5)^5) & (-(-1)^5 + (5)^5) \\ (-2(-1)^5 + 2(5)^5) & (-2(-1)^5 - (5)^5) \end{pmatrix}$$

$$= -\frac{1}{3} \begin{pmatrix} -6249 & 3126 \\ 6252 & -3123 \end{pmatrix}$$

$$M^5 = \begin{pmatrix} 2083 & -1042 \\ -2084 & 1041 \end{pmatrix}$$