




10.2 Functional Group Chemistry

Mark Schemes



| | |
|------------|---------------------------------|
| Course | DP IB Chemistry |
| Section | 10. Organic Chemistry |
| Topic | 10.2 Functional Group Chemistry |
| Difficulty | Medium |

Exam Papers Practice

To be used by all students preparing for DP IB Chemistry HL
Students of other boards may also find this useful

1

The correct answer is **C** because:


- A primary alcohol has a hydroxyl group (-OH) attached to a saturated carbon atom that has only 1 other carbon atom attached to it
- Secondary alcohol has a hydroxyl group (-OH) attached to a saturated carbon atom that has 2 other carbon atoms
- Xylitol has both primary and secondary hydroxyl groups
- Primary alcohols are oxidised to aldehydes and carboxylic acids
- Secondary alcohols are oxidised to ketones

2

The correct answer is **B** because:

- Bromine is an electrophile (electron loving species)
- It reacts with the alkene isomers via electrophilic addition by breaking the C=C double bond
- The best way to approach a question like this is to completely draw out the structural formula of each product:

Exam Papers Practice



- So, the reaction between HBr and isomers **X** and **Y** form:
 - (CH₃)₂CHCBr₃
 - (CH₃)₂CBrCHBr₂
 - CH₃CBr₂CHBrCH₃



3

The correct answer is **C** because:

- In the initiation step, free radical Cl^{\bullet} atoms are formed:
- Initiation step: $Cl_2 \rightarrow 2Cl^{\bullet}$
 - Initiation = net formation of radicals
- These free radical Cl^{\bullet} atoms remove a hydrogen from the ethane making an ethyl free radical, $C_2H_5^{\bullet}$
- Propagation step: $C_2H_6 + Cl^{\bullet} \rightarrow C_2H_5^{\bullet} + HCl$
 - Propagation = no change in the number of free radicals
- Two of these radicals can then pair-up in a termination step to form butane $CH_3CH_2CH_2CH_3$ (or C_4H_{10}):
- Termination step: $C_2H_5^{\bullet} + C_2H_5^{\bullet} \rightarrow C_4H_{10}$
 - Termination = net destruction of free radicals
- Therefore, option **C** is the only option which shows a possible compound that could be present

A, B and D are incorrect as

it's not possible to produce these products from ethane and chlorine

The correct answer is **C** because:

- In the initiation step, free radical Cl^{\bullet} atoms are formed:
- Initiation step: $Cl_2 \rightarrow 2Cl^{\bullet}$
 - Initiation = net formation of radicals
- These free radical Cl^{\bullet} atoms remove a hydrogen from the ethane making an ethyl free radical, $C_2H_5^{\bullet}$
- Propagation step: $C_2H_6 + Cl^{\bullet} \rightarrow C_2H_5^{\bullet} + HCl$
 - Propagation = no change in the number of free radicals
- Two of these radicals can then pair-up in a termination step to form butane $CH_3CH_2CH_2CH_3$ (or C_4H_{10}):
- Termination step: $C_2H_5^{\bullet} + C_2H_5^{\bullet} \rightarrow C_4H_{10}$
 - Termination = net destruction of free radicals
- Therefore, option **C** is the only option which shows a possible compound that could be present

A, B and D are incorrect as it's not possible to produce these products from ethane and chlorine



4

The correct answer is **A** because:

- In dichlorodifluoromethane, there are:
 - Two C–F bonds and two C–Cl bonds
- Both of these are incredibly strong bonds, and the carbon-fluorine bond energy is the larger of the two
- The C–F bond strength is the reason for the inertness of CCl_2F_2 and CFCs in general
- They are so strong that such bonds don't break under the energy of ordinary visible light
 - UV light is required to break them

B is incorrect as the polarity of a bond tells us how electronegative an atom is, which contributes to the strength of the bond, but is not the direct reason for the inertness of the molecule overall

C is incorrect as halogens are electronegative and create bond polarity, but this is not directly the reason for the inertness


D is incorrect as this is another consequence of the inertness of CCl_2F_2 , not a reason for it

Exam Papers Practice

5

The correct answer is **D** because:

- The reaction between bromoethane and sodium hydroxide is **nucleophilic substitution**
- It begins with the nucleophilic attack of OH^- from NaOH on the slightly positive carbon atom in the C-Br bond
- The leaving group is the Br^- ion, which is a species with a lone pair of electrons, thus both of the attacking and leaving groups are nucleophiles
- Therefore, **both** the attacking group and leaving groups are nucleophiles



6

Exam Papers Practice

The correct answer is **A** because:

- **Step 1:** Write out the equation for the complete combustion of propane
 - $\text{C}_3\text{H}_8 + \text{O}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$
- **Step 2:** Balance the equation
 - $\text{C}_3\text{H}_8 + 5\text{O}_2 \rightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$
- **Step 3:** Determine moles of oxygen using molar ratios
 - For every 1 mole of propane, 5 moles of oxygen are required
 - So, 0.1 mol of propane completely combusts in 0.5 mol of oxygen
- **Step 4:** Multiply moles of oxygen by 24 dm^3 to determine volume occupied by oxygen
 - Volume of oxygen = $0.5 \times 24 \text{ dm}^3 = 12 \text{ dm}^3$

7

The correct answer is **A** because:

- **Step 1:** Write the chemical equation
 - $\text{C}_2\text{H}_5\text{OH} + [\text{O}] \rightarrow \text{CH}_3\text{CHO} + \text{H}_2\text{O}$
- **Step 2:** Calculate the moles of ethanol
 - Moles = $\frac{\text{Mass}}{M_r}$
 - Moles = $\frac{2.3}{(12 \times 2 + 6 + 16)} = 0.05 \text{ mol}$
- **Step 3:** Calculate the mass of ethanal
 - From the chemical equation, we can see that 1 mol of ethanol produces 1 mol of ethanal
 - So, the number of moles of ethanal is 0.05 mol
- **Step 4:** Calculate the theoretical yield of ethanal
 - Mass = Moles \times M_r
 - So, the mass of ethanal = $0.05 \times ((12 \times 2) + (1 \times 4) + 16) = 0.05 \times 44 \text{ g}$
 - Therefore, the theoretical yield of ethanal is 2.2 g
- **Step 5:** Calculate the actual yield
 - % yield = $\frac{\text{actual}}{\text{theoretical}}$
 - So, the actual yield of ethanal = $\frac{50}{100} \times 2.2 \text{ g} = 1.1 \text{ g}$

Exam Paper Practice

8

The correct answer is **B** because:

- Primary alcohols are oxidised to aldehydes and then to carboxylic acids
- Secondary alcohols are only oxidised to ketones, and cannot be oxidised further
- Tertiary alcohols cannot be oxidised
- Therefore, the only alcohol which would give only **one** oxidation product is the secondary alcohol, butan-2-ol

A & C are incorrect as butan-1-ol and 2-methylpropan-1-ol are primary alcohols, so they could produce **two** products when oxidised: aldehydes and carboxylic acids

D is incorrect as 2-methylpropan-2-ol is a tertiary alcohol, so, it cannot be oxidised

Exam Papers Practice

9

The correct answer is **C** because:

- **Acidified potassium dichromate(VI)** is an oxidising agent
 - An **oxidising agent** is a species that **oxidises** another species and gets **reduced** itself
 - The reaction mixture turns green as the **orange $\text{Cr}_2\text{O}_7^{2-}$** ions (dichromate) are reduced to **green Cr^{3+}** ions
 - Only **primary alcohols, secondary alcohols** and **aldehydes** can be oxidised by acidified potassium dichromate(VI)
- **Hydroxyethanal** contains a **primary alcohol** and **aldehyde** group
- Since the compound is heated **under reflux** until no further oxidation takes place, both the primary alcohol and the aldehyde are oxidised to **carboxylic acid groups**
 - Primary alcohol \rightarrow aldehyde \rightarrow carboxylic acid
 - Aldehyde \rightarrow carboxylic acid
- The correct answer is therefore **C**


A is incorrect as **only** the **aldehyde** group has been oxidised

B is incorrect as this compound contains the **incorrect** number of **carbon atom** in the carbon chain (there should be only 2 carbon atoms and not 3)

D is incorrect as the molecule is not fully oxidised; the aldehyde can undergo further oxidation to a carboxylic acid

10

The correct answer is **B** because:



11


The correct answer is **A** because:

- Compound Q is an **ester** made from propanoic acid, CH₃CH₂CO₂H, and methanol, CH₃OH
- The naming of the ester starts with the alkyl attached to the alcohol – in this case methyl – and ends with the carboxylic acid, in this case propanoate
 - '-oate' is the suffix of compounds produced from carboxylic acids
- Therefore, the name is **methyl propanoate**
- The boiling point of methyl propanoate is **lower** than butanoic acid because it has only **dipole-dipole attractions** between the molecules, whereas butanoic acid has stronger **hydrogen bonds**



12

The correct answer is **C** because:



Exam Papers Practice