



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions


Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2.2 Quadratic Functions & Graphs



IB Maths - Revision Notes

AA SL



2.2.1 Quadratic Functions

Quadratic Functions & Graphs

What are the key features of quadratic graphs?

- A **quadratic** graph can be written in the form $y = ax^2 + bx + c$ where $a \neq 0$
- The value of a affects the shape of the curve
 - If a is **positive** the shape is **concave up** \cup
 - If a is **negative** the shape is **concave down** \cap
- The **y-intercept** is at the point $(0, c)$
- The **zeros or roots** are the solutions to $ax^2 + bx + c = 0$
 - These can be found by
 - Factorising
 - Quadratic formula
 - Using your GDC
 - These are also called the x -intercepts
 - There can be 0, 1 or 2 x -intercepts
 - This is determined by the value of the **discriminant**
- There is an **axis of symmetry** at $X = -\frac{b}{2a}$
 - This is given in your **formula booklet**
 - If there are two x -intercepts then the axis of symmetry goes through the midpoint of them
- The **vertex** lies on the axis of symmetry
 - It can be found by **completing the square**
 - The x -coordinate is $X = -\frac{b}{2a}$
- The y -coordinate can be found using the GDC or by calculating y when $X = -\frac{b}{2a}$
- If a is **positive** then the vertex is the **minimum point**
- If a is **negative** then the vertex is the **maximum point**

Copyright © 2024 Exam Papers Practice



Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

What are the equations of a quadratic function?

- $f(x) = ax^2 + bx + c$
 - This is the **general form**
 - It clearly shows the y-intercept $(0, c)$
 - You can find the axis of symmetry by $x = -\frac{b}{2a}$
 - This is given in the formula booklet
- $f(x) = a(x - p)(x - q)$
 - This is the **factorised form**
 - It clearly shows the roots $(p, 0)$ & $(q, 0)$
 - You can find the axis of symmetry by $x = \frac{p + q}{2}$
- $f(x) = a(x - h)^2 + k$
 - This is the **vertex form**



- It clearly shows the vertex (h, k)
- The axis of symmetry is therefore $x = h$
- It clearly shows how the function can be transformed from the graph $y = x^2$
 - Vertical stretch by scale factor a
 - Translation by vector $\begin{pmatrix} h \\ k \end{pmatrix}$

How do I find an equation of a quadratic?

- If you have the **roots** $x = p$ and $x = q$...
 - Write in **factorised form** $y = a(x - p)(x - q)$
 - You will need a third point to find the value of a
- If you have the **vertex** (h, k) then...
 - Write in **vertex form** $y = a(x - h)^2 + k$
 - You will need a second point to find the value of a
- If you have **three random points** (x_1, y_1) , (x_2, y_2) & (x_3, y_3) then...
 - Write in the **general form** $y = ax^2 + bx + c$
 - Substitute the three points into the equation
 - Form and solve a system of three linear equations to find the values of a , b & c

Exam Tip

- Use your GDC to find the roots and the turning point of a quadratic function
 - You do not need to factorise or complete the square
 - It is good exam technique to sketch the graph from your GDC as part of your working

Copyright


© 2024 Exam Papers Practice



Worked example

The diagram below shows the graph of $y = f(x)$, where $f(x)$ is a quadratic function.

The intercept with the y -axis and the vertex have been labelled.



Write down an expression for $y = f(x)$.

Copyright

© 2024 Exam Papers Practice

We have the vertex so use $y = a(x-h)^2 + k$

$$\text{Vertex } (-1, 8) : y = a(x - (-1))^2 + 8$$

$$y = a(x + 1)^2 + 8$$

Substitute the second point

$$x = 0, y = 6 : 6 = a(0 + 1)^2 + 8$$

$$6 = a + 8$$

$$a = -2$$

$$y = -2(x + 1)^2 + 8$$

2.2.2 Factorising & Completing the Square

Factorising Quadratics

Why is factorising quadratics useful?

- Factorising gives **roots (zeroes or solutions)** of a quadratic
- It gives the **x-intercepts** when drawing the graph

How do I factorise a monic quadratic of the form $x^2 + bx + c$?

- A monic quadratic is a quadratic where the coefficient of the x^2 term is 1
- You might be able to spot the factors by **inspection**
 - Especially if c is a **prime number**
- Otherwise find two numbers m and n .
 - A sum equal to b
 - $p + q = b$
 - A product equal to c
 - $pq = c$
- Rewrite bx as $mx + nx$
- Use this to factorise $x^2 + mx + nx + c$
- A shortcut is to write:
 - $(x + p)(x + q)$

How do I factorise a non-monic quadratic of the form $ax^2 + bx + c$?

- A non-monic quadratic is a quadratic where the coefficient of the x^2 term is not equal to 1
- If a , b & c have a common factor then first factorise that out to leave a quadratic with coefficients

Copyright that have **no common factors**

© 2024 You might be able to spot the factors by **inspection**

- Especially if a and/or c are **prime numbers**
- Otherwise find two numbers m and n .
 - A sum equal to b
 - $m + n = b$
 - A product equal to ac
 - $mn = ac$
- Rewrite bx as $mx + nx$
- Use this to factorise $ax^2 + mx + nx + c$
- A shortcut is to write:

$$\frac{(ax + m)(ax + n)}{a}$$



- Then factorise common factors from numerator to cancel with the a on the denominator

How do I use the difference of two squares to factorise a quadratic of the form $a^2x^2 - c^2$?

- The **difference of two squares** can be used when...
 - There is **no x term**
 - The **constant term is a negative**
- Square root the two terms a^2x^2 and c^2
- The two factors are the **sum of square roots** and the **difference of the square roots**
- A shortcut is to write:
 - $(ax + c)(ax - c)$

Exam Tip

- You can deduce the factors of a quadratic function by using your GDC to find the solutions of a quadratic equation
 - Using your GDC, the quadratic equation $6x^2 + x - 2 = 0$ has solutions $x = -\frac{2}{3}$ and $x = \frac{1}{2}$
 - Therefore the factors would be $(3x + 2)$ and $(2x - 1)$
 - i.e. $6x^2 + x - 2 = (3x + 2)(2x - 1)$

Worked example

Factorise fully:

a) $x^2 - 7x + 12$.

Copyright

© 2024 Exam Papers Practice

Find two numbers m and n such that

$$m+n = b = -7 \quad mn = c = 12$$

$$-4 + -3 = -7 \quad -4 \times -3 = 12$$

Split $-7x$ up and factorise

$$x^2 - 4x - 3x + 12$$

$$x(x-4) - 3(x-4)$$

$$(x-3)(x-4)$$

Shortcut

$$(x+m)(x+n)$$

$$(x-3)(x-4)$$

b) $4x^2 + 4x - 15$.



Find two numbers m and n such that

$$m+n=b=4 \quad mn=ac=4 \times -15=-60$$

$$10 + -6 = 4 \quad 10 \times -6 = -60$$

Split $4x$ up and factorise Shortcut

$$4x^2 + 10x - 6x - 15$$

$$\frac{(ax+m)(ax+n)}{a}$$

$$2x(2x+5) - 3(2x+5)$$

$$\frac{(4x+10)(4x-6)}{4}$$

$$(2x-3)(2x+5)$$

$$\frac{2(2x+5)2(2x-3)}{4}$$

$$(2x-3)(2x+5)$$

c) $18 - 50x^2$.

Factorise the common factor

$$2(9 - 25x^2)$$

Use difference of two squares

$$2(3-5x)(3+5x)$$

Copyright

© 2024 Exam Papers Practice



Completing the Square

Why is completing the square for quadratics useful?

- Completing the square gives the **maximum/minimum** of a quadratic function
 - This can be used to define the **range** of the function
- It gives the **vertex** when drawing the graph
- It can be used to **solve quadratic equations**
- It can be used to derive the **quadratic formula**

How do I complete the square for a monic quadratic of the form $x^2 + bx + c$?

- Half the value of b and write $\left(x + \frac{b}{2}\right)^2$
 - This is because $\left(x + \frac{b}{2}\right)^2 = x^2 + bx + \frac{b^2}{4}$
- Subtract the unwanted $\frac{b^2}{4}$ term and add on the constant c
 - $\left(x + \frac{b}{2}\right)^2 - \frac{b^2}{4} + c$

How do I complete the square for a non-monic quadratic of the form $ax^2 + bx + c$?

- Factorise out the a from the terms involving x
 - $a\left(x^2 + \frac{b}{a}x\right) + c$
 - Leaving the a alone will **avoid working with lots of fractions**
- Complete the square on the quadratic term

- Half $\frac{b}{a}$ and write $\left(x + \frac{b}{2a}\right)^2$
 - This is because $\left(x + \frac{b}{2a}\right)^2 = x^2 + \frac{b}{a}x + \frac{b^2}{4a^2}$

- Subtract the unwanted $\frac{b^2}{4a^2}$ term

- Multiply by a and add the constant c

- $a\left[\left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a^2}\right] + c$

- $a\left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a} + c$

Exam Tip

- Some questions may not use the phrase "completing the square" so ensure you can recognise a quadratic expression or equation written in this form
 - $a(x - h)^2 + k (= 0)$



 **Worked example**

Complete the square:

a) $x^2 - 8x + 3$.

Half b and subtract its square

$$(x - 4)^2 - 4^2 + 3$$

$$(x - 4)^2 - 13$$

b) $3x^2 + 12x - 5$.

Factorise the 3 from the x terms

$$3(x^2 + 4x) - 5$$

Complete the square on $x^2 + 4x$

$$3((x+2)^2 - 2^2) - 5$$

Simplify

$$3((x+2)^2 - 4) - 5$$

$$3(x+2)^2 - 12 - 5$$

$$3(x+2)^2 - 17$$

Copyright

© 2024 Exam Papers Practice

Exam Papers Practice



2.2.3 Solving Quadratics

Solving Quadratic Equations

How do I decide the best method to solve a quadratic equation?

- A quadratic equation is of the form $ax^2 + bx + c = 0$
- If it is a calculator paper then use your GDC to solve the quadratic
- If it is a non-calculator paper then...
 - you can always use the **quadratic formula**
 - you can **factorise** if it can be factorised with integers
 - you can always **complete the square**

How do I solve a quadratic equation by the quadratic formula?

- If necessary **rewrite** in the form $ax^2 + bx + c = 0$
- **Clearly identify** the values of a, b & c
- **Substitute** the values into the formula
 - $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
 - This is given in the **formula booklet**
- **Simplify** the solutions as much as possible

How do I solve a quadratic equation by factorising?

- **Factorise** to rewrite the quadratic equation in the form $a(x - p)(x - q) = 0$
- Set each factor to zero and **solve**
 - $x - p = 0 \Rightarrow x = p$
 - $x - q = 0 \Rightarrow x = q$

How do I solve a quadratic equation by completing the square?

- **Complete the square** to rewrite the quadratic equation in the form $a(x - h)^2 + k = 0$
- Get the squared term by itself
 - $(x - h)^2 = -\frac{k}{a}$
- If $\left(-\frac{k}{a}\right)$ is **negative** then there will be **no solutions**



- If $\left(-\frac{k}{a}\right)$ is **positive** then there will be **two values** for $x - h$
 - $x - h = \pm \sqrt{-\frac{k}{a}}$
- **Solve** for x
 - $x = h \pm \sqrt{-\frac{k}{a}}$

Exam Tip

- When using the quadratic formula with awkward values or fractions you may find it easier to deal with the " $b^2 - 4ac$ " (**discriminant**) first
 - This can help avoid numerical and negative errors, improving accuracy

Worked example

Solve the equations:

a) $4x^2 + 4x - 15 = 0.$

This can be factorised

$(2x + 5)(2x - 3) = 0$

$2x + 5 = 0$ or $2x - 3 = 0$

$x = -\frac{5}{2}$ or $x = \frac{3}{2}$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice b) $3x^2 + 12x - 5 = 0.$

This can not be factorised but $3x^2$ and $12x$ have a common factor so complete the square

$3(x+2)^2 - 17 = 0$

$(x+2)^2 = \frac{17}{3}$ *Rearrange*

$x + 2 = \pm \sqrt{\frac{17}{3}}$ *Remember ±*

$x = -2 \pm \sqrt{\frac{17}{3}}$

c) $7 - 3x - 5x^2 = 0.$



This can not be factorised so use formula

Formula booklet

Solutions of a quadratic equation	$ax^2 + bx + c = 0 \Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, a \neq 0$
-----------------------------------	--

$$a = -5 \quad b = -3 \quad c = 7$$

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(-5)(7)}}{2(-5)}$$

$$= \frac{3 \pm \sqrt{9 + 140}}{-10}$$

$$x = -\frac{3 \pm \sqrt{149}}{10}$$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

2.2.4 Quadratic Inequalities

Quadratic Inequalities

What affects the inequality sign when rearranging a quadratic inequality?

- The inequality sign is **unchanged** by...
 - **Adding/subtracting** a term to both sides
 - **Multiplying/dividing** both sides by a **positive term**
- The inequality sign **flips** (< changes to >) when...
 - **Multiplying/dividing** both sides by a **negative term**

How do I solve a quadratic inequality?

- **STEP 1: Rearrange** the inequality into quadratic form with a **positive squared term**
 - $ax^2 + bx + c > 0$
 - $ax^2 + bx + c \geq 0$
 - $ax^2 + bx + c < 0$
 - $ax^2 + bx + c \leq 0$
- **STEP 2:** Find the **roots** of the quadratic equation
 - Solve $ax^2 + bx + c = 0$ to get x_1 and x_2 where $x_1 < x_2$
- **STEP 3: Sketch** a graph of the quadratic and label the roots
 - As the squared term is positive it will be **concave up** so "U" shaped
- **STEP 4: Identify** the **region** that satisfies the inequality
 - If you want the graph to be **above the x-axis** then choose the region to be the **two intervals outside** of the two roots
 - If you want the graph to be **below the x-axis** then choose the region to be the **interval between** the two roots

Copyright © 2024 Exam Papers Practice

- For $ax^2 + bx + c > 0$
 - The solution is $x < x_1$ or $x > x_2$
- For $ax^2 + bx + c \geq 0$
 - The solution is $x \leq x_1$ or $x \geq x_2$
- For $ax^2 + bx + c < 0$
 - The solution is $x_1 < x < x_2$
- For $ax^2 + bx + c \leq 0$
 - The solution is $x_1 \leq x \leq x_2$

How do I solve a quadratic inequality of the form $(x - h)^2 < n$ or $(x - h)^2 > n$?

- The safest way is by following the steps above
 - Expand and rearrange
- A **common mistake** is writing $x - h < \pm\sqrt{n}$ or $x - h > \pm\sqrt{n}$



- This is **NOT correct!**
- The correct solution to $(x-h)^2 < n$ is
 - $|x-h| < \sqrt{n}$ which can be written as $-\sqrt{n} < x-h < \sqrt{n}$
 - The **final solution** is $h - \sqrt{n} < x < h + \sqrt{n}$
- The correct solution to $(x-h)^2 > n$ is
 - $|x-h| > \sqrt{n}$ which can be written as $x-h < -\sqrt{n}$ or $x-h > \sqrt{n}$
 - The **final solution** is $x < h - \sqrt{n}$ or $x > h + \sqrt{n}$

Exam Tip

- It is easiest to sketch the graph of a quadratic when it has a positive x^2 term, so rearrange first if necessary
- Use your GDC to help select the correct region(s) for the inequality
- Some makes/models of GDC may have the ability to solve inequalities directly
 - However unconventional notation may be used to display the answer (e.g. $6 > x > 3$ rather than $3 < x < 6$)
 - The safest method is to **always** sketch the graph



 **Worked example**

Find the set of values which satisfy $3x^2 + 2x - 6 > x^2 + 4x - 2$.

STEP 1: Rearrange

$$(3x^2 + 2x - 6) - (x^2 + 4x - 2) > 0$$

This way gives $a > 0$


$$2x^2 - 2x - 4 > 0$$
$$x^2 - x - 2 > 0$$

Divide by factor of 2


STEP 2: Find the roots

$$x^2 - x - 2 = 0$$
$$(x - 2)(x + 1) = 0$$
$$x = 2 \text{ or } x = -1$$

STEP 3: Sketch



STEP 4: Identify region



$$x < -1 \text{ or } x > 2$$

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice



2.2.5 Discriminants

Discriminants

What is the discriminant of a quadratic function?

- The discriminant of a quadratic is denoted by the Greek letter Δ (upper case delta)
- For the quadratic function the discriminant is given by
 - $\Delta = b^2 - 4ac$
 - This is given in the **formula booklet**
- The discriminant is the expression that is square rooted in the **quadratic formula**


How does the discriminant of a quadratic function affect its graph and roots?

- If $\Delta > 0$ then $\sqrt{b^2 - 4ac}$ and $-\sqrt{b^2 - 4ac}$ are **two distinct values**
 - The equation $ax^2 + bx + c = 0$ has **two distinct real solutions**
 - The graph of $y = ax^2 + bx + c$ has **two distinct real roots**
 - This means the graph **crosses** the x-axis **twice**
- If $\Delta = 0$ then $\sqrt{b^2 - 4ac}$ and $-\sqrt{b^2 - 4ac}$ are **both zero**
 - The equation $ax^2 + bx + c = 0$ has **one repeated real solution**
 - The graph of $y = ax^2 + bx + c$ has **one repeated real root**
 - This means the graph **touches** the x-axis at **exactly one point**
 - This means that the **x-axis** is a **tangent** to the graph
- If $\Delta < 0$ then $\sqrt{b^2 - 4ac}$ and $-\sqrt{b^2 - 4ac}$ are **both undefined**
 - The equation $ax^2 + bx + c = 0$ has **no real solutions**
 - The graph of $y = ax^2 + bx + c$ has **no real roots**
 - This means the graph **never touches** the **x-axis**
 - This means that graph is **wholly above** (or **below**) the **x-axis**

Copyright


© 2024 Exam Papers Practice

IF $b^2 - 4ac > 0$




TWO DISTINCT
REAL ROOTS

IF $b^2 - 4ac = 0$



ONE REAL ROOT
(REPEATED ROOTS)

IF $b^2 - 4ac < 0$



NO REAL ROOTS

Exam Papers Practice

Copyright

© 2024 Exam Papers Practice

Forming equations and inequalities using the discriminant

- Often at least one of the coefficients of a quadratic is **unknown**
 - Questions usually use the letter k for the unknown constant
- You will be given a fact about the quadratic such as:
 - The **number of solutions** of the equation
 - The **number of roots** of the graph
- To find the **value or range of values** of k
 - Find an **expression for the discriminant**
 - Use $\Delta = b^2 - 4ac$
 - Decide whether $\Delta > 0$, $\Delta = 0$ or $\Delta < 0$
 - If the question says there are **real roots** but does not specify how many then use $\Delta \geq 0$
- **Solve** the resulting equation or inequality



Exam Tip

- Questions will rarely use the word discriminant so it is important to recognise when its use is required
 - Look for
 - a number of roots or solutions being stated
 - whether and/or how often the graph of a quadratic function intercepts the X -axis
- Be careful setting up inequalities that concern "two real roots" ($\Delta \geq 0$) as opposed to "two real distinct roots" ($\Delta > 0$)

Worked example

A function is given by $f(x) = 2kx^2 + kx - k + 2$, where k is a constant. The graph of $y = f(x)$ has two distinct real roots.

a) Show that $9k^2 - 16k > 0$.

Two distinct real roots $\Rightarrow \Delta > 0$

Formula booklet

Discriminant	$\Delta = b^2 - 4ac$
--------------	----------------------

$$a = 2k \quad b = k \quad c = (-k + 2)$$

$$\Delta = k^2 - 4(2k)(-k + 2)$$

$$= k^2 + 8k^2 - 16k$$

$$= 9k^2 - 16k$$

$$\Delta > 0 \Rightarrow 9k^2 - 16k > 0$$

Copyright

© 2024 Exam Papers Practice

b) Hence find the set of possible values of k .

Solve the inequality

$$9k^2 - 16k = 0$$

$$k(9k - 16) = 0$$

$$k = 0 \text{ or } k = \frac{16}{9}$$

$$k < 0 \text{ or } k > \frac{16}{9}$$

