



# 2.2 Quadratic Functions & Graphs

### Contents

- \* 2.2.1 Quadratic Functions
- \* 2.2.2 Factorising & Completing the Square
- \* 2.2.3 Solving Quadratics
- \* 2.2.4 Quadratic Inequalities
- \* 2.2.5 Discriminants

Tights Reserved



### 2.2.1 Quadratic Functions

## **Quadratic Functions & Graphs**

#### What are the key features of quadratic graphs?

- A quadratic graph can be written in the form  $y = ax^2 + bx + c$  where  $a \ne 0$
- The value of a affects the shape of the curve
  - If a is **positive** the shape is **concave up** ∪
  - If a is **negative** the shape is **concave down**  $\cap$
- The **y-intercept** is at the point (0, c)
- The **zeros or roots** are the solutions to  $ax^2 + bx + c = 0$ 
  - These can be found by
    - Factorising
    - Quadratic formula
    - Using your GDC
  - These are also called the *x*-intercepts
  - There can be 0, 1 or 2 *x*-intercepts
    - This is determined by the value of the discriminant
- There is an axis of symmetry at  $x = -\frac{b}{2a}$ 
  - This is given in your formula booklet
  - If there are two x-intercepts then the axis of symmetry goes through the midpoint of them
- The **vertex** lies on the axis of symmetry
  - It can be found by **completing the square**
  - The x-coordinate is  $X = -\frac{b}{2a}$
  - The y-coordinate can be found using the GDC or by calculating y when  $x = -\frac{b}{2a}$
  - If a is **positive** then the vertex is the **minimum point**
  - If a is negative then the vertex is the maximum point



#### What are the equations of a quadratic function?

$$f(x) = ax^2 + bx + c$$

- This is the general form
- It clearly shows the y-intercept (0, c)
- You can find the axis of symmetry by  $X = -\frac{b}{2a}$ 
  - This is given in the formula booklet

$$f(x) = a(x-p)(x-q)$$

- This is the **factorised form**
- It clearly shows the roots (p, 0) & (q, 0)
- You can find the axis of symmetry by  $X = \frac{p+q}{2}$

$$f(x) = a(x-h)^2 + k$$

- This is the **vertex form**
- It clearly shows the vertex (h, k)
- The axis of symmetry is therefore X = h
- It clearly shows how the function can be transformed from the graph  $y = x^2$ 
  - Vertical stretch by scale factor a
  - Translation by vector  $\begin{pmatrix} h \\ k \end{pmatrix}$


## How do I find an equation of a quadratic?

- If you have the **roots** x = p and x = q...
  - Write in **factorised form** y = a(x p)(x q)
  - You will need a third point to find the value of a
- If you have the **vertex** (h, k) then...
  - Write in vertex form  $y = a(x h)^2 + k$
  - You will need a second point to find the value of a
- If you have **three random points**  $(x_1, y_1), (x_2, y_2) \& (x_3, y_3)$  then...
  - Write in the **general form**  $y = ax^2 + bx + c$
  - Substitute the three points into the equation
  - Form and solve a system of three linear equations to find the values of a, b & c



The diagram below shows the graph of y = f(x), where f(x) is a quadratic function.

The intercept with the y-axis and the vertex have been labelled.



Write down an expression for y = f(x).



We have the vertex so use 
$$y = a(x-h)^2 + k$$
  
Vertex  $(-1,8): y = a(x-(-1))^2 + 8$   
 $y = a(x+1)^2 + 8$ 

Substitute the second point

$$x = 0$$
,  $y = 6$ :  $6 = a(0+1)^2 + 8$   
 $6 = a + 8$   
 $a = -2$ 

$$y = -2(x+1)^2 + 8$$

Rights Reserved



# 2.2.2 Factorising & Completing the Square

## **Factorising Quadratics**

#### Why is factorising quadratics useful?

- Factorising gives roots (zeroes or solutions) of a quadratic
- It gives the **x-intercepts** when drawing the graph

### How do I factorise a monic quadratic of the form $x^2 + bx + c$ ?

- A monic quadratic is a quadratic where the coefficient of the  $x^2$  term is 1
- You might be able to spot the factors by inspection
  - Especially if c is a **prime number**
- Otherwise find two numbers *m* and *n* ...
  - A sum equal to b
    - p+q=b
  - A product equal to c
    - pq = c
- Rewrite bx as mx + nx
- Use this to factorise  $x^2 + mx + nx + c$
- A shortcut is to write:
  - (x+p)(x+q)

## How do I factorise a non-monic quadratic of the form $ax^2 + bx + c$ ?

- A non-monic quadratic is a quadratic where the coefficient of the  $x^2$  term is not equal to 1
- If a, b & c have a common factor then first factorise that out to leave a quadratic with coefficients that have **no common factors**
- You might be able to spot the factors by **inspection** 
  - Especially if a and/or c are **prime numbers**
- Otherwise find two numbers m and n..
  - A sum equal to b
    - m+n=b
  - A product equal to ac
    - = mn = ac
- Rewrite bx as mx + nx
- Use this to factorise  $ax^2 + mx + nx + c$
- A shortcut is to write:

$$\frac{(ax+m)(ax+n)}{a}$$


• Then factorise common factors from numerator to cancel with the a on the denominator

How do I use the difference of two squares to factorise a quadratic of the form  $a^2x^2 - c^2$ ?

ACY C



- The difference of two squares can be used when...
  - There is **no** *x* **term**
  - The constant term is a negative
- Square root the two terms  $a^2 x^2$  and  $c^2$
- The two factors are the **sum of square roots** and the **difference of the square roots**
- A shortcut is to write:
  - (ax + c)(ax c)





Factorise fully:

a) 
$$x^2 - 7x + 12$$
.

Find two numbers m and n such that

$$m+n=b=-7$$
  $mn=c=12$ 
 $-4+-3=-7$   $-4\times-3=12$ 

Split  $-7\times$  up and factorise Shortcut

 $x^2-4x-3x+12$   $(x+m)(x+n)$ 
 $x(x-4)-3(x-4)$   $(x-3)(x-4)$ 

### b) $4x^2 + 4x - 15$ .

Find two numbers m and n such that  

$$m+n=b=4$$
  $mn=ac=4x-15=-60$   
 $10+-6=4$   $10x-6=-60$   
Split 4 x up and factorise Shortcut  
 $4x^2+10x-6x-15$   $\frac{(ax+m)(ax+n)}{a}$   
 $2x(2x+5)-3(2x+5)$   $\frac{(4x+10)(4x-6)}{4}$   
 $\frac{(2x-3)(2x+5)}{4}$ 

c) 
$$18 - 50x^2$$
.



Factorise the common factor  $2(9-25x^2)$  Use difference of two squares 2(3-5x)(3+5x)

Rights Reserved



## Completing the Square

#### Why is completing the square for quadratics useful?

- Completing the square gives the **maximum/minimum** of a quadratic function
  - This can be used to define the range of the function
- It gives the **vertex** when drawing the graph
- It can be used to solve quadratic equations
- It can be used to derive the quadratic formula

## How do I complete the square for a monic quadratic of the form $x^2 + bx + c$ ?

• Half the value of b and write 
$$\left(x + \frac{b}{2}\right)^2$$

This is because 
$$\left(x + \frac{b}{2}\right)^2 = x^2 + bx + \frac{b^2}{4}$$

■ Subtract the unwanted  $\frac{b^2}{4}$  term and add on the constant c

$$\left(x + \frac{b}{2}\right)^2 - \frac{b^2}{4} + c$$

# How do I complete the square for a non-monic quadratic of the form $ax^2 + bx + c$ ?

• Factorise out the a from the terms involving x

$$a\left(x^2 + \frac{b}{a}x\right) + x$$

- Leaving the c alone will avoid working with lots of fractions
- Complete the square on the quadratic term

• Half 
$$\frac{b}{a}$$
 and write  $\left(x + \frac{b}{2a}\right)^2$ 

This is because 
$$\left(x + \frac{b}{2a}\right)^2 = x^2 + \frac{b}{a}x + \frac{b^2}{4a^2}$$

Subtract the unwanted 
$$\frac{b^2}{4a^2}$$
 term

Multiply by a and add the constant c

$$a \left[ \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right] + c$$

$$a\left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a} + c$$

ACY CHI



Complete the square:

a) 
$$x^2 - 8x + 3$$
.

Half b and subtract its square 
$$(x-4)^2-4^2+3$$

$$(x-4)^2-13$$

b) 
$$3x^2 + 12x - 5$$
.

$$3(x^2+4x)-5$$

$$3((x+2)^2-2^2)-5$$

$$3((x+2)^2-4)-5$$

$$3(x+2)^2-12-5$$

$$3(x+2)^2-17$$



# 2.2.3 Solving Quadratics

## **Solving Quadratic Equations**

#### How do I decide the best method to solve a quadratic equation?

- A quadratic equation is of the form  $ax^2 + bx + c = 0$
- If it is a calculator paper then use your GDC to solve the quadratic
- If it is a non-calculator paper then...
  - you can always use the quadratic formula
  - you can factorise if it can be factorised with integers
  - you can always complete the square

#### How do I solve a quadratic equation by the quadratic formula?

- If necessary **rewrite** in the form  $ax^2 + bx + c = 0$
- Clearly identify the values of a, b & c
- Substitute the values into the formula

$$X = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- This is given in the formula booklet
- Simplify the solutions as much as possible

#### How do I solve a quadratic equation by factorising?

- Factorise to rewrite the quadratic equation in the form a(x-p)(x-q)=0
- Set each factor to zero and **solve**

$$x - p = 0 \Rightarrow x = p$$

$$x - q = 0 \Rightarrow x = q$$

## How do I solve a quadratic equation by completing the square?

- Complete the square to rewrite the quadratic equation in the form  $a(x-h)^2+k=0$
- Get the squared term by itself


$$(x-h)^2 = -\frac{k}{a}$$

- If  $\left(-\frac{k}{a}\right)$  is **negative** then there will be **no solutions**
- If  $\left(-\frac{k}{a}\right)$  is **positive** then there will be **two values** for x-h



$$x - h = \pm \sqrt{-\frac{k}{a}}$$
Solve for x

$$x = h \pm \sqrt{-\frac{k}{a}}$$





Solve the equations:

a) 
$$4x^2 + 4x - 15 = 0$$
.

This can be factorised

$$(2x + 5)(2x - 3) = 0$$

$$2x+5=0$$
 or  $2x-3=0$ 

$$x = -\frac{5}{2} \quad \text{or} \quad x = \frac{3}{2}$$

b) 
$$3x^2 + 12x - 5 = 0$$
.

This can not be factorised but  $3x^2$  and 12x have a common

factor so complete the square

$$3(x+2)^2 - 17 = 0$$

$$3(x+2)^2 - 17 = 0$$
  
 $(x+2)^2 = \frac{17}{3}$  Rearrange

$$x + 2 = \pm \sqrt{\frac{17}{3}}$$
 Remember  $\pm$ 

$$x = -2 \pm \sqrt{\frac{17}{3}}$$

c) 
$$7 - 3x - 5x^2 = 0$$
.



# This can not be factorised so use formula

# Formula booklet

Solutions of a quadratic equation  $ax^2 + bx + c = 0 \implies x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} , a \neq 0$ 

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(-5)(7)}}{2(-5)}$$

$$=\frac{3 \pm \sqrt{9+140}}{-10}$$

$$x = -\frac{3 \pm \sqrt{144}}{10}$$

Topice. All Rights Reserved



## 2.2.4 Quadratic Inequalities

## **Quadratic Inequalities**

#### What affects the inequality sign when rearranging a quadratic inequality?

- The inequality sign is **unchanged** by...
  - Adding/subtracting a term to both sides
  - Multiplying/dividing both sides by a positive term
- The inequality sign **flips** (< changes to >) when...
  - Multiplying/dividing both sides by a negative term

#### How do I solve a quadratic inequality?

- STEP 1: Rearrange the inequality into quadratic form with a positive squared term
  - $ax^2 + bx + c > 0$
  - $ax^2 + bx + c \ge 0$
  - $ax^2 + bx + c < 0$
  - $ax^2 + bx + c \le 0$
- STEP 2: Find the roots of the quadratic equation
  - Solve  $ax^2 + bx + c = 0$  to get  $x_1$  and  $x_2$  where  $x_1 < x_2$
- STEP 3: Sketch a graph of the quadratic and label the roots
  - As the squared term is positive it will be **concave up** so "U" shaped
- STEP 4: Identify the region that satisfies the inequality
  - If you want the graph to be above the x-axis then choose the region to be the two intervals outside of the two roots
  - If you want the graph to be **below the x-axis** then choose the region to be the **interval between** the two roots
  - For  $ax^2 + bx + c > 0$ 
    - The solution is  $x < x_1$  or  $x > x_2$
  - For  $ax^2 + bx + c \ge 0$ 
    - The solution is  $x \le x_1$  or  $x \ge x_2$
  - For  $ax^2 + bx + c < 0$ 
    - The solution is  $x_1 < x < x_2$
  - For  $ax^2 + bx + c \le 0$ 
    - The solution is  $x_1 \le x \le x_2$


# How do I solve a quadratic inequality of the form $(x - h)^2 < n$ or $(x - h)^2 > n$ ?

- The safest way is by following the steps above
  - Expand and rearrange
- A common mistake is writing  $x h < \pm \sqrt{n}$  or  $x h > \pm \sqrt{n}$ 
  - This is NOT correct!
- The correct solution to  $(x h)^2 < n$  is



$$|x-h| < \sqrt{n} \text{ which can be written as } -\sqrt{n} < x-h < \sqrt{n}$$

- The final solution is  $h \sqrt{n} < x < h + \sqrt{n}$ The correct solution to  $(x h)^2 > n$  is
- - $|x-h| > \sqrt{n} \text{ which can be written as } x-h < -\sqrt{n} \text{ or } x-h > \sqrt{n}$  The final solution is  $x < h \sqrt{n} \text{ or } x > h + \sqrt{n}$





Find the set of values which satisfy  $3x^2 + 2x - 6 > x^2 + 4x - 2$ .

$$(3x^2+2x-6) - (x^2+4x-2) > 0$$
 This way gives  $a>0$ 

$$2x^2-2x-4>0$$
 gives  $x^2-x-2>0$  Divide by factor of 2


STEP 2: Find the roots

$$x^2 - x - 2 = 0$$


$$(x-2)(x+1)=0$$

$$x=2$$
 or  $x=-1$ 

STEP 3: Sketch



STEP 4: Identify region



$$x < -1$$
 or  $x > 2$ 



#### 2.2.5 Discriminants

#### **Discriminants**

#### What is the discriminant of a quadratic function?

- The discriminant of a quadratic is denoted by the Greek letter ∆ (upper case delta)
- For the quadratic function the discriminant is given by
  - $\Delta = b^2 4ac$ 
    - This is given in the formula booklet
- The discriminant is the expression that is square rooted in the **quadratic formula**

#### How does the discriminant of a quadratic function affect its graph and roots?

- If  $\Delta > 0$  then  $\sqrt{b^2 4ac}$  and  $-\sqrt{b^2 4ac}$  are two distinct values
  - The equation  $ax^2 + bx + c = 0$  has two distinct real solutions
  - The graph of  $y = ax^2 + bx + c$  has two distinct real roots
    - This means the graph **crosses** the x-axis **twice**
- If  $\Delta = 0$  then  $\sqrt{b^2 4ac}$  and  $-\sqrt{b^2 4ac}$  are **both zero** 
  - The equation  $ax^2 + bx + c = 0$  has one repeated real solution
  - The graph of  $y = ax^2 + bx + c$  has one repeated real root
    - This means the graph touches the x-axis at exactly one point
    - This means that the *x*-axis is a tangent to the graph
- If  $\Delta$  < 0 then  $\sqrt{b^2-4ac}$  and  $-\sqrt{b^2-4ac}$  are **both undefined** 
  - The equation  $ax^2 + bx + c = 0$  has no real solutions
  - The graph of  $y = ax^2 + bx + c$  has no real roots
    - This means the graph **never touches** the **x-axis**
    - This means that graph is **wholly above** (or **below**) the **x-axis**



#### Forming equations and inequalities using the discriminant

- Often at least one of the coefficients of a quadratic is **unknown** 
  - Questions usually use the letter *k* for the unknown constant
- You will be given a fact about the quadratic such as:
  - The **number of solutions** of the equation
  - The **number of roots** of the graph
- To find the value or range of values of k
  - Find an expression for the discriminant
    - Use  $\Delta = b^2 4ac$
  - Decide whether  $\Delta > 0$ ,  $\Delta = 0$  or  $\Delta < 0$ 
    - If the question says there are **real roots** but does not specify how many then use  $\Delta \ge 0$
  - Solve the resulting equation or inequality

Reserved



A function is given by  $f(x) = 2kx^2 + kx - k + 2$ , where k is a constant. The graph of y = f(x) has two distinct real roots.

a) Show that  $9k^2 - 16k > 0$ .

Two distinct real roots 
$$\Rightarrow \Delta > 0$$

Formula booklet Discriminant  $\Delta = b^2 - 4ac$ 
 $a = 2k$   $b = k$   $c = (-k+2)$ 
 $\Delta = k^2 - 4(2k)(-k+2)$ 
 $= k^2 + 8k^2 - 16k$ 
 $= 9k^2 - 16k$ 
 $\Delta > 0 \Rightarrow 9k^2 - 16k > 0$ 

b) Hence find the set of possible values of k.

Solve the inequality 
$$9k^2-16k=0$$
 $k(9k-16)=0$ 
 $k=0$  or  $k=\frac{16}{9}$ 
 $k<0$  or  $k>\frac{16}{9}$