

Energy Levels & Photon Emission

TOPIC QUESTIONS

Level	AS Level
Subject	Physics
Exam Board	AQA
Paper Type	Multiple Choice

Time Allowed : 30min

EXAM PAPERS PRACTICE


1. A particle has a de Broglie wavelength of 1.2 nm. What is the momentum of this particle?

- A. $5.5 \times 10^{-31} \text{ kg m s}^{-1}$
- B. $5.5 \times 10^{-22} \text{ kg m s}^{-1}$
- C. $5.5 \times 10^{-28} \text{ kg m s}^{-1}$
- D. $5.5 \times 10^{-25} \text{ kg m s}^{-1}$




EXAM PAPERS PRACTICE

2. The diagram below shows the energy levels of a mercury atom.




What transition produces a photon with the **longest** wavelength?



EXAM PAPERS PRACTICE


3. The diagram shows an energy-level diagram for a hydrogen atom.



How many discrete photon energies could be produced from these energy levels?

- A. 3
- B. 5
- C. 6
- D. 9

4. The diagram shows some energy levels of an atom.



The transition E_4 to E_1 corresponds to the emission of infrared radiation. A transition corresponding to the emission of X-rays would be

- A. E_3 to E_1
- B. E_3 to E_2
- C. E_4 to E_0
- D. E_4 to E_2

5. The diagram below is an example of an electron diffraction pattern.




Which row correctly states how the **momentum** and therefore **wavelength** of the incident electrons

For more help, please visit www.exampaperspractice.co.uk

should change to produce a pattern with a **smaller** radius?

	Momentum	Wavelength
A	Decreases	Decreases
B	Decreases	Increases
C	Increases	Increases
D	Increases	Decreases

6. The diagram below shows a simplified diagram of a fluorescent tube.



What is the purpose of the phosphor coating?

- A. This is where the electrons are ionised
- B. Electrons in the phosphor coating de-excite to release visible light photons
- C. Electrons in the phosphor coating de-excite to release UV light photons
- D. Without the phosphor coating, the electrons in the mercury atoms would not be excited

7. When a nucleus of the radioactive isotope ${}_{28}^{65}\text{Ni}$ decays, a β^- particle and an electron antineutrino are emitted.

How many protons and neutrons are there in the resulting daughter nucleus?

	Number of protons	Number of neutrons
A	28	65

B	29	65
C	29	36
D	30	35

8. What interactions are involved in the production of a strange particle and its decay into non-strange particles?

	Production	Decay
A	strong	weak
B	strong	strong
C	weak	strong
D	weak	weak

9. An atom of ${}^{16}_7\text{N}$ gains 3 electrons.

What is the specific charge of the ion?

- A $1.80 \times 10^7 \text{ C kg}^{-1}$
- B $-1.80 \times 10^7 \text{ C kg}^{-1}$
- C $4.19 \times 10^7 \text{ C kg}^{-1}$
- D $-4.19 \times 10^7 \text{ C kg}^{-1}$

10. What is the de Broglie wavelength of an alpha particle with kinetic energy 4.8 MeV? A. $1.5 \times 10^{-9} \text{ m}$

- B. $2.6 \times 10^{-21} \text{ m}$
- C. $3.0 \times 10^{-15} \text{ m}$
- D. $6.5 \times 10^{-15} \text{ m}$


11. A beam of electrons R is accelerated from rest by a potential difference of 200 V and another beam of electrons S is accelerated from rest by a potential difference of 1800 V.

When they behave like waves, what is the ratio of their respective wavelengths


$\lambda_S : \lambda_R$? A. 9 : 1

- B. 1 : 9
- C. 3 : 1
- D. 1 : 3

12. The energy level diagram below shows six possible transitions between the energy levels of an atom. Each transition between levels creates a photon of definite energy and frequency.



If the increase in frequency is from left to right, which of the following spectrum corresponds most closely to the transitions?



13. Transitions between three energy levels in a particular atom give rise to the three spectral lines of frequencies, in decreasing magnitude f_1 , f_2 and f_3 with wavelengths λ_1 , λ_2 and λ_3 .

Which of the following correctly related f_1 , f_2 and f_3 and λ_1 , λ_2 and λ_3

	Frequency	Wavelength
A	$f_3 = f_1 + f_2$	$\frac{1}{\lambda_1} = \frac{1}{\lambda_2} + \frac{1}{\lambda_3}$
B	$f_3 = f_1 - f_2$	$\frac{1}{\lambda_1} = \frac{1}{\lambda_2} + \frac{1}{\lambda_3}$
C	$f_1 = f_2 + f_3$	$\lambda_1 = \lambda_2 + \lambda_3$
D	$f_1 = f_2 + f_3$	$\frac{1}{\lambda_3} = \frac{1}{\lambda_1} - \frac{1}{\lambda_2}$

14. The diagram shows some of the energy levels for a hydrogen atom.



Not to scale

A free electron of kinetic energy 12.5 eV collides with a hydrogen atom in its ground state. The hydrogen atom is excited from its ground state to the first excited state. The kinetic energy of the free electron after the collision is

- A. 2.29 eV
- B. 22.9 eV
- C. 60.0 eV
- D. 3.38 eV

15. The ionization energy of hydrogen is 13.6eV

What is the de Broglie wavelength of the slowest electron that can ionise a hydrogen atom when it collides with it?

- A. 0.16 nm
- B. 0.33 nm
- C. 0.59 nm
- D. 0.78 nm



EXAM PAPERS PRACTICE