EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2.5 Transformations of Graphs

IB Maths - Revision Notes

AA SL

2.5.1 Translations of Graphs

Translations of Graphs

What are translations of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a **translation**:
  - the graph is **moved** (up or down, left or right) in the xy plane
    - Its position **changes**
  - the shape, size, and orientation of the graph remain **unchanged**
- A particular translation (how far left/right, how far up/down) is specified by a **translation vector** $\begin{pmatrix} x \\ y \end{pmatrix}$:
  - x is the **horizontal** displacement
    - **Positive** moves **right**
    - **Negative** moves **left**
  - y is the **vertical** displacement
    - **Positive** moves **up**
    - **Negative** moves **down**



What effects do horizontal translations have on the graphs and functions?

- A **horizontal translation** of the graph $y = f(x)$ by the vector $\begin{pmatrix} a \\ 0 \end{pmatrix}$ is represented by
  - $y = f(x - a)$
- The ***x*-coordinates change**
  - The value *a* is **subtracted** from them
- The ***y*-coordinates stay the same**
- The coordinates (x, y) become $(x + a, y)$
- **Horizontal** asymptotes **stay the same**
- **Vertical** asymptotes **change**
  - $x = k$ becomes $x = k + a$


y
y=f(x)
0
4
x
(2,−3)
y
y=f(x−2)
HORIZONTAL TRANSLATION BY THE VECTOR $\binom{2}{0}$
(2,0)
0
(6,0)
x
(4,−3)
x−COORDINATES CHANGE, y−COORDINATES STAY THE SAME

What effects do vertical translations have on the graphs and functions?

- A **vertical translation** of the graph $y = f(x)$ by the vector $\begin{pmatrix} 0 \\ b \end{pmatrix}$ is represented by
  - $y - b = f(x)$
  - This is often rearranged to $y = f(x) + b$
- The ***x*-coordinates stay the same**
- The ***y*-coordinates change**
  - The value *b* is **added** to them
- The coordinates (x, y) become $(x, y + b)$
- **Horizontal** asymptotes **change**
  - $y = k$ becomes $y = k + b$
- **Vertical** asymptotes **stay the same**





Exam Tip

- To get full marks in an exam make sure you use correct mathematical terminology
  - For example: Translate by the vector $\begin{pmatrix} 2 \\ -4 \end{pmatrix}$

Worked example

The diagram below shows the graph of $y = f(x)$.



a) Sketch the graph of $y = f(x+3)$.

$y = f(x+k)$ translation by $\begin{pmatrix} -k \\ 0 \end{pmatrix}$

Translate $y = f(x)$ by $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$

A becomes $(-4, 5)$

B becomes $(0, -3)$



b) Sketch the graph of $y = f(x) + 3$.

$y = f(x) + k$ translation by $\begin{pmatrix} 0 \\ k \end{pmatrix}$

Translate $y = f(x)$ by $\begin{pmatrix} 0 \\ 3 \end{pmatrix}$

A becomes $(-1, 8)$

B becomes $(3, 0)$



2.5.2 Reflections of Graphs

Reflections of Graphs

What are reflections of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a **reflection**:
  - the graph is **flipped** about one of the coordinate axes
    - Its orientation **changes**
  - the size of the graph remains **unchanged**
- A particular reflection is specified by an **axis of symmetry**:
  - $y = 0$
    - This is the *x*-axis
  - $x = 0$
    - This is the *y*-axis



What effects do horizontal reflections have on the graphs and functions?

- A **horizontal reflection** of the graph $y = f(x)$ about the *y*-axis is represented by
  - $y = f(-x)$
- The ***x*-coordinates change**
  - Their **sign** changes
- The ***y*-coordinates stay the same**

- The coordinates (x, y) become $(-x, y)$
- **Horizontal** asymptotes **stay the same**
- **Vertical** asymptotes **change**
  - $x = k$ becomes $x = -k$


y
y=f(x)
0
4
x
(2,−3)
y
y=f(−x)
REFLECTION IN THE y−AXIS
(−4,0)
(0,0)
0
x
(−2,−3)
x−COORDINATES CHANGE, y−COORDINATES STAY THE SAME
POINTS ON THE y−AXIS ARE UNAFFECTED

What effects do vertical reflections have on the graphs and functions?

- A **vertical reflection** of the graph $y = f(x)$ about the *x*-axis is represented by
  - $-y = f(x)$
  - This is often rearranged to $y = -f(x)$
- The ***x*-coordinates stay the same**
- The ***y*-coordinates change**
  - Their **sign** changes
- The coordinates (x, y) become $(x, -y)$
- **Horizontal** asymptotes **change**
  - $y = k$ becomes $y = -k$
- **Vertical** asymptotes **stay the same**


y
y=f(x)
0
4
x
(2,–3)
y
y–COORDINATES CHANGE,
x–COORDINATES STAY
THE SAME
(2,3)
(0,0)
(4,0)
0
x
POINTS ON THE
x–AXIS ARE
UNAFFECTED
y=–f(x)
REFLECTION IN
THE x–AXIS

Worked example

The diagram below shows the graph of $y = f(x)$.



a) Sketch the graph of $y = -f(x)$.

$y = -f(x)$ reflection in x-axis

A becomes $(-1, -5)$

B becomes $(3, 3)$



b) Sketch the graph of $y = f(-x)$.

$y = f(-x)$ reflection in y-axis

A becomes $(1, 5)$

B becomes $(-3, -3)$



2.5.3 Stretches of Graphs

Stretches of Graphs

What are stretches of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a **stretch**:
  - the graph is **stretched** about one of the coordinate axes by a scale factor
    - Its size **changes**
  - the orientation of the graph remains **unchanged**
- A particular stretch is specified by a **coordinate axis** and a **scale factor**:
  - The **distance** between a **point** on the graph and the **specified coordinate axis** is **multiplied** by the **constant scale factor**
  - The graph is stretched in the **direction** which is **parallel** to the **other coordinate axis**
  - For scale factors **bigger than 1**
    - the points on the graph get **further away** from the **specified coordinate axis**
  - For scale factors **between 0 and 1**
    - the points on the graph get **closer** to the **specified coordinate axis**
    - This is also sometimes called a **compression** but in your exam you must use the term **stretch** with the appropriate scale factor



What effects do horizontal stretches have on the graphs and functions?

- A **horizontal stretch** of the graph $y = f(x)$ by a scale factor *q* centred about the *y*-axis is represented by
  - $y = f\left(\frac{x}{q}\right)$
- The ***x*-coordinates change**
  - They are **divided** by *q*
- The ***y*-coordinates stay the same**

- The coordinates (x, y) become (qx, y)
- **Horizontal** asymptotes **stay the same**
- **Vertical** asymptotes **change**
  - $x = k$ becomes $x = qk$



What effects do vertical stretches have on the graphs and functions?

- A **vertical stretch** of the graph $y = f(x)$ by a scale factor *p* centred about the *x*-axis is represented by
  - $\frac{y}{p} = f(x)$
  - This is often rearranged to $y = pf(x)$
- The ***x*-coordinates stay the same**
- The ***y*-coordinates change**
  - They are **multiplied** by *p*
- The coordinates (x, y) become (x, py)
- **Horizontal** asymptotes **change**
  - $y = k$ becomes $y = pk$
- **Vertical** asymptotes **stay the same**



Exam Tip

- To get full marks in an exam make sure you use correct mathematical terminology
  - For example: Stretch vertically by scale factor ½
  - Do not use the word "compress" in your exam

Worked example

The diagram below shows the graph of $y = f(x)$.



a) Sketch the graph of $y = 2f(x)$.

$y = kf(x)$ vertical stretch scale factor k

Stretch $y=f(x)$ vertically scale factor 2

A becomes $(-1, 10)$

B becomes $(3, -6)$



b) Sketch the graph of $y = f(2x)$.

$y = f(kx)$ horizontal stretch scale factor $\frac{1}{k}$

Stretch $y=f(x)$ horizontally scale factor $\frac{1}{2}$

A becomes $(-\frac{1}{2}, 5)$

B becomes $(\frac{3}{2}, -3)$



2.5.4 Composite Transformations of Graphs

Composite Transformations of Graphs

What transformations do I need to know?

- $y = f(x+k)$ is **horizontal translation** by vector $\begin{pmatrix} -k \\ 0 \end{pmatrix}$
  - If k is **positive** then the graph moves **left**
  - If k is **negative** then the graph moves **right**
- $y = f(x) + k$ is **vertical translation** by vector $\begin{pmatrix} 0 \\ k \end{pmatrix}$
  - If k is **positive** then the graph moves **up**
  - If k is **negative** then the graph moves **down**
- $y = f(kx)$ is a **horizontal stretch** by scale factor $\frac{1}{k}$ centred about the *y*-axis
  - If $k > 1$ then the graph gets **closer** to the *y*-axis
  - If $0 < k < 1$ then the graph gets **further** from the *y*-axis
- $y = kf(x)$ is a **vertical stretch** by scale factor k centred about the *x*-axis
  - If $k > 1$ then the graph gets **further** from the *x*-axis
  - If $0 < k < 1$ then the graph gets **closer** to the *x*-axis
- $y = f(-x)$ is a **horizontal reflection** about the *y*-axis
  - A **horizontal reflection** can be viewed as a special case of a **horizontal stretch**
- $y = -f(x)$ is a **vertical reflection** about the *x*-axis
  - A **vertical reflection** can be viewed as a special case of a **vertical stretch**

How do horizontal and vertical transformations affect each other?

- **Horizontal and vertical transformations** are **independent** of each other
  - The horizontal transformations involved will need to be applied in their correct order
  - The vertical transformations involved will need to be applied in their correct order
- Suppose there are **two horizontal** transformation **H_1 then H_2** and **two vertical** transformations **V_1 then V_2** then they can be applied in the following orders:
  - Horizontal then vertical:
    - $H_1 H_2 V_1 V_2$
  - Vertical then horizontal:
    - $V_1 V_2 H_1 H_2$
  - Mixed up (provided that **H_1 comes before H_2** and **V_1 comes before V2**):
    - $H_1 V_1 H_2 V_2$
    - $H_1 V_1 V_2 H_2$
    - $V_1 H_1 V_2 H_2$
    - $V_1 H_1 H_2 V_2$

Exam Tip

- In an exam you are more likely to get the correct solution if you deal with one transformation at a time and sketch the graph after each transformation

Worked example

The diagram below shows the graph of $y = f(x)$.



Sketch the graph of $y = \frac{1}{2} f\left(\frac{x}{2}\right)$.

A vertical and horizontal transformation can be done in any order

$y = \frac{1}{2} f(x)$: vertical stretch scale factor $\frac{1}{2}$

$y = f\left(\frac{x}{2}\right)$: horizontal stretch scale factor 2

A becomes $\left(-2, \frac{5}{2}\right)$

B becomes $\left(6, -\frac{3}{2}\right)$



Composite Vertical Transformations af(x)+b

How do I deal with multiple vertical transformations?

- **Order matters** when you have **more than one vertical transformations**
- If you are asked to find the equation then **build up the equation** by looking at the transformations in order
  - A **vertical stretch** by scale factor *a* followed by a **translation** of $\begin{pmatrix} 0 \\ b \end{pmatrix}$
    - Stretch: $y = af(x)$
    - Then translation: $y = [af(x)] + b$
    - Final equation: $y = af(x) + b$
  - A **translation** of $\begin{pmatrix} 0 \\ b \end{pmatrix}$ followed by a **vertical stretch** by scale factor *a*
    - Translation: $y = f(x) + b$
    - Then stretch: $y = a[f(x) + b]$
    - Final equation: $y = af(x) + ab$
- If you are asked to determine the **order**
  - The order of vertical transformations **follows the order of operations**
  - First write the equation in the form $y = af(x) + b$
    - **First stretch vertically** by scale factor *a*
    - If *a* is negative then the **reflection and stretch** can be **done in any order**
    - **Then translate** by $\begin{pmatrix} 0 \\ b \end{pmatrix}$

Worked example

The diagram below shows the graph of $y = f(x)$.



Sketch the graph of $y = 3f(x) - 2$.

The order vertical transformations follows the order of operations

$y = 3f(x)$: Vertical stretch scale factor 3

$y = f(x) - 2$: Translate $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$

A becomes $(-1, 13)$

B becomes $(3, -11)$