



# 2.5 Transformations of Graphs

### **Contents**

- \* 2.5.1 Translations of Graphs
- \* 2.5.2 Reflections of Graphs
- \* 2.5.3 Stretches of Graphs
- \* 2.5.4 Composite Transformations of Graphs



## 2.5.1 Translations of Graphs

### **Translations of Graphs**

#### What are translations of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a translation:
  - the graph is **moved** (up or down, left or right) in the xy plane
    - Its position changes
  - the shape, size, and orientation of the graph remain **unchanged**
- A particular translation (how far left/right, how far up/down) is specified by a **translation vector**  $\begin{pmatrix} X \\ Y \end{pmatrix}$ 
  - x is the **horizontal** displacement
    - Positive moves right
    - Negative moves left
  - y is the **vertical** displacement
    - Positive moves up
    - Negative moves down



#### What effects do horizontal translations have on the graphs and functions?

- A horizontal translation of the graph y = f(x) by the vector  $\begin{pmatrix} a \\ 0 \end{pmatrix}$  is represented by
  - y = f(x a)
- The x-coordinates change
  - The value a is **subtracted** from them
- The y-coordinates stay the same
- The coordinates (x, y) become (x + a, y)
- Horizontal asymptotes stay the same
- Vertical asymptotes change
  - x = k becomes x = k + a




#### What effects do vertical translations have on the graphs and functions?

- A **vertical translation** of the graph y = f(x) by the vector  $\begin{pmatrix} 0 \\ b \end{pmatrix}$  is represented by
  - y b = f(x)
  - This is often rearranged to y = f(x) + b
- The x-coordinates stay the same
- The y-coordinates change
  - The value b is **added** to them
- The coordinates (x, y) become (x, y + b)
- Horizontal asymptotes change
  - y = k becomes y = k + b
- Vertical asymptotes stay the same



The diagram below shows the graph of y = f(x).



a) Sketch the graph of y = f(x+3).




y = f(x + k) translation by  $\begin{pmatrix} -k \\ 0 \end{pmatrix}$ 

Translate y=f(x) by  $\binom{-3}{0}$ 

A becomes (-4,5)

B becomes (0, -3)




b) Sketch the graph of y = f(x) + 3.

y = f(x) + k translation by  $\binom{0}{k}$ 

Translate y=f(x) by  $\binom{0}{3}$ 

A becomes (-1,8)

B becomes (3,0)





# 2.5.2 Reflections of Graphs

## **Reflections of Graphs**

#### What are reflections of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a reflection:
  - the graph is **flipped** about one of the coordinate axes
    - Its orientation changes
  - the size of the graph remains unchanged
- A particular reflection is specified by an **axis of symmetry**:
  - y=0
    - This is the *x*-axis
  - X = 0
    - This is the *y*-axis

#### What effects do horizontal reflections have on the graphs and functions?

- A horizontal reflection of the graph y = f(x) about the y-axis is represented by
  - y = f(-x)
- The x-coordinates change
  - Their **sign** changes
- The y-coordinates stay the same
- The coordinates (x, y) become (-x, y)
- Horizontal asymptotes stay the same
- Vertical asymptotes change




#### What effects do vertical reflections have on the graphs and functions?

- A vertical reflection of the graph y = f(x) about the x-axis is represented by
  - -y = f(x)
  - This is often rearranged to y = -f(x)
- The x-coordinates stay the same
- The y-coordinates change
  - Their **sign** changes
- The coordinates (x, y) become (x, -y)
- Horizontal asymptotes change
  - y = k becomes y = -k
- Vertical asymptotes stay the same



The diagram below shows the graph of y = f(x).




a) Sketch the graph of y = -f(x).




# y = -f(x) reflection in x-axis

- A becomes (-1,-5)
- B becomes (3, 3)



- Sketch the graph of y = f(-x). b)
  - y = f(-x) reflection in y-axis

  - A becomes (1,5)
    B becomes (-3,-3)






### 2.5.3 Stretches of Graphs

### Stretches of Graphs

#### What are stretches of graphs?

- When you alter a function in certain ways, the effects on the graph of the function can be described by **geometrical transformations**
- For a stretch:
  - the graph is **stretched** about one of the coordinate axes by a scale factor
    - Its size changes
  - the orientation of the graph remains unchanged
- A particular stretch is specified by a **coordinate axis** and a **scale factor**:
  - The distance between a point on the graph and the specified coordinate axis is multiplied by the constant scale factor
  - The graph is stretched in the direction which is parallel to the other coordinate axis
  - For scale factors bigger than 1
    - the points on the graph get further away from the specified coordinate axis
  - For scale factors **between 0 and 1** 
    - the points on the graph get closer to the specified coordinate axis
    - This is also sometimes called a compression but in your exam you must use the term stretch with the appropriate scale factor



#### What effects do horizontal stretches have on the graphs and functions?

• A horizontal stretch of the graph y = f(x) by a scale factor q centred about the y-axis is represented by



$$y = f\left(\frac{x}{q}\right)$$

- The x-coordinates change
  - They are **divided** by q
- The y-coordinates stay the same
- The coordinates (x, y) become (qx, y)
- Horizontal asymptotes stay the same
- Vertical asymptotes change
  - x = k becomes x = qk



#### What effects do vertical stretches have on the graphs and functions?


• A vertical stretch of the graph y = f(x) by a scale factor p centred about the x-axis is represented by

$$\frac{y}{p} = f(x)$$

- This is often rearranged to y = pf(x)
- The x-coordinates stay the same
- The y-coordinates change
  - They are multiplied by p
- The coordinates (x, y) become (x, py)
- Horizontal asymptotes change
  - y = k becomes y = pk
- Vertical asymptotes stay the same



The diagram below shows the graph of y = f(x).



a) Sketch the graph of y = 2f(x).




# y=kf(x) vertical stretch scale factor k

Stretch y=f(x) vertically scale factor 2

A becomes (-1,10)

B becomes (3,-6)




b) Sketch the graph of y = f(2x).

y = f(kx) horizontal stretch scale factor  $\frac{1}{k}$ 

Stretch y=f(x) horizontally scale factor  $\frac{1}{2}$ 

A becomes  $\left(-\frac{1}{2},5\right)$ 

B becomes  $(\frac{3}{\lambda}, -3)$ 





### 2.5.4 Composite Transformations of Graphs

### **Composite Transformations of Graphs**

#### What transformations do I need to know?

$$y = f(x+k) \text{ is horizontal translation by vector} \begin{pmatrix} -k \\ 0 \end{pmatrix}$$

- If k is **positive** then the graph moves **left**
- If k is **negative** then the graph moves **right**

$$y = f(x) + k \text{ is } \mathbf{vertical translation by vector} \begin{pmatrix} 0 \\ k \end{pmatrix}$$


- If k is **positive** then the graph moves **up**
- If k is **negative** then the graph moves **down**
- y = f(kx) is a **horizontal stretch** by scale factor  $\frac{1}{k}$  centred about the y-axis
  - If k > 1 then the graph gets closer to the y-axis
  - If **0 < k < 1** then the graph gets **further** from the y-axis
- y = kf(x) is a **vertical stretch** by scale factor k centred about the x-axis
  - If k > 1 then the graph gets further from the x-axis
  - If **0 < k < 1** then the graph gets **closer** to the x-axis
- y = f(-x) is a **horizontal reflection** about the y-axis
  - A horizontal reflection can be viewed as a special case of a horizontal stretch
- y = -f(x) is a **vertical reflection** about the x-axis
  - A vertical reflection can be viewed as a special case of a vertical stretch

#### How do horizontal and vertical transformations affect each other?

- Horizontal and vertical transformations are independent of each other
  - The horizontal transformations involved will need to be applied in their correct order
  - The vertical transformations involved will need to be applied in their correct order
- Suppose there are two horizontal transformation H<sub>1</sub>then H<sub>2</sub> and two vertical transformations V<sub>1</sub>then
   V<sub>2</sub> then they can be applied in the following orders:
  - Horizontal then vertical:
    - H<sub>1</sub>H<sub>2</sub>V<sub>1</sub>V<sub>2</sub>
  - Vertical then horizontal:
    - $V_1V_2H_1H_2$
  - Mixed up (provided that H<sub>1</sub> comes before H<sub>2</sub> and V<sub>1</sub> comes before V2):
    - H<sub>1</sub>V<sub>1</sub>H<sub>2</sub>V<sub>2</sub>
    - H<sub>1</sub>V<sub>1</sub>V<sub>2</sub>H<sub>2</sub>
    - V<sub>1</sub>H<sub>1</sub>V<sub>2</sub>H<sub>2</sub>
    - V<sub>1</sub>H<sub>1</sub>H<sub>2</sub>V<sub>2</sub>



The diagram below shows the graph of y = f(x).



Sketch the graph of 
$$y = \frac{1}{2} f\left(\frac{x}{2}\right)$$
.




A vertical and horizontal transformation can be done in any order

 $y = \frac{1}{2}f(x)$ : vertical stretch scale factor  $\frac{1}{2}$  $y = f(\frac{x}{2})$ : horizontal stretch scale factor 2

A becomes  $\left(-2, \frac{5}{2}\right)$ 

B becomes  $(6, -\frac{3}{2})$ 






## Composite Vertical Transformations af(x)+b

#### How do I deal with multiple vertical transformations?

- Order matters when you have more than one vertical transformations
- If you are asked to find the equation then **build up the equation** by looking at the transformations in order
  - A **vertical stretch** by scale factor a followed by a **translation** of  $\begin{pmatrix} 0 \\ b \end{pmatrix}$ 
    - Stretch: y = af(x)
    - Then translation: y = [af(x)] + b
    - Final equation: y = af(x) + b
  - A **translation** of  $\begin{pmatrix} 0 \\ b \end{pmatrix}$  followed by a **vertical stretch** by scale factor a
    - Translation: y = f(x) + b
    - Then stretch: y = a[f(x) + b]
    - Final equation: y = af(x) + ab
- If you are asked to determine the order
  - The order of vertical transformations follows the order of operations
  - First write the equation in the form y = af(x) + b
    - First stretch vertically by scale factor a
    - If a is negative then the **reflection and stretch** can be **done in any order**
    - Then translate by  $\begin{pmatrix} 0 \\ b \end{pmatrix}$



The diagram below shows the graph of y = f(x).



Sketch the graph of y = 3f(x) - 2.




The order vertical transformations follows the order of operations

y = 3f(x): Vertical stretch scale factor 3

y = f(x) - 2: Translate  $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$ 

A becomes (-1, 13)

B becomes (3,-11)






## Composite Horizontal Transformations f(ax+b)

#### How do I deal with multiple horizontal transformations?

- Order matters when you have more than one horizontal transformations
- If you are asked to find the equation then **build up the equation** by looking at the transformations in order
  - A horizontal stretch by scale factor  $\frac{1}{a}$  followed by a translation of  $\begin{pmatrix} -b \\ 0 \end{pmatrix}$ 
    - Stretch: y = f(ax)
    - Then translation: y = f(a(x+b))
    - Final equation: y = f(ax + ab)
  - A translation of  $\begin{pmatrix} -b \\ 0 \end{pmatrix}$  followed by a horizontal stretch by scale factor  $\frac{1}{a}$ 
    - Translation: y = f(x + b)
    - Then stretch: y = f((ax) + b)
    - Final equation: y = f(ax + b)
- If you are asked to determine the order
  - First write the equation in the form y = f(ax + b)
  - The order of horizontal transformations is the reverse of the order of operations
    - First translate by  $\begin{pmatrix} -b \\ 0 \end{pmatrix}$
    - Then stretch by scale factor  $\frac{1}{a}$
    - If a is negative then the reflection and stretch can be done in any order



The diagram below shows the graph of y = f(x).



Sketch the graph of y = f(2x - 1).



The order of horizontal transformations is the reverse of the order of operations

y = f(x-1): Translate  $\binom{1}{0}$ y = f(2x): Horizontal stretch scale factor  $\frac{1}{2}$ 

A becomes (0,5)

B becomes (2,-3)

