

Boost your performance and confidence with these topic-based exam questions


Practice questions created by actual examiners and assessment experts

Detailed mark schemes

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2.8 Photosynthesis



IB Biology - Revision Notes


www.exampaperspractice.co.uk




2.8.1 Photosynthesis

Photosynthesis Defined

- Simple, inorganic compounds are converted into complex organic ones by photosynthesis
 - The energy required is provided by **light**
- Photosynthesis occurs in autotrophic organisms such as plants, algae and cyanobacteria
 - H₂O and CO₂ are the **raw materials**
- Photosynthesis is a form of energy conversion, from light energy to chemical energy, stored in biomass
- Energy is stored within the bonds of these organic compounds
- Photosynthesis can be thought of as the **exact reverse of respiration**
 - Respiration is the process by which **energy is** released from organic molecules in living cells



The basic equation of photosynthesis as it takes place in a leaf



The chemical equation for photosynthesis



Remember, energy is never created or destroyed; it is only ever converted from one form to another!



Visible Light Wavelengths

- Chloroplasts contain **pigments** in order to **absorb light**
- Pigments are coloured, which means they absorb some wavelengths (or colours) of the white light that the Sun radiates
 - The remaining light is **reflected**, giving the pigment its colour
- Chloroplasts contain several different photosynthetic pigments, so that they can absorb multiple different wavelengths of light
 - The main photosynthetic pigment is **chlorophyll**
- Violet light has the shortest wavelength of light in the visible spectrum (around 400nm)
- Red light has the longest wavelength of light in the visible spectrum (around 700nm)
- Green light has a wavelength in the middle of this range (around 550nm)
- The absorption of light varies with wavelength, as does the rate of photosynthesis that a plant can carry out
- When plants are exposed to light of a specific wavelength, the rate of photosynthesis can be measured as well as the absorbance (the % of the light that is absorbed by the plants)
- There are peaks in both plots at the **blue** and **red** ends of the spectrum, where photosynthesis can occur
- There are troughs in both plots for green light, which is not absorbed and so cannot provide energy for photosynthesis



You don't have to memorise the wavelengths of different colours of light, but you need to know that visible light has a wavelength of between 400 and 700 nanometres (nm).

© 2024 Exam Papers Practice



Chlorophyll

- Plant cells contain **chloroplasts** which are the site of photosynthesis
- The main photosynthetic pigment is **chlorophyll**
- Chlorophyll absorbs red and blue light most effectively and reflects green light more than other colours
 - Chlorophyll appears green because it absorbs red and blue light
 - The green light is reflected away and so leaves appear green to the eye
 - This explains why the majority of plants are green (with variations in the shades of green that we can see)
- Red and blue light provides the energy needed for photosynthesis
- Chlorophyll exists in **two main forms**, **a** and **b**
- There are two groups of pigments: primary pigments known as chlorophylls and accessory pigments known as carotenoids
- Chlorophylls absorb wavelengths in the blue-violet and red regions of the light spectrum
- Carotenoids absorb wavelengths of light mainly in the blue-violet region of the spectrum
- The combination of pigments maximises the amount of white light energy that can be captured



Remember - chlorophyll is not the only photosynthetic pigment, others exist to maximise light energy absorption.

Photolysis of Water

- Oxygen is produced in photosynthesis from the photolysis of water
- Copyright Photo means 'with light'
- © 2024 Exar**Lysis** emeanstibreaking apart'
 - Water is broken apart using light energy; this is called photolysis
 - This releases **electrons** (e⁻), **protons** (H⁺) and the waste product, **oxygen gas**

$$2H_2O \rightarrow 4e^- + 4H^+ + O_2$$

- Whilst oxygen is a waste product, the electrons and protons play a crucial role in the further reactions of photosynthesis
 - Though oxygen is a waste product, in practice, a plant will use some of the oxygen it produces in photosynthesis for its own respiration (during the day)



2.8.2 Photosynthesis Continued

Effects of Photosynthesis

- Changes to the Earth's atmosphere, oceans and rock deposition occur due to photosynthesis
- The first life forms emerged around 4 billion years ago
 - At the time, there was no oxygen in the atmosphere
- About 3.5 billion years ago photosynthetic prokaryotes became the first organisms to carry out photosynthesis
 - This began the release of oxygen into the atmosphere
- Millions of years later algae and plants evolved and also carried out photosynthesis
- Around 2.2 billion years ago, the oxygen concentration in the atmosphere reached 2%
 - This is known as the Great Oxidation Event
- Other changes to the Earth occurred due to photosynthesis
 - Minerals in the oceans were oxidised
 - Photosynthetic bacteria released oxygen into the ocean
 - When dissolved iron was oxidised it formed iron oxide which is a red precipitate that lies on the sea bed
 - Over time a distinctive rock formation was produced the banded iron formation. Layers of red iron oxide alternate with other mineral oxides
 - Banded iron formations are the most important source of iron ores (and consequently our supply of steel)
 - Methane and CO₂ levels in the air fell, which resulted in an Ice Age
 - Because methane and CO₂ are important greenhouse gases
- ractice
- By 600 million years ago, life had evolved into large multicellular organisms, many of which were photosynthetic (plants)
 - This pushed the oxygen concentration of the air up to 20%, peaking at 35% 300 million years ago
 - This contributed to the large size of the animals that roamed the Earth at that time
 - The current atmospheric oxygen level is around 21%, due to **increased human activity** eg. burning of fossil fuels, deforestation which remove oxygen from the atmosphere



Energy Requirements

- Chemical reactions can be **exothermic** or **endothermic**
- Photosynthesis is an example of an endothermic reaction and an anabolic reaction, where the required energy input is in the form of light energy
- Energy is needed to produce carbohydrates and other carbon compounds from carbon dioxide
- The energy is not lost it is stored in chemical form in the carbohydrates that are produced

Limiting Factors

Temperature, light intensity and carbon dioxide concentration are possible limiting factors on the rate of photosynthesis

- Each of these factors can limit the rate of photosynthesis when they are below the optimal level
 - Temperature
 - Light intensity
 - Carbon dioxide concentration
- These are known as the **limiting factors** of photosynthesis
 - A limiting factor is a variable that holds back the rate of a chemical reaction
 - If that variable is increased, the reaction rate also increases
- Under any set of conditions, only one of these factors will be limiting the rate of photosynthesis
 - At night, light intensity will be very low, so that is the limiting factor
 - On a cold, sunny day, temperature will be the limiting factor
 - An increase in the light intensity will not increase the rate of photosynthesis because the temperature is, at that point, the limiting factor

Analogy - Limiting Factors

Imagine you're on a group hike and you have to stick together with your two teammates. You are not allowed to finish the hike separately. If your legs are feeling weak, you will be walking slowly. The other two hikers will have to slow down to walk at your speed; the team progresses at the speed of the slowest member. However, after lunch, you may be feeling strengthened by the food and your legs feel good; this time, it's your teammate who is the slowest, so has taken over from you as the limiting factor. Your team only finishes when all three of you finish, and that determines your overall rate of progress.



When writing about limiting factors, it's important to mention '**light intensity**', not just 'light'. Other aspects of light such as wavelength can play a role so it's important to specific about intensity.



2.8.3 Skills: Photosynthesis


Absorption & Action Spectrums

- An **absorption spectrum** (for a particular pigment) shows how much light of different wavelengths is absorbed by the pigment
- An **action spectrum** shows how each wavelength of light affects the rate of photosynthesis that it can power
- These are two graphs that use the wavelength of light on the x-axis
 - Violet light is the lowest wavelength of the visible spectrum, at 400 nanometres (nm)
 - Red light is the highest wavelength of the visible spectrum, at 700 nanometres (nm)

Drawing an absorption spectrum for chlorophyll

Step 1: Draw and label the axes


- Draw an x-axis
- Label the axis wavelength
- Add the units / nm
- Make 400 the smallest value and 700 the largest value
 - Label 500 and 600 nm on the x-axis
- Draw a y-axis
- Labelit % absorption
- Make 0 the lowest value and 100 the highest value
 - No units are required because the y-axis is showing a percentage scale





Step 2: Draw the Plot

- There should be two absorbance peaks
 - One peak at either end, in the blue and red areas of the spectrum
 - And a **trough** in the middle, which represents green light
 - As below, with a smooth curve




© 2024 Exam P An absorbance spectrum for photosynthesis (colour range labels are not required)

Drawing an action spectrum for photosynthesis


Step 1: Draw and label the axes

- Draw an x-axis
- Label the axis wavelength
- Add the units / nm
- Make 400 the smallest value and 700 the largest value
 - Label 500 and 600 nm on the x-axis
- Draw a y-axis
- Labelit Rate of photosynthesis / % of maximum rate
- Make 0 the lowest value and 100 the highest value
 - No units are required because the y-axis is showing a percentage scale





- One peak at either end, in the blue and red areas of the spectrum
- And a **trough** in the middle, which represents green light
- As below, with a smooth curve



Step 2: Sketch the Curve. An absorbance spectrum for photosynthesis (colour range labels are not required)



Exam Tip

Remember – the pigments themselves have a distinctive colour. This is different from the colours of light that they *absorb*. Key points to remember:

- 1. Label 400 700nm on the x-axis, in 100nm increments
- 2. Use a % scale on the y-axis
- 3. Smooth curve
- 4. Peaks at either end
- 5. Trough in the middle for green light

Investigating Photosynthesis

- An aquatic plant such as Elodea or Camboba is a good choice for investigating photosynthesis in
 plants, because the rate of photosynthesis can be measured by counting oxygen bubbles that
 come off a cutting of this plant
 - Oxygen output from terrestrial plants (that grow on land) would not be observable


NOS: Experimental design - controlling relevant variables in photosynthesis experiments is essential

- When designing an experiment it is crucial that all variables (apart from the independent and dependent variables being investigated) are controlled
 - The independent variable is the factor that is **deliberately manipulated** between a specific range throughout the experiment
 - The dependent variable is the factor that is measured during the experiment (to see if it is
 affected by the changes to the independent variable)
- Other variables must be controlled so that it can be said the independent variable is the only Copyright factor affecting the dependent variable during the experiment
- © 202 Changes in light intensity, carbon dioxide concentration and temperature are all limiting factors that affect the rate of photosynthesis and can be altered experimentally to measure the effect on the rate of photosynthesis

Effect of light intensity - experimental design

- Basic Experimental Setup
 - Aquatic plant cutting in water
 - Powdered sodium hydrogencarbonate (NaHCO₃)
 - Glass funnel
 - Boiling tube
 - Lampforillumination
 - Glass tank filled with water





Measuring the Effect of Light Intensity on the Rate of Photosynthesis in Pondweed

Research Question

Does the rate of photosynthesis (number of bubbles released per min) of *Elodea* increase as the light intensity increases?

© 2 Method Papers Practice

- Place a piece of aquatic plant (*Elodea* or *Cabomba* are often used), into a beaker of water
- Use a light a set distance from the plant
- Record the number of bubbles observed in three minutes
- Repeat steps for different distances

Improvements

- Use a gas syringe to collect and measure the volume of gas produced
- For **reliability** of data, **repeat** the experiment at least twice for each distance and calculate the mean number of bubbles
- Use of a data logger to measure results continuously




Variables to Be Controlled

- Temperature
 - The glass tank filled with water absorbs any heat that is emitted from the lamp
 - Modern LED bulbs can be used as they give off less heat than filament bulbs
- CO₂ concentration
 - The water used around the plant is first boiled and re-cooled to remove any dissolved carbon dioxide
 - A set mass of sodium hydrogencarbonate is added to the water that surrounds the plant
 - To make the concentration approx. 0.1 mol dm⁻³, which is not a limiting concentration


Results

- A graph of the number of bubbles produced per minute against the distance between the lamp and the plant used can be drawn to see the pattern or trend
 - Distance between the lamp and the plant is linked to the light intensity



A graph of distance from the lamp against number of bubbles per minute






The effect of light intensity on the rate of photosynthesis

Carbon dioxide concentration

- The same basic experimental setup can be used, but with varying use of the following variables
- Copy ig Start with boiled and re-cooled water as before
- © 20.24 Add successive masses of sodium hydrogencarbonate to increase the concentration in increments of 0.01 mol dm⁻³, and record the rate of photosynthesis in bubbles minute⁻¹
 - **Keep the temperature constant** at 25°C using a water bath, monitoring with a thermometer in the water surrounding the aquatic plant
 - Keep the light intensity constant by keeping the lamp a fixed distance from the plant






The effect of carbon dioxide concentration on the rate of photosynthesis

Temperature

- The same basic experimental setup can be used, but with varying use of the following variables
- Copy in glStart with boiled and re-cooled water as before, with **sodium hydrogencarbonate at a fixed** © 2024 **concentration** of 0-1 mol dm⁻³, and record the rate of photosynthesis in bubbles minute⁻¹
 - Vary the temperature from 5°C to 50°C using water baths, monitoring with a thermometer in the water surrounding the plant
 - Keep the light intensity constant by keeping the lamp a fixed distance from the plant
 - The rate will tail off after around 40°C due to the denaturation of important proteins involved in photosynthesis





The effect of temperature on the rate of photosynthesis

Papers Practice

CopyriThe key to this part of the spec is to appreciate how an experimental investigation can be 202 controlled so that any effects we observe **are directly due to the one variable** that we are deliberately changing.



2.8.4 Skills: Separating Photosynthetic Pigments

Practical 4: Separation of Photosynthetic Pigments

Separation of photosynthetic pigments by chromatography

- Plants contain several different photosynthetic pigments, which absorb different wavelengths of light
- There are two groups of pigments: chlorophylls and carotenoids
- Carotenoids surround the chlorophyll and absorb both similar and different wavelengths of light to chlorophyll
 - This expands the range of wavelengths that can be absorbed from light for use in photosynthesis

Chloroplast Pigments Table


Pigment group	Name of pigment	Colour of pigment
Chlorophylls	Chlorophyll a Chlorophyll b	Yellow-green Blue-green
Carotenoids	β carotene Xanthophyll	Orange Yellow

Copyright

© 2024 Exam Paners Practice

- Chlorophylls absorb wavelengths in the blue-violet and red regions of the light spectrum
 - They reflect green light, causing plants to appear green
- Carotenoids absorb wavelengths of light mainly in the blue-violet region of the spectrum





Exam Pages Practice

600

700

Copy**Chlorophyll and carotenoids absorb light across the visible light spectrum to use in the light-dependent**© 2024 Exam Papers Practice reaction of photosynthesis

Chromatography

400

- Chromatography is an experimental technique that is used to separate mixtures
 - Different components within the mixture travel through the material at different speeds
 - This causes the different components to **separate**

500

- A retardation factor (R_f value) can be calculated for each component of the mixture
- Two of the most common techniques for separating these photosynthetic pigments are:
 - Paper chromatography the mixture of pigments is passed through paper (cellulose)
 - Thin-layer chromatography (TLC) the mixture of pigments is passed through a thin layer of adsorbent (eg. silica gel), through which the mixture travels faster and separates more distinctly



 Paper chromatography can be used to separate photosynthetic pigments although TLC gives better results

Apparatus

- Leaf sample
- Distilled water
- Pestle and mortar
- Filterpaper
- Capillary tube
- Chromatographysolvent
- Propanone
- Pencil
- Ruler

Method

- Draw a straight line in pencil approximately 1cm above the bottom of the filter paper being used
 - Do not use a pen as the ink will separate into pigments within the experiment and obscure the
 results
- Cut a section of leaf and place it in a mortar
 - It is important to choose a healthy leaf that has been in direct sunlight so you can be sure it contains many active photosynthetic cells
- Add 20 drops of propanone and use the pestle to grind up the leaf sample and release the pigments
 - Propanone is an organic solvent and therefore fats, such as the lipid membrane, dissolve in it
 - The combination of propanone and mechanical pressure breaks down the cell and chloroplasts to release the pigments

Copyright

- © 202 Extract some of the pigment using a capillary tube and spot it onto the centre of the pencil line you have drawn
 - Suspend the paper in the chromatography solvent so that the level of the solvent is below the pencil line and leave the paper until the solvent has reached the top of the paper
 - The mixture is dissolved in the solvent (called the mobile phase) and the dissolved mixture then passes through a static material (called the stationary phase)
 - Remove the paper from the solvent and draw a pencil line marking where the solvent moved up to
 - The pigment should have separated out and there should be different spots on the paper at different heights above the pencil line, these are the separate pigments
 - Calculate the Rf value for each spot




R_f value = $\frac{\text{distance travelled by component (pigment)}}{\text{distance travelled by the solvent}}$


Always measure to the centre of each spot

Results

- Chromatography can be used to **separate and identify chloroplast pigments** that have been extracted from a leaf as each pigment will have a unique R_f value
- The R_f value demonstrates how far a dissolved pigment travels through the stationary phase
 - Molecules with a higher affinity to the stationary phase, such as large molecules, will travel slower and therefore have a smaller R_f value
 - Molecules that are more soluble in the mobile phase will travel faster and therefore have a larger R_f value
- Although specific R_f values depend on the solvent that is being used, in general:
 - Carotenoids have the highest R_f values (usually close to 1)
 - Chlorophyll bhas a much lower Rf value
 - Chlorophyll a has an R_f value somewhere between those of carotenoids and chlorophyll b
 - Small R_f values indicate the pigment is less soluble and/or larger in size







Paper chromatography is used to separate photosynthetic pigments. These pigments can be identified by their R_f values. In this example, a line of the mixture (rather than a spot) is added to the paper.

Limitations

- Copyright Paper chromatography is not as specific as other chromatography techniques
 - It is sufficient to separate and distinguish different pigments and to calculate their R_f value
 - Chromatography does not give data on the amount of each pigment present or the wavelengths that they absorb
 - Colorimetry can be used to calculate these values

Exam Tip

Remember – the pigments themselves have colour (as described in the table). This is different from the colours of light that they *absorb*. You don't have to remember specific R_f values, just know that they differ between each type of pigment.