



Cambridge International AS & A Level

MATHEMATICS

9709/02

Paper 2 Pure Mathematics 2

For examination from 2020

MARK SCHEME

Maximum Mark: 50

Specimen

This document has **10** pages. Blank pages are indicated.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

<p>GENERIC MARKING PRINCIPLE 1:</p> <p>Marks must be awarded in line with:</p> <ul style="list-style-type: none"> ● the specific content of the mark scheme or the generic level descriptors for the question ● the specific skills defined in the mark scheme or in the generic level descriptors for the question ● the standard of response required by a candidate as exemplified by the standardisation scripts.
<p>GENERIC MARKING PRINCIPLE 2:</p> <p>Marks awarded are always whole marks (not half marks, or other fractions).</p>
<p>GENERIC MARKING PRINCIPLE 3:</p> <p>Marks must be awarded positively:</p> <ul style="list-style-type: none"> ● marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate ● marks are awarded when candidates clearly demonstrate what they know and can do ● marks are not deducted for errors ● marks are not deducted for omissions ● answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.
<p>GENERIC MARKING PRINCIPLE 4:</p> <p>Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.</p>
<p>GENERIC MARKING PRINCIPLE 5:</p> <p>Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).</p>

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mark Scheme Notes

Marks are of the following three types.

- M** Method mark, given for a valid method applied to the problem. Method marks can still be given even if there are numerical errors, algebraic slips or errors in units. However the method must be applied to the specific problem, e.g. by substituting the relevant quantities into a formula. Correct use of a formula without the formula being quoted earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, given for an accurate answer or accurate intermediate step following a correct method. Accuracy marks cannot be given unless the relevant method mark has also been given.
- B** Mark for a correct statement or step.
- DM or DB** M marks and B marks are generally independent of each other. The notation DM or DB means a particular M or B mark is dependent on an earlier M or B mark (indicated by *). When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT below).
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures (sf) or would be correct to 3 sf if rounded (1 decimal point (dp) for angles in degrees). As stated above, an A or B mark is not given if a correct numerical answer is obtained from incorrect working.
 - Common alternative solutions are shown in the Answer column as: **EITHER** Solution 1 **OR** Solution 2 **OR** Solution 3 ...? Round brackets appear in the Partial Marks column around the marks for each alternative solution.
 - Square brackets [] around text show extra information not needed for the mark to be awarded.
 - The total number of marks available for each question is shown at the bottom of the Marks column in bold type.

The following abbreviations may be used in a mark scheme.

- AG** Answer given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid).
- CAO** Correct answer only (emphasising that no 'follow through' from an error is allowed).
- CWO** Correct working only
- FT** Follow through after error (see Mark Scheme Notes for further details).
- ISW** Ignore subsequent working
- OE** Or equivalent form
- SC** Special case
- SOI** Seen or implied

Question	Answer	Marks	Partial Marks	Guidance
1(a)	Substitute $x = -1$ and equate to zero	1	M1	
	Obtain answer $a = 7$	1	A1	
		2		
1(b)	Substitute $x = -3$ and evaluate expression	1	M1	
	Obtain answer 18	1	A1	
		2		
Question		Marks	Partial Marks	Guidance
2	Use $\sin 2\theta = 2 \sin \theta \cos \theta$	1	B1	
	Simplify to obtain form $c_1 \sin^2 \theta = c_2$ or equivalent	1	M1	
	Find at least one value of θ from equation of form $\sin \theta = k$	1	M1	
	Obtain 35.3° and 144.7°	1	A1	And no others between 0° and 180° . Unsupported answer receives 0 marks.
		4		

Question	Answer	Marks	Partial Marks	Guidance
3(a)(i)	Draw two V-shaped graphs, one with vertex on negative x -axis, the other with vertex on positive x -axis	1	M1	
	Show graphs in correct relation to each other, with lines (roughly) parallel as appropriate	1	A1	
		2		
3(a)(ii)	State coordinates $(-2a, 0)$, $(\frac{3}{2}a, 0)$, $(0, 4a)$, $(0, 3a)$	1	B1	Accept values inserted at correct points on axes
3(b)	Square both sides to obtain linear equation or inequality and attempt solution OR solve linear equation $2x + 4a = -(2x - 3a)$ OR equivalent OR use symmetry of diagram to determine x -coordinate of point of intersection	1	M1	
	Obtain value $-\frac{1}{4}a$ and no other	1	A1	
	State answer $x > -\frac{1}{4}a$	1	A1	
		3		

Question	Answer	Marks	Partial Marks	Guidance
4(a)	Recognise quadratic equation in 5^x and attempt solution for 5^x	1	M1	
	Obtain $5^x = 3$ only	1	A1	
	Use logarithms to solve equation of the form $5^x = k$ where $k > 0$	1	M1	
	Obtain 0.683	1	A1	
4(b)		4		
	Use $2 \ln x = \ln x^2$	1	B1	
	Use law for addition or subtraction of logarithms	1	M1*	
	Arrange equation not involving logarithms to the form $y = \dots$	1	DM1	
	Obtain $y = \frac{5}{x^2 - 1}$	1	A1	Or equivalent explicit $y = \dots$ form
		4		

Question	Answer	Marks	Partial Marks	Guidance
5(a)	Attempt use of quotient rule or equivalent	1	M1	
	Obtain $\frac{2(x+2)\cos 2x - \sin 2x}{(x+2)^2}$ or equivalent	1	A1	
	Equate numerator to zero and attempt rearrangement	1	M1	
	Confirm given result $\tan 2x = 2x + 4$	1	A1	AG Showing necessary detail
		4		
5(b)	Consider sign of $\tan 2x - 2x - 4$ for 0.6 and 0.7 or equivalent	1	M1	
	Obtain -2.63 and 0.40 or equivalent and justify conclusion	1	A1	
		2		
5(c)	Use iteration process correctly at least once	1	M1	
	Obtain final answer 0.694	1	A1	
	Show sufficient iterations to 5 decimal places to justify answer OR show a sign change in the interval (0.6935, 0.6945)	1	A1	
		3		

Question	Answer	Marks	Partial Marks	Guidance
6(a)	Use product rule to differentiate y	1	M1	
	Obtain $\frac{dy}{dt} = 4e^t + 4te^t$ or equivalent	1	A1	
	Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$	1	M1	
	Obtain given answer $\frac{dy}{dx} = \frac{2(t+1)}{e^t}$ correctly	1	A1	AG
		4		
6(b)	Substitute $t = 0$ to evaluate derivative and find coordinates of point	1	B1	
	Obtain $\frac{dy}{dx} = 2$ and coordinates $(1, 0)$	1	B1	
	Form equation of normal at their point, using negative reciprocal of their $\frac{dy}{dx}$	1	M1	
	State correct equation of normal $y = -\frac{1}{2}x + \frac{1}{2}$ or equivalent	1	A1	
		4		

Question	Answer	Marks	Partial Marks	Guidance
7(a)	Replace $\tan^2 x$ by $\sec^2 x - 1$	1	B1	
	Express $\cos^2 x$ in the form $\pm \frac{1}{2} \pm \frac{1}{2} \cos 2x$	1	M1	
	Obtain given answer $\sec^2 x + \frac{1}{2} \cos 2x - \frac{1}{2}$ correctly	1	A1	AG Showing necessary detail
	Attempt integration of expression	1	M1	
	Obtain $\tan x + \frac{1}{4} \sin 2x - \frac{1}{2}x$	1	A1	
	Use limits correctly for integral involving at least $\tan x$ and $\sin 2x$	1	M1	
	Obtain $\frac{5}{4} - \frac{1}{8}\pi$ or exact equivalent	1	A1	
		7		
7(b)	State or imply volume is $\int \pi(\tan x + \cos x)^2 dx$	1	B1	Presence of π implied by its appearance later if not shown initially
	Attempt expansion and simplification	1	M1	
	Integrate to obtain one additional term of form $k \cos x$	1	M1	
	Obtain $\pi(\frac{5}{4} - \frac{1}{8}\pi) + \pi(2 - \sqrt{2})$ or equivalent	1	A1	
			4	

BLANK PAGE