



# 5.1 Differentiation

# Contents

- \* 5.1.1 Introduction to Differentiation
- \* 5.1.2 Applications of Differentiation



### 5.1.1 Introduction to Differentiation

### Introduction to Derivatives

Before introducing a derivative, an understanding of a limit is helpful

#### What is a limit?

- The **limit** of a **function** is the value a function (of *X*) approaches as *X* approaches a particular value from either side
  - Limits are of interest when the function is undefined at a particular value
  - For example, the function  $f(x) = \frac{x^4 1}{x 1}$  will approach a limit as x approaches 1 from both

below and above but is undefined at x = 1 as this would involve dividing by zero

### What might I be asked about limits?

- You may be asked to predict or estimate limits from a table of function values or from the graph of v = f(x)
- You may be asked to use your GDC to plot the graph and use values from it to estimate a limit


#### What is a derivative?

- Calculus is about rates of change
  - the way a car's position on a road changes is its speed (velocity)
  - the way the car's speed changes is its acceleration
- The gradient (rate of change) of a (non-linear) function varies with X
- The derivative of a function is a function that relates the gradient to the value of X
- The derivative is also called the gradient function

#### How are limits and derivatives linked?

- Consider the point P on the graph of y = f(x) as shown below
  - $[PQ_i]$  is a series of chords





- The gradient of the function f(x) at the point P is equal to the gradient of the tangent at point P
- The gradient of the tangent at point P is the limit of the gradient of the chords  $[PQ_i]$  as point Q

'slides' down the curve and gets ever closer to point P

- The gradient of the function changes as X changes
- The derivative is the function that calculates the gradient from the value X

#### What is the notation for derivatives?

• For the function y = f(x), the **derivative**, with respect to x, would be written as

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f'(x)$$

Different variables may be used

e.g. If 
$$V = f(s)$$
 then  $\frac{\mathrm{d}V}{\mathrm{d}s} = f'(s)$ 



### Worked example

The graph of y = f(x) where  $f(x) = x^3 - 2$  passes through the points P(2, 6), A(2.3, 10.167), B(2.1, 7.261) and C(2.05, 6.615125).

a) Find the gradient of the chords [PA], [PB] and [PC].

Gradient of a line (chord) is  $\frac{42-41}{x_2-x_1}$ [PA]:  $\frac{10\cdot167-6}{2\cdot3-2} = 13\cdot89$   $2\cdot3-2$ [PB]:  $\frac{7\cdot261-6}{2\cdot1-2} = 12\cdot61$ [PC]:  $\frac{6\cdot615125-6}{2\cdot05-2} = 12\cdot3$ [PC]:  $\frac{6\cdot615125-6}{2\cdot05-2} = 12\cdot3$ [PC] 12\cdot3025

b) Estimate the gradient of the tangent to the curve at the point P.

There will be a limit the gradient of the chord reaches as the difference in the x-coordinates approaches zero.

Estimate of gradient of tangent at x=2 is 12



# Differentiating Powers of x

#### What is differentiation?

• **Differentiation** is the process of finding an expression of the **derivative** (gradient function) from the expression of a function

#### How do I differentiate powers of x?

- **Powers** of *X* are **differentiated** according to the following formula:
  - If  $f(x) = x^n$  then  $f'(x) = nx^{n-1}$  where  $n \in \mathbb{Q}$
  - This is given in the **formula booklet**
- If the power of X is multiplied by a constant then the derivative is also multiplied by that constant
  - If  $f(x) = ax^n$  then  $f'(x) = anx^{n-1}$  where  $n \in \mathbb{Q}$  and a is a constant
- The alternative notation (to f'(x)) is to use  $\frac{dy}{dx}$ 
  - If  $y = ax^n$  then  $\frac{dy}{dx} = anx^{n-1}$

e.g. If 
$$y = -4x^{\frac{1}{2}}$$
 then  $\frac{dy}{dx} = -4 \times \frac{1}{2} \times x^{\frac{1}{2}-1} = -2x^{-\frac{1}{2}}$ 

- Don't forget these **two** special cases:
  - If f(x) = ax then f'(x) = a

• e.g. If 
$$y = 6x$$
 then  $\frac{dy}{dx} = 6$ 

- If f(x) = a then f'(x) = 0• e.g. If y = 5 then  $\frac{dy}{dx} = 0$
- These allow you to differentiate **linear terms** in *X* and **constants**
- Functions involving **roots** will need to be rewritten as **fractional powers** of *X* first
  - e.g. If  $f(x) = 2\sqrt{x}$  then rewrite as  $f(x) = 2x^{\frac{1}{2}}$  and differentiate
- Functions involving fractions with denominators in terms of X will need to be rewritten as negative powers of X first

• e.g. If 
$$f(x) = \frac{4}{x}$$
 then rewrite as  $f(x) = 4x^{-1}$  and differentiate

How do I differentiate sums and differences of powers of x?




- The formulae for differentiating powers of *X* apply to **all rational** powers so it is possible to differentiate any expression that is a **sum** or **difference** of **powers** of *X* 
  - e.g. If  $f(x) = 5x^4 3x^{\frac{2}{3}} + 4$  then  $f'(x) = 5 \times 4x^{4-1} - 3 \times \frac{2}{3}x^{\frac{2}{3}-1} + 0$  $f'(x) = 20x^3 - 2x^{-\frac{1}{3}}$

.

- **Products** and **quotients cannot** be differentiated in this way so would need **expanding/simplifying** first
  - e.g. If  $f(x) = (2x-3)(x^2-4)$  then expand to  $f(x) = 2x^3 3x^2 8x + 12$  which is a sum/difference of powers of X and can be differentiated







# 5.1.2 Applications of Differentiation


# **Finding Gradients**

### How do I find the gradient of a curve at a point?

- The gradient of a curve at a point is the gradient of the tangent to the curve at that point
- Find the gradient of a curve at a point by substituting the value of X at that point into the curve's derivative function
- For example, if  $f(x) = x^2 + 3x 4$ 
  - then f'(x) = 2x + 3
  - and the gradient of y = f(x) when x = 1 is f'(1) = 2(1) + 3 = 5
  - and the gradient of y = f(x) when x = -2 is f'(-2) = 2(-2) + 3 = -1
- Although your GDC won't find a derivative function for you, it is possible to use your GDC to evaluate


the derivative of a function at a point, using  $\frac{d}{dx}(\Box)_{x=\Box}$ 





(c) Find the gradient of y = f(x) when x = -2.5.








### **Increasing & Decreasing Functions**

#### What are increasing and decreasing functions?

- A function, f(x), is increasing if f'(x) > 0
  - This means the value of the function ('output') increases as X increases
- A function, f(x), is decreasing if f'(x) < 0
  - This means the value of the function ('output') decreases as X increases
- A function, f(x), is stationary if f'(x) = 0



#### How do I find where functions are increasing, decreasing or stationary?

 To identify the intervals on which a function is increasing or decreasing STEP 1
 Find the derivative f'(x)

#### **STEP 2** Solve the inequalities



f'(x) > 0 (for increasing intervals) and/or f'(x) < 0 (for decreasing intervals)

- Most functions are a combination of increasing, decreasing and stationary
  - a range of values of X (interval) is given where a function satisfies each condition
  - e.g. The function  $f(x) = x^2$  has derivative f'(x) = 2x so
    - f(x) is decreasing for x < 0
    - f(x) is stationary at x = 0

• f(x) is increasing for x > 0






$$f(x) = x^2 - x - 2$$

a) Determine whether f(x) is increasing or decreasing at the points where x = 0 and x = 3.

Differentiate  

$$f'(x) = 2x - 1$$
  
At  $x = 0$ ,  $f'(0) = 2x0 - 1 = -1 < 0$  : decreasing  
At  $x = 3$ ,  $f'(3) = 2x3 - 1 = 5 > 0$  : increasing  
 $\therefore$  At  $x = 0$ ,  $f(x)$  is decreasing  
At  $x = 3$ ,  $f(x)$  is increasing

b) Find the values of X for which f(x) is an increasing function.






# **Tangents & Normals**

### What is a tangent?

- At any point on the graph of a (non-linear) **function**, the **tangent** is the straight line that **touches** the graph at a point **without crossing** through it
- Its gradient is given by the derivative function



### How do I find the equation of a tangent?

- To find the equation of a straight line, a point and the gradient are needed
- The gradient, m, of the tangent to the function y = f(x) at  $(x_1, y_1)$  is  $f'(x_1)$
- Therefore find the equation of the tangent to the function y = f(x) at the point  $(x_1, y_1)$  by


substituting the gradient, 
$$f'(x_1)$$
, and point  $(x_1, y_1)$  into  $y - y_1 = m(x - x_1)$ , giving:  
 $y - y_1 = f'(x_1)(x - x_1)$ 

• (You could also substitute into y = mx + c but it is usually quicker to substitute into  $y - y_1 = m(x - x_1)$ )

#### What is a normal?

• At any point on the graph of a (non-linear) function, the **normal** is the straight line that passes through that point and is **perpendicular** to the **tangent** 





### How do I find the equation of a normal?

- The gradient of the normal to the function y = f(x) at  $(x_1, y_1)$  is  $\frac{-1}{f'(x_1)}$
- Therefore find the equation of the normal to the function y = f(x) at the point  $(x_1, y_1)$  by using

$$y - y_1 = \frac{-1}{f'(x_1)}(x - x_1)$$



# Worked example

The function f(x) is defined by

$$f(x) = 2x^4 + \frac{3}{x^2} \qquad x \neq 0$$

a) Find an equation for the tangent to the curve y = f(x) at the point where x = 1, giving your answer in the form y = mx + c.

First find 
$$f'(x)$$
 by differentiating  
 $f(x) = 2x^{14} + 3x^{-2}$  Rewrite as powers of x  
 $f'(x) = 8x^3 - 6x^{-3}$   
For a tangent, "y-y<sub>1</sub> =  $f(a)(x-x_1)$ "  
At  $x = 1$ ,  $y = 2(1)^{14} + \frac{3}{(1)^2} = 5$   
 $f'(1) = 8(1)^3 - \frac{6}{(1)^3} = 2$   
 $(1)^3$   
 $\therefore y - 5 = 2(x - 1)$   
Tangent at  $x = 1$ , is  $y = 2x + 3$ 

b) Find an equation for the normal at the point where x = 1, giving your answer in the form ax + by + d = 0, where a, b and d are integers.



For a normal, "y-y\_1 =  $\frac{-1}{f'(a)}(x-x_1)$ " Using results from part (a):  $y-5 = \frac{-1}{2}(x-1)$   $y = -\frac{1}{2}x + \frac{11}{2}$  2y = -x + 11"Equation of normal is x + 2y - 11 = 0