



**EXAM PAPERS PRACTICE**

Boost your performance and confidence with these topic-based exam questions


Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

## **5.12 Further Limits (inc l'Hôpital's Rule)**



# **IB Maths - Revision Notes**

---

**AA HL**



## 5.12.1 Further Limits

### L'Hôpital's Rule

#### What is L'Hôpital's Rule?

- L'Hôpital's rule is a method involving calculus that allows us to find the value of certain limits
- Specifically, it allows us to attempt to evaluate the limit of a quotient  $\frac{f(x)}{g(x)}$  for which our usual limit evaluation techniques would return one of the **indeterminate forms**  $\frac{0}{0}$  or  $\frac{\pm\infty}{\pm\infty}$ .

#### How do I evaluate a limit using L'Hôpital's Rule?

- STEP 1: Check that the limit of the quotient results in one of the indeterminate forms given above

- I.e., check that  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{f(a)}{g(a)} = \frac{0}{0}$  or  $\frac{\pm\infty}{\pm\infty}$

- STEP 2: Find the derivatives of the numerator and denominator of the quotient

- STEP 3: Check whether the limit  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$  exists

- STEP 4: If that limit does exist, then  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

- STEP 5: If  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \frac{f'(a)}{g'(a)} = \frac{0}{0}$  or  $\frac{\pm\infty}{\pm\infty}$  then you may repeat the process by considering

$$\lim_{x \rightarrow a} \frac{f''(x)}{g''(x)}$$

(and possibly higher order derivatives after that)

- As long as the limits continue giving indeterminate forms you may continue applying L'Hôpital's rule
- Each time this happens find the next set of derivatives and consider the limit again

#### Exam Tip

- Some limits of an indeterminate form can also be evaluated using the **Maclaurin series** for the numerator and denominator
- If an exam question does not specify a method to use, then you are free to use whichever method you prefer

**Worked example**

Use l'Hôpital's rule to evaluate each of the following limits:

a)  $\lim_{x \rightarrow 0} \frac{\sin x}{e^x - 1}$

STEP 1:  $\lim_{x \rightarrow 0} \frac{\overset{f(x)}{\sin x}}{\underset{g(x)}{e^x - 1}} = \frac{\sin(0)}{e^0 - 1} = \frac{0}{0}$  ← indeterminate form

STEP 2:  $\frac{d}{dx}(\sin x) = \cos x$      $\frac{d}{dx}(e^x - 1) = e^x$

STEP 3:  $\lim_{x \rightarrow 0} \frac{\cos x}{e^x} = \frac{\cos(0)}{e^0} = \frac{1}{1} = 1$  ← limit exists

STEP 4:  $\lim_{x \rightarrow 0} \frac{\sin x}{e^x - 1} = 1$

b)  $\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x}$

STEP 1:  $\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x} = \frac{0^3}{-2(0) + \sin 0} = \frac{0}{0}$  ← indeterminate form

STEP 2:  $\frac{d}{dx}(x^3) = 3x^2$      $\frac{d}{dx}(-2x + \sin 2x) = -2 + 2\cos 2x$

STEP 3:  $\lim_{x \rightarrow 0} \frac{3x^2}{-2 + 2\cos 2x} = \frac{3(0)^2}{-2 + 2\cos 0} = \frac{0}{0}$  ← indeterminate form

STEP 4: That limit is still an indeterminate form, so proceed to STEP 5.

STEP 5:  $\lim_{x \rightarrow 0} \frac{6x}{-4\sin 2x} = \frac{6(0)}{-4\sin 0} = \frac{0}{0}$      $\frac{d}{dx}(3x^2) = 6x$   
 $\frac{d}{dx}(-2 + 2\cos 2x) = -4\sin 2x$

That's still an indeterminate form, so repeat again:

$$\lim_{x \rightarrow 0} \frac{6}{-8\cos 2x} = \frac{6}{-8\cos 0} = -\frac{3}{4}$$
     $\frac{d}{dx}(6x) = 6$   
 $\frac{d}{dx}(-4\sin 2x) = -8\cos 2x$

And that limit exists, so

$$\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x} = -\frac{3}{4}$$

## Limits Using a Maclaurin Series

### How do I evaluate a limit using Maclaurin series?

- Limits of the form  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$  or  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$  may sometimes be evaluated by using Maclaurin series
- Usually this will be in a situation where attempting to evaluate the limit in the usual way returns an **indeterminate form**  $\frac{0}{0}$  or  $\frac{\pm\infty}{\pm\infty}$ .
- In such a case:
  - STEP 1: Find the Maclaurin series for  $f(x)$  and  $g(x)$
  - STEP 2: Rewrite  $\frac{f(x)}{g(x)}$  using the Maclaurin series in the numerator and denominator
  - STEP 3: Use algebra to simplify your new expression for  $\frac{f(x)}{g(x)}$  as far as possible
  - STEP 4: Evaluate the limit using your simplified form of the expression

#### Exam Tip

- Some limits of an indeterminate form can also be evaluated using **L'Hôpital's Rule**
- If an exam question does not specify a method to use, then you are free to use whichever method you prefer



### Worked example

Use Maclaurin series to evaluate the limit

$$\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x}$$

Maclaurin series for special functions	$e^x = 1 + x + \frac{x^2}{2!} + \dots$	$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$
	$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$	$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$
	$\arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$	

from exam formula booklet

$$\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x} = \frac{0^3}{-2(0) + \sin 0} = \frac{0}{0} \leftarrow \text{indeterminate form}$$

STEP 1:  $-2x + \sin 2x = -2x + \left(2x - \frac{(2x)^3}{3!} + \frac{(2x)^5}{5!} - \dots\right)$

$$= -\frac{4}{3}x^3 + \frac{4}{15}x^5 - \dots$$

# Exam Papers Practice

Copyright © 2024 Exam Papers Practice

STEP 2:  $\frac{x^3}{-2x + \sin 2x} = \frac{x^3}{-\frac{4}{3}x^3 + \frac{4}{15}x^5 - \dots}$

STEP 3:  $\frac{x^3}{-\frac{4}{3}x^3 + \frac{4}{15}x^5 - \dots} \times \frac{\frac{1}{x^3}}{\frac{1}{x^3}} = \frac{1}{-\frac{4}{3} + \frac{4}{15}x^2 - \dots}$

STEP 4:  $\lim_{x \rightarrow 0} \frac{1}{-\frac{4}{3} + \frac{4}{15}x^2 - \dots} = \frac{1}{-\frac{4}{3} + \frac{4}{15}(0)^2 - \dots} = \frac{1}{-\frac{4}{3}} = -\frac{3}{4}$

higher powers of x will also be zero

$$\lim_{x \rightarrow 0} \frac{x^3}{-2x + \sin 2x} = -\frac{3}{4}$$