



5.6 Kinematics

Contents

- ★ 5.6.1 Kinematics Toolkit
- * 5.6.2 Calculus for Kinematics



5.6.1 Kinematics Toolkit

Displacement, Velocity & Acceleration

What is kinematics?

- Kinematics is the branch of mathematics that models and analyses the motion of objects
- Common words such as distance, speed and acceleration are used in kinematics but are used according to their technical definition

What definitions do I need to be aware of?

- Firstly, only motion of an object in a **straight line** is considered
 - this could be a **horizontal** straight line
 - the **positive** direction would be to the **right**
 - or this could be a **vertical** straight line
 - the positive direction would be upwards

Particle

- A particle is the general term for an object
 - some questions may use a **specific** object such as a **car** or a **ball**

Time t seconds

- ullet Displacement, velocity and acceleration are all functions of time t
- Initially time is zero t=0

Displacement $S \, \text{m}$

- The displacement of a particle is its distance relative to a fixed point
 - the fixed point is often (but not always) the particle's initial position
- **Displacement** will be zero s = 0 if the object is at or has returned to its initial position
- Displacement will be negative if its position relative to the fixed point is in the negative direction (left or down)

Distance d_{m}

- Use of the word **distance** needs to be considered carefully and could refer to
 - the distance **travelled** by a particle
 - the (straight line) distance the particle is from a particular point
- Be careful not to confuse displacement with distance
 - if a bus route starts and ends at a bus depot, when the bus has returned to the depot, its

 displacement will be zero but the distance the bus has travelled will be the length of the route
- Distance is always positive

Velocity Vms⁻¹

lacktriangledown The **velocity** of a particle is the **rate of change** of its **displacement** at time t



- Velocity will be negative if the particle is moving in the negative direction
- A **velocity** of **zero** means the particle is **stationary** V = 0

Speed $|V| \text{ m s}^{-1}$

- Speed is the magnitude (a.k.a. absolute value or modulus) of velocity
 - as the particle is **moving** in a **straight line**, **speed** is the **velocity ignoring** the **direction**
 - if v = 4, |v| = 4
 - if v = -6, |v| = 6

Acceleration a m s⁻²

- lacktriangledown The **acceleration** of a particle is the **rate of change** of its **velocity** at time t
- Acceleration can be **negative** but this alone cannot fully describe the particle's motion
 - if **velocity** and **acceleration** have the **same** sign the particle is **accelerating** (speeding up)
 - if **velocity** and **acceleration** have **different** signs then the particle is **decelerating** (slowing down)
 - if acceleration is zero a = 0 the particle is moving with constant velocity
 - in all cases the **direction** of **motion** is determined by the **sign** of **velocity**

Are there any other words or phrases in kinematics I should know?

- Certain words and phrases can imply values or directions in kinematics
 - a particle described as "at rest" means that its velocity is zero, v = 0
 - a particle described as moving "due east" or "right" or would be moving in the positive horizontal direction
 - this also means that v > 0
 - a particle "dropped from the top of a cliff" or "down" would be moving in the negative vertical direction
 - this also means that v < 0

What are the key features of a velocity-time graph?


- The **gradient** of the graph equals the **acceleration** of an object
- A straight line shows that the object is accelerating at a constant rate
- A horizontal line shows that the object is moving at a constant velocity
- The area between graph and the x-axis tells us the change in displacement of the object
 - Graph above the x-axis means the object is moving forwards
 - Graph **below** the x-axis means the object is moving **backwards**
- The total displacement of the object from its starting point is the sum of the areas above the x-axis minus the sum of the areas below the x-axis
- The total distance travelled by the object is the sum of all the areas
- If the graph touches the x-axis then the object is stationary at that time
- If the graph is **above** the **x-axis** then the object has positive velocity and is **travelling forwards**
- If the graph is **below** the **x-axis** then the object has negative velocity and is **travelling backwards**



Worked example

A particle is projected vertically upwards from ground level, taking 8 seconds to return to the ground.

The velocity-time graph below illustrates the motion of the particle for these 8 seconds.



- How many seconds does the particle take to reach its maximum height? i) Give a reason for your answer.
- State, with a reason, whether the particle is accelerating or decelerating at time t=3. ii)



i. At maximum height, velocity is zero V= 0 at t=4

- "The particle takes 4 seconds to reach its maximum height. This is because its velocity is 0 m s⁻¹ at 4 seconds.
- ii. At L=3, velocity is POSITIVE

 Acceleration is the gradient of velocity

 At L=3, acceleration is NEGATIVE
 - .. At 3 seconds the particle is decelerating on its velocity and occeleration have different signs.



5.6.2 Calculus for Kinematics

Differentiation for Kinematics

How is differentiation used in kinematics?

- Displacement, velocity and acceleration are related by calculus
- In terms of differentiation and derivatives
 - velocity is the rate of change of displacement

$$v = \frac{\mathrm{d}s}{\mathrm{d}t} \text{ or } v(t) = s'(t)$$

acceleration is the rate of change of velocity

$$a = \frac{\mathrm{d}v}{\mathrm{d}t} \text{ or } a(t) = v'(t)$$

• so acceleration is also the second derivative of displacement

$$a = \frac{\mathrm{d}^2 s}{\mathrm{d}t^2} \text{ or } a(t) = s''(t)$$

- If a graph is not given you can use your GDC to draw one
 - you can then use your GDC's graphing features to find gradients
 - velocity is the gradient on a displacement (-time) graph
 - acceleration is the gradient on a velocity (-time) graph



Worked example

The displacement, s m, of a particle at t seconds, is modelled by $s(t) = 2t^3 - 27t^2 + 84t$

- i. Find v(t) and a(t).
- ii. Find the times at which the particle is at rest.

i.
$$v(t) = s'(t) = 6t^2 - 5+t + 8+t = 6(t^2 - 9t + 1+t)$$

 $a(t) = v'(t) = 12t - 5+t = 6(2t - 9)$

$$v(t) = 6(t-7)(t-2)$$

$$a(t) = 6(2t-9)$$
It's not essential to factorise the final answers

ii. The particle is at rest when u(t)=0

. The particle is at rest at 2 seconds and 7 seconds



Integration for Kinematics

How is integration used in kinematics?

• Since **velocity** is the **derivative** of **displacement** ($V = \frac{\mathrm{d}s}{\mathrm{d}t}$) it follows that

$$s = \int v \, \mathrm{d}t$$

Similarly, velocity will be an antiderivative of acceleration

$$v = \int a \, \mathrm{d}t$$

How would I find the constant of integration in kinematics problems?

- A **boundary** or **initial** condition would need to be known
 - phrases involving the word "initial", or "initially" are referring to time being zero, i.e. t=0
 - you might also be given information about the object at some other time (this is called a **boundary** condition)
 - substituting the values in from the initial or boundary condition would allow the constant of integration to be found

How are definite integrals used in kinematics?


- Definite integrals can be used to find the displacement of a particle between two points in time
 - $\int_{t_1}^{t_2} v(t) dt$ would give the **displacement** of the particle **between** the times $t = t_1$ and $t = t_2$
 - This can be found using a velocity-time graph by subtracting the total area below the horizontal axis from the total area above
 - $\int_{t_1}^{t_2} |v(t)| \, \mathrm{d}t$ gives the **distance** a particle has **travelled** between the times $t = t_1$ and $t = t_2$
 - This can be found using a velocity velocity-time graph by adding the total area below the horizontal axis to the total area above
 - Use a GDC to plot the modulus graph y = |v(t)|



Worked example


A particle moving in a straight horizontal line has velocity ($V \, {
m m \, s^{-1}}$) at time t seconds modelled by $v(t) = 8 \, t^3 - 12 \, t^2 - 2 \, t$.

- i. Given that the initial position of the particle is at the origin, find an expression for its displacement from the origin at time t seconds.
- ii. Find the displacement of the particle from the origin in the first five seconds of its motion.
- iii. Find the distance travelled by the particle in the first five seconds of its motion.





Use your GOC to sketch a velocity (-time) graph and use it to check to see if your arguers are cersible.



i. "initial" - E=0, "origin" - g=0

$$s(t) = \int v(t) dt = \int (8t^3 - 12t^2 - 2t) dt$$

$$s(t) = 2t^4 - 4t^3 - t^2 + c$$

at t=0, s=0, : c=0

where c is a constant

ii. "first fine seconds" -
$$t_1=0$$
, $t_2=5$
Using a GDC this would be
 $s=\int_{0.5}^{5} (8t^3-12t^2-2t) dt$

iii. Using a GOC this would be

$$d = \int_{0}^{5} |8E^{3} - 12E^{2} - 2E| dE$$

d for distance