

EDEXCEL INTERNATIONAL GCSE (9-1)

FURTHER PURE MATHEMATICS



Student Book Ali Datoo



PEARSON EDEXCEL INTERNATIONAL GCSE (9–1)

FURTHER PURE MATHEMATICS

Student Book

Ali Datoo

Greg Attwood, Keith Pledger, David Wilkins, Alistair Macpherson, Bronwen Moran, Joseph Petran and Geoff Staley

Published by Pearson Education Limited, 80 Strand, London, WC2R 0RL.

Copies of official specifications for all Pearson qualifications may be found on the website: https://qualifications.pearson.com

Text © Pearson Education Limited 2017

Edited by Linnet Bruce, Keith Gallick and Susan Lyons

Designed by Cobalt id

Typeset by Tech-Set Ltd, Gateshead, UK

Cover design by Pearson Education Limited

Picture research by Debbie Gallagher

Cover photo © Getty Images: Naahushi Kavuri / EyeEm

Inside front cover: Shutterstock.com: Dmitry Lobanov

The rights of Ali Datoo, Greg Attwood, Keith Pledger, David Wilkins, Alistair Macpherson, Bronwen Moran, Joseph Petran and Geoff Staley to be identified as authors of this work have been asserted by them in accordance with the Copyright, Designs and Patents Act 1988.

First published 2017

20 19 18 17

1098765432

British Library Cataloguing in Publication Data

A catalogue record for this book is available from the British Library ISBN 978 0 435 18854 2

Copyright notice

All rights reserved. No part of this publication may be reproduced in any form or by any means (including photocopying or storing it in any medium by electronic means and whether or not transiently or incidentally to some other use of this publication) without the written permission of the copyright owner, except in accordance with the provisions of the Copyright, Designs and Patents Act 1988 or under the terms of a licence issued by the Copyright Licensing Agency, Barnard's Inn, 86 Fetter Lane, London EC4A 1EN (www.cla.co.uk). Applications for the copyright owner's written permission should be addressed to the publisher.

Printed in Slovakia by Neografia

Acknowledgements

The author and publisher would like to thank the following individuals and organisations for permission to reproduce photographs:

(Key: b-bottom; c-centre; I-left; r-right; t-top)

123RF.com: ealisa 143; Alamy Stock Photo: Ammit 3; Getty Images: Christoph Hetzmannseder 2, David Crunelle / EyeEm 160, Ditto 24, EyeEm 136, hohl 74, Julia Antonio / EyeEm 60, Krzysztof Dydynski 106, Martin Barraud 124, Ozgur Donmaz 36, sebastian-julian 96, View Pictures / UIG 182; Shutterstock.com: Africa Studio 75, Akhenaton Images 90, Aleksandr Markin 128, andrea crisante 107t, Anette Holmberg 37, Brian Kinney 161, cigdem 148, IM_photo 123, Jacques Durocher 49t, jordache 180, Maridav 83, Miami2you 130, migrean 107b, Mikael Damkier 160, Nando 11, Nattawadee Supchapo 137, Neil Lockhart 61, Nicku 80, oksmit 125, Ondrej Prosicky 183, Patricia Hofmeester 131, Pavel L Photo and Video 154, photofriday 77t, Rawpixel.com 9, Sergey Edentod 47, Steve Noakes 25, Svetlana Privezentseva 49b, Syda Productions 77b, Viacheslav Nikolaenko 92, wavebreakmedia 121, worradirek 10, Zhukov Oleg 5

All other images @ Pearson Education

Endorsement statement

In order to ensure that this resource offers high-quality support for the associated Pearson qualification, it has been through a review process by the awarding body. This process confirms that this resource fully covers the teaching and learning content of the specification or part of a specification at which it is aimed. It also confirms that it demonstrates an appropriate balance between the development of subject skills, knowledge and understanding, in addition to preparation for assessment.

Endorsement does not cover any guidance on assessment activities or processes (e.g. practice questions or advice on how to answer assessment questions), included in the resource nor does it prescribe any particular approach to the teaching or delivery of a related course.

While the publishers have made every attempt to ensure that advice on the qualification and its assessment is accurate, the official specification and associated assessment guidance materials are the only authoritative source of information and should always be referred to for definitive guidance.

Pearson examiners have not contributed to any sections in this resource relevant to examination papers for which they have responsibility.

Examiners will not use endorsed resources as a source of material for any assessment set by Pearson. Endorsement of a resource does not mean that the resource is required to achieve this Pearson qualification, nor does it mean that it is the only suitable material available to support the qualification, and any resource lists produced by the awarding body shall include this and other appropriate resources.

| ABOUT THIS BOOK | vi |
|--|-----|
| CHAPTER 1: SURDS AND LOGARITHMIC FUNCTIONS | 2 |
| CHAPTER 2: THE QUADRATIC FUNCTION | 24 |
| CHAPTER 3: INEQUALITIES AND IDENTITIES | 36 |
| CHAPTER 4: SKETCHING POLYNOMIALS | 60 |
| CHAPTER 5: SEQUENCES AND SERIES | 74 |
| CHAPTER 6: THE BINOMIAL SERIES | 96 |
| CHAPTER 7: SCALAR AND VECTOR QUANTITIES | 106 |
| CHAPTER 8: RECTANGULAR CARTESIAN COORDINATES | 124 |
| CHAPTER 9: DIFFERENTIATION | 136 |
| CHAPTER 10: INTEGRATION | 160 |
| CHAPTER 11: TRIGONOMETRY | 182 |
| GLOSSARY | 212 |
| ANSWERS | 216 |
| INDEX | 238 |
| | |

| CHAPTER 1 | | FACTORISE A POLYNOMIAL BY USING THE FACTOR THEOREM | 52 |
|--|----------|--|----------|
| WRITE A NUMBER EXACTLY USING A SURD | 4 | USING THE REMAINDER THEOREM, FIND THE | |
| RATIONALISE THE DENOMINATOR OF A SURD | 5 | REMAINDER WHEN A POLYNOMIAL IS DIVIDED | F.4 |
| BE FAMILIAR WITH THE FUNCTIONS a^x AND $\log_b x$ AND RECOGNISE THE SHAPES OF THEIR GRAPHS | 7 | BY $(ax - b)$ EXAM PRACTICE QUESTIONS | 54 56 |
| BE FAMILIAR WITH EXPRESSIONS OF THE TYPE e^x AND USE THEM IN GRAPHS | 9 | CHAPTER SUMMARY | 58 |
| BE ABLE TO USE GRAPHS OF FUNCTIONS TO SOLVE EQUATIONS | 12 | CHAPTER 4 SKETCH CUBIC CURVES OF THE FORM | |
| WRITING AN EXPRESSION AS A LOGARITHM | 14 | $y = ax^3 + bx^2 + cx + d \text{ OR}$ | |
| UNDERSTAND AND USE THE LAWS OF LOGARITHMS | 15 | y = (x + a)(x + b)(x + c) SKETCH AND INTERPRET GRAPHS OF CUBIC | 62 |
| CHANGE THE BASE OF A LOGARITHM | 17 | FUNCTIONS OF THE FORM $y=x^3$ | 64 |
| SOLVE EQUATIONS OF THE FORM $a^x=b$ | 18 | SKETCH THE RECIPROCAL FUNCTION $y=rac{k}{x}$ | |
| EXAM PRACTICE QUESTIONS | 21 | WHERE & IS A CONSTANT | 65 |
| CHAPTER SUMMARY | 23 | SKETCH CURVES OF DIFFERENT FUNCTIONS TO SHOW POINTS OF INTERSECTION AND SOLUTIONS TO EQUATIONS | 67 |
| CHAPTER 2 | | APPLY TRANSFORMATIONS TO CURVES | 69 |
| FACTORISE QUADRATIC EXPRESSIONS WHERE | | EXAM PRACTICE QUESTIONS | 71 |
| THE COEFFICIENT OF x^2 IS GREATER THAN 1 | 26 | CHAPTER SUMMARY | 73 |
| COMPLETE THE SQUARE AND USE THIS TO SOLVE QUADRATIC EQUATIONS | 27 | CHAPTER 5 | |
| SOLVE QUADRATIC EQUATIONS USING THE QUADRATIC FORMULA | 28 | IDENTIFY AN ARITHMETIC SEQUENCE | 76 |
| UNDERSTAND AND USE THE DISCRIMINANT TO IDENTIFY WHETHER THE ROOTS ARE (i) EQUAL | | FIND THE COMMON DIFFERENCE, FIRST TERM AND n th TERM OF AN ARITHMETIC SERIES | 78 |
| AND REAL, (ii) UNEQUAL AND REAL OR (iii) NOT REAL UNDERSTAND THE ROOTS $lpha$ AND eta AND HOW TO | 29 | FIND THE SUM OF AN ARITHMETIC SERIES AND BE ABLE TO USE Σ NOTATION | 80 |
| USE THEM | 31 | IDENTIFY A GEOMETRIC SEQUENCE | 84 |
| EXAM PRACTICE QUESTIONS CHAPTER SUMMARY | 34 35 | FIND THE COMMON RATIO, FIRST TERM AND nth TERM OF A GEOMETRIC SEQUENCE | 85 |
| | 00 | FIND THE SUM OF A GEOMETRIC SERIES | 87 |
| CHAPTER 3 | | FIND THE SUM TO INFINITY OF A CONVERGENT GEOMETRIC SERIES | 90 |
| SOLVE SIMULTANEOUS EQUATIONS, ONE LINEAR AND ONE QUADRATIC | 38 | EXAM PRACTICE QUESTIONS | 93 |
| SOLVE LINEAR INEQUALITIES | 39 | CHAPTER SUMMARY | 95 |
| SOLVE QUADRATIC INEQUALITIES | 41 | | |
| GRAPH LINEAR INEQUALITIES IN TWO VARIABLES | 44 | CHAPTER 6 | |
| DIVIDE A POLYNOMIAL BY $(x \pm p)$ | 50 | USE $\binom{n}{r}$ TO WORK OUT THE COEFFICIENTS IN THE BINOMIAL EXPANSION | 98 |


| USE THE BINOMIAL EXPANSION TO EXPAND $(1 + x)^n$ | 100 | USE THE QUOTIENT RULE TO DIFFERENTIATE MORE | |
|---|-----|--|-----|
| DETERMINE THE RANGE OF VALUES FOR WHICH x IS | 101 | COMPLICATED FUNCTIONS | 148 |
| TRUE AND VALID FOR AN EXPANSION | 101 | FIND THE EQUATION OF THE TANGENT AND NORMAL TO THE CURVE $y = f(x)$ | 149 |
| EXAM PRACTICE QUESTIONS | 104 | FIND THE STATIONARY POINTS OF A CURVE AND | |
| CHAPTER SUMMARY | 105 | CALCULATE WHETHER THEY ARE MINIMUM OR | |
| CHAPTER 7 | | MAXIMUM STATIONARY POINTS | 151 |
| VECTOR NOTATION AND HOW TO DRAW VECTOR | | APPLY WHAT YOU HAVE LEARNT ABOUT TURNING POINTS TO SOLVE PROBLEMS | 153 |
| DIAGRAMS | 108 | EXAM PRACTICE QUESTIONS | 156 |
| PERFORM SIMPLE VECTOR ARITHMETIC AND | | CHAPTER SUMMARY | 159 |
| UNDERSTAND THE DEFINITION OF A UNIT VECTOR | 110 | OTAL TEN SOMMANT | 100 |
| USE VECTORS TO DESCRIBE THE POSITION OF A POINT IN TWO DIMENSIONS | 113 | CHAPTER 10 | |
| USE VECTORS TO DEMONSTRATE SIMPLE | | INTEGRATION AS THE REVERSE OF DIFFERENTIATION | 162 |
| PROPERTIES OF GEOMETRICAL FIGURES WRITE DOWN AND USE CARTESIAN COMPONENTS | 114 | UNDERSTAND HOW CALCULUS IS RELATED TO PROBLEMS INVOLVING KINEMATICS | 164 |
| OF A VECTOR IN TWO DIMENSIONS | 117 | USE INTEGRATION TO FIND AREAS | 166 |
| EXAM PRACTICE QUESTIONS | 120 | USE INTEGRATION TO FIND A VOLUME OF REVOLUTION | 171 |
| CHAPTER SUMMARY | 123 | RELATE RATES OF CHANGE TO EACH OTHER | 174 |
| | | EXAM PRACTICE QUESTIONS | 178 |
| CHAPTER 8 | | CHAPTER SUMMARY | 181 |
| WORK OUT THE GRADIENT OF A STRAIGHT LINE | 125 | | |
| FIND THE EQUATION OF A STRAIGHT LINE | 127 | CHAPTER 11 | |
| UNDERSTAND THE RELATIONSHIP BETWEEN PERPENDICULAR LINES | 128 | MEASURE ANGLES IN RADIANS | 184 |
| FIND THE DISTANCE BETWEEN TWO POINTS ON A LINE | 131 | CALCULATE ARC LENGTH AND THE AREA OF A CIRCLE USING RADIANS | 185 |
| FIND THE COORDINATES OF A POINT THAT DIVIDES A | | UNDERSTAND THE BASIC TRIGONOMETRICAL RATIOS | |
| LINE IN A GIVEN RATIO | 132 | AND SINE, COSINE AND TANGENT GRAPHS | 190 |
| EXAM PRACTICE QUESTIONS | 133 | USE SINE AND COSINE RULES | 193 |
| CHAPTER SUMMARY | 135 | USE SINE AND COSINE RULES TO SOLVE PROBLEMS IN 3D | 197 |
| CHAPTER 9 | | USE TRIGONOMETRY IDENTITIES TO SOLVE PROBLEMS | 199 |
| FIND THE GRADIENT FUNCTION OF A CURVE AND | | SOLVE TRIGONOMETRIC EQUATIONS | 202 |
| DIFFERENTIATE A FUNCTION THAT HAS MULTIPLE POWERS OF x | 138 | USE TRIGONOMETRIC FORMULAE TO SOLVE EQUATIONS | 206 |
| DIFFERENTIATE e^{ax} , $\sin ax$ AND $\cos ax$ | 141 | EXAM PRACTICE QUESTIONS | 208 |
| USE THE CHAIN RULE TO DIFFERENTIATE MORE | | CHAPTER SUMMARY | 210 |
| COMPLICATED FUNCTIONS | 143 | GLOSSARY | 212 |
| USE THE PRODUCT RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS | 146 | ANSWERS | 216 |

ABOUT THIS BOOK

This book is written for students following the Pearson Edexcel International GCSE (9–1) Further Pure Maths specification and covers both years of the course. The specification and sample assessment materials for Further Pure Maths can be found on the Pearson Qualifications website.

In each chapter, there are concise explanations and worked examples, plus numerous exercises that will help you build up confidence.


There are also exam practice questions and a chapter summary to help with exam preparation. Answers to all exercises are included at the back of the book as well as a glossary of Maths-specific terminology.




Examples provide a clear, instructional framework.
The blue highlighted text gives further explanation of the method.

Language is graded for speakers of English as an additional language (EAL), with advanced Maths-specific terminology highlighted and defined in the glossary at the back of the book.

Key Points boxes summarise the essentials.





ASSESSMENT OVERVIEW

The following tables give an overview of the assessment for the Edexcel International GCSE in Further Pure Mathematics.

We recommend that you study this information closely to help ensure that you are fully prepared for this course and know exactly what to expect in the assessment.

| PAPER 1 | PERCENTAGE | MARK | TIME | AVAILABILITY |
|--|------------|------|---------|--|
| Written examination paper Paper code 4PM1/01C Externally set and assessed by Edexcel | 50% | 100 | 2 hours | January and June examination series First assessment June 2019 |
| | | | | |
| PAPER 2 | PERCENTAGE | MARK | TIME | AVAILABILITY |

CONTENT SUMMARY

- Number
- · Algebra and calculus
- · Geometry and calculus

ASSESSMENT

- Each paper will consist of around 11 questions with varying mark allocations per questions, which will be stated on the paper
- Each paper will contain questions from any part of the specification content, and the solution of any questions may require knowledge of more than one section of the specification content
- · A formulae sheet will be included in the written examinations
- A calculator may be used in the examinations

ASSESSMENT OBJECTIVES AND WEIGHTINGS

| ASSESSMENT OBJECTIVE | DESCRIPTION | % IN INTERNATIONAL GCSE |
|----------------------|--|-------------------------|
| AO1 | Demonstrate a confident knowledge of the techniques of pure mathematics required in the specification | 30%-40% |
| AO2 | Apply a knowledge of mathematics to the solutions of problems for which an immediate method of solution is not available and which may involve knowledge of more than one topic in the specification | 20%–30% |
| AO3 | Write clear and accurate mathematical solutions | 35%–50% |

RELATIONSHIP OF ASSESSMENT OBJECTIVES TO UNITS

| UNIT NUMBER | ASSESSMENT OBJECTIVE | | | | |
|------------------------------|----------------------|---------|-----------|--|--|
| | AO1 | AO2 | AO3 | | |
| Paper 1 | 15%–20% | 10%-15% | 17.5%–25% | | |
| Paper 2 | 15%–20% | 10%-15% | 17.5%–25% | | |
| Total for International GCSE | 30%-40% | 20%-30% | 35%-50% | | |

ASSESSMENT SUMMARY

The Edexcel International GCSE in Further Pure Mathematics requires students to demonstrate application and understanding of the following topics.

Number

• Use numerical skills in a purely mathematical way and in real-life situations.

Algebra and calculus

- Use algebra and calculus to set up and solve problems.
- Develop competence and confidence when manipulating mathematical expressions.
- · Construct and use graphs in a range of situations.

Geometry and trigonometry

- Understand the properties of shapes, angles and transformations.
- Use vectors and rates of change to model situations.
- · Use coordinate geometry.
- · Use trigonometry.

Students will be expected to have a thorough knowledge of the content common to the Pearson Edexcel International GCSE in Mathematics (Specification A) (Higher Tier) or Pearson Edexcel International GCSE in Mathematics (Specification B).

Questions may be set which assumes knowledge of some topics covered in these specifications, however knowledge of statistics and matrices will not be required.

Students will be expected to carry out arithmetic and algebraic manipulation, such as being able to change the subject of a formula and evaluate numerically the value of any variable in a formula, given the values of the other variables. The use and notation of set theory will be adopted where appropriate.

CALCULATORS

Students will be expected to have access to a suitable electronic calculator for all examination papers. The electronic calculator should have these functions as a minimum:

+, -, \times , \div , π , x^2 , \sqrt{x} , $\frac{1}{x}$, x^y , $\ln x$, e^x , sine, cosine and tangent and their inverses in degrees and decimals of a degree or radians.

Prohibitions

Calculators with any of the following facilities are prohibited in all examinations:

databanks

retrieval of text or formulae

QWERTY keyboards

built-in symbolic algebra manipulations

symbolic differentiation or integration.

FORMULAE SHEET

These formulae will be provided for you during the examination.

MENSURATION

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times$ slant height

Volume of sphere $=\frac{4}{3}\pi r^3$

SERIES

Arithmetic series

Sum to n terms $S_n = \frac{n}{2}[2a + (n-1)d]$

Geometric series

Sum to
$$n$$
 terms, $S_n = \frac{a(1-r^n)}{(1-r)}$

Sum to infinity,
$$S_{\infty} = \frac{a}{1 - r}$$

Binomial series

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \ldots + \frac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots$$
 for $|x| < 1, n \in \mathbb{Q}$

CALCULUS

Quotient rule (differentiation)

$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$$

TRIGONOMETRY

Cosine rule

In triangle *ABC*: $a^2 = b^2 + c^2 - 2bc \cos A$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$sin(A + B) = sin A cos B + cos A sin B$$

$$sin(A - B) = sin A cos B - cos A sin B$$

$$cos(A + B) = cos A cos B - sin A sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

LOGARITHMS

$$\log_a x = \frac{\log_b x}{\log_b a}$$

FORMULAE TO KNOW

The following are formulae that you are expected to know and remember during the examination. These formulae will not be provided for you. Note that this list is not exhaustive.

LOGARITHMIC FUNCTIONS AND INDICES

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \frac{x}{y} = \log_a x - \log_a y$$

$$\log_a x^k = k \log_a x$$

$$\log_a \frac{1}{x} = -\log_a x$$

$$\log_a a = 1$$

$$\log_a 1 = 0$$

$$\log_a b = \frac{1}{\log_b a}$$

OUADRATIC EQUATIONS

$$ax^2 + bx + c = 0$$
 has roots given by $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

When the roots of $ax^2 + bx + c = 0$ are α and β then $\alpha + \beta = -\frac{b}{a}$ and $\alpha\beta = \frac{c}{a}$ and the equation can be written $x^2 - (\alpha + \beta)x + \alpha\beta = 0$

SERIES

Arithmetic series: nth term = l = a + (n - 1)d

Geometric series: nth term = $a r^{n-1}$

COORDINATE GEOMETRY

The gradient of the line joining two points (x_1, y_1) and (x_2, y_2) is $\frac{y_2 - y_1}{x_2 - x_1}$

The distance d between two points (x_1, y_1) and (x_2, y_2) is given by

$$d^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$$

The coordinates of the point dividing the line joining (x_1, y_1) and (x_2, y_2) in the ratio m:n are

$$\left(\frac{n x_{\scriptscriptstyle 1} - m x_{\scriptscriptstyle 2}}{m+n}, \frac{n y_{\scriptscriptstyle 1} - m y_{\scriptscriptstyle 2}}{m+n}\right)$$

CALCULUS

Differentiation: function derivative

 x^n nx^{n-1} $\sin ax$ $a\cos ax$ $\cos ax$ $-a\sin ax$

 e^{ax} ae^{ax}

f(x)g(x) f'(x)g(x) + f(x)g'(x)

f(g(x)) f'(g(x))g'x

Integration: function derivative

 $\frac{1}{n+1}x^{n+1}+c \quad n\neq -1$

 $\sin ax$ $-\frac{1}{a}\cos ax + c$

 $\cos ax$ $\frac{1}{a}\sin ax + c$

 e^{ax} $\frac{1}{a}e^{ax} + c$

AREA AND VOLUME

Area between a curve and the *x*-axis = $\int_a^b y dx$, $y \ge 0$

 $\left|\int_a^b y \mathrm{d}x\right|,\,y<0$

Area between a curve and the y-axis $=\int\limits_{c}^{d}x\mathrm{d}y$, $x\geqslant0$

 $\left|\int_{c}^{d}x\mathrm{d}y\right|,\ x<0$

Area between g(x) and $f(x) = \int_a^b |g(x) - f(x)| dx$

Volume of revolution = $\int_a^b \pi y^2 dx$ or $\int_c^d \pi x^2 dy$

TRIGONOMETRY

Radian measure: length of arc = $r\theta$

area of sector $=\frac{1}{2}r^2\theta$

In a triangle *ABC*: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

 $\cos^2\theta + \sin^2\theta = 1$

area of a triangle $=\frac{1}{2}ab \sin C$

CHAPTER 1



1 SURDS AND LOGARITHMIC FUNCTIONS

The Richter scale, which describes the energy released by an earthquake, uses the base 10 logarithm as its unit. An earthquake of **magnitude** 9 is 10 times as powerful as one of magnitude 8, and 100 000 times as powerful as one of magnitude 4.

The **devastating** 2004 earthquake in the Indian Ocean had a magnitude of 9. Thankfully, such events are rare. The most common earthquakes, which occur over 100 000 times a year, are magnitude 2 to 3, so humans can hardly feel them.



LEARNING OBJECTIVES

- Write a number exactly using surds
- Rationalise the denominator of a surd
- Be familiar with the **functions** a^x and $\log_b x$ and recognise the shapes of their graphs
- Be familiar with functions including e^x and similar terms, and use them in graphs
- Use graphs of functions to solve equations
- Rewrite expressions including powers using logarithms instead
- Understand and use the laws of logarithms
- · Change the base of a logarithm
- Solve equations of the form $a^x = b$

STARTER ACTIVITIES

- Simplify
 - a $y^6 \times y^5$
 - $(x^2)^4$

- **b** $2q^3 \times 4q^4$
- $(a^4)^2 \div a^3$

- c $3k^2 \times 3k^7 \times 3k^{-3}$
- $\mathbf{f} = 64x^4y^6 \div 4xy^2$

- Simplify
 - $(m^3)^{\frac{1}{2}}$

b $3p^{\frac{1}{2}} \times p^3$

c $28c^{\frac{2}{3}} \div 7c^{\frac{1}{3}}$

d $6b^{\frac{1}{2}} \times 3b^{-\frac{1}{2}}$

- $e 27p^{\frac{2}{3}} \div 9p^{\frac{1}{6}}$
- f $5y^6 \times 3y^{-7}$

- 3 > Evaluate
 - a $16^{\frac{1}{2}}$

b $125\frac{1}{3}$

c 8⁻²

 $(-2)^{-3}$

 $\left(\frac{6}{7}\right)^0$

f 81⁻⁴

 $g \left(\frac{9}{16}\right)^{-}$

WRITE A NUMBER EXACTLY USING A SURD

A surd is a number that cannot be simplified to remove a square root (or a cube root, fourth root etc). Surds are irrational numbers.

HINT

An irrational number is a number that cannot be expressed as a fraction, for example π .

| NUMBER | DECIMAL | IS IT A SURD? |
|----------------------|-------------|---------------|
| √1 | 1 | No |
| $\sqrt{2}$ | √2 1.414213 | |
| $\sqrt{4}$ | 2 | No |
| $\sqrt{\frac{1}{4}}$ | 0.5 | No |
| $\sqrt{\frac{2}{3}}$ | 0.816496 | Yes |

You can manipulate surds using these rules:

$$\sqrt{ab} = \sqrt{a} \times \sqrt{b}$$

$$\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$$

EXAMPLE 1

Simplify

SKILLS CRITICAL

THINKING

a √12

 $\sqrt{20}$

 $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

a
$$\sqrt{12}$$

$$= \sqrt{4 \times 3}$$

$$=\sqrt{4}\times\sqrt{3}$$

$$=2\sqrt{3}$$

Use the rule $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$

$$\sqrt{4} = 2$$

b
$$\frac{\sqrt{20}}{2}$$

$$=\frac{\sqrt{4\times5}}{2}$$

$$=\frac{2\times\sqrt{5}}{2}$$

$$=\sqrt{5}$$

 $\sqrt{20} = \sqrt{4} \times \sqrt{5}$

$$\sqrt{4} = 2$$

 $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

$$=5\sqrt{6}-2\sqrt{6}\sqrt{4}+\sqrt{6}\sqrt{49}$$

$$=\sqrt{6}(5-2\sqrt{4}+\sqrt{49})$$

$$=\sqrt{6}(5-2\times2+7)$$

$$= 8\sqrt{6}$$

 $\sqrt{6}$ is a common factor

Work out the square roots $\sqrt{4}$ and $\sqrt{49}$

$$5 - 4 + 7 = 8$$

EXERCISE 1

SKILLS

CRITICAL THINKING Simplify without using a calculator

Simplify without using a calculator

a
$$\frac{\sqrt{60}}{2}$$

b
$$\frac{\sqrt{135}}{2}$$

$$\frac{\sqrt{96}}{6}$$

√128 8

Simplify without using a calculator

a
$$6\sqrt{3} - 2\sqrt{3}$$

b
$$7\sqrt{3} - \sqrt{12} + \sqrt{48}$$

b
$$7\sqrt{3} - \sqrt{12} + \sqrt{48}$$
 c $\sqrt{112} + 2\sqrt{172} - \sqrt{63}$

d
$$6\sqrt{48} - 3\sqrt{12} + 2\sqrt{27}$$
 e $3\sqrt{578} - \sqrt{162} + 4\sqrt{32}$ **f** $2\sqrt{5} \times 3\sqrt{5}$

$$9.3\sqrt{578} - \sqrt{162} + 4\sqrt{32}$$

$$12\sqrt{5}\times3\sqrt{5}$$

g
$$6\sqrt{7} \times 4\sqrt{7}$$

h
$$4\sqrt{8} \times 6\sqrt{8}$$

Simplify without using a calculator

a
$$6(4-\sqrt{12})$$

b
$$9(6-3\sqrt{29})$$

c
$$4(1+\sqrt{3})+3(3+2\sqrt{3})$$

d
$$3(\sqrt{2} - \sqrt{7}) - 5(\sqrt{2} + \sqrt{7})$$
 e $(4 + \sqrt{3})(4 - \sqrt{3})$

$$(4 + \sqrt{3})(4 - \sqrt{3})$$

f
$$(2\sqrt{7} - \sqrt{6})(\sqrt{7} - 2\sqrt{6})$$
 g $(\sqrt{8} + \sqrt{5})(\sqrt{8} - \sqrt{5})$

g
$$(\sqrt{8} + \sqrt{5})(\sqrt{8} - \sqrt{5})$$

5 A garden is $\sqrt{30}$ m long and $\sqrt{8}$ m wide. The garden is covered in grass except for a small rectangular pond which is $\sqrt{2}$ m long and $\sqrt{6}$ m wide.

> Express the area of the pond as a percentage of the area of the garden.



6 Find the value of $2p^2 - 3pq$ when $p = \sqrt{2} + 3$ and $q = \sqrt{2} - 2$

HINT

In the denominator, the multiplication gives the difference of two squares, with the result $a^2 - b^2$, which means the surd disappears.

RATIONALISE THE DENOMINATOR OF A SURD

Rationalising the denominator of a surd means removing a root from the denominator of a fraction. You will usually need to rationalise the denominator when you are asked to simplify it.

The rules for rationalising the denominator of a surd are:

- For fractions in the form $\sqrt{\frac{1}{a}}$, multiply the numerator and denominator by \sqrt{a}
- For fractions in the form $\frac{1}{a+\sqrt{b}}$, multiply the numerator and denominator by $a-\sqrt{b}$
- For fractions in the form $\frac{1}{a-\sqrt{b}}$, multiply the numerator and denominator by $a+\sqrt{b}$

EXAMPLE 2

Rationalise the denominator of

a
$$\frac{1}{\sqrt{3}}$$

b
$$\frac{1}{3+\sqrt{2}}$$

$$\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$$

a
$$\frac{1}{\sqrt{3}}$$

$$= \frac{1 \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}}$$

Multiply the top and bottom by $\sqrt{3}$

$$=\frac{\sqrt{3}}{3}$$

$$\sqrt{3} \times \sqrt{3} = (\sqrt{3})^2 = 3$$

b
$$\frac{1}{3+\sqrt{2}}$$

$$=\frac{1\times(3-\sqrt{2})}{(3+\sqrt{2})(3-\sqrt{2})}$$

$$=\frac{3-\sqrt{2}}{9-3\sqrt{2}+3\sqrt{2}-2}$$

$$=\frac{3-\sqrt{2}}{3-3\sqrt{2}}$$

Multiply the top and bottom by $3 - \sqrt{2}$

$$\sqrt{2} \times \sqrt{2} = 2$$

$$9-2=7$$
, $-3\sqrt{2}+3\sqrt{2}=0$

c
$$\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$$

= $\frac{(\sqrt{5} + \sqrt{2})(\sqrt{5} + \sqrt{2})}{(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2})}$

Multiply the top and bottom by $\sqrt{5} + \sqrt{2}$

$$=\;\frac{5+\sqrt{5}\sqrt{2}\,+\sqrt{2}\sqrt{5}\,+2}{5-2}$$

$$\sqrt{5}\sqrt{2}\,-\sqrt{2}\sqrt{5}\,=\,0$$
 in the denominator

$$=\frac{7+2\sqrt{10}}{3}$$

$$\sqrt{5}\sqrt{2} = \sqrt{10}$$

EXERCISE 2

Rationalise

SKILLS EXECUTIVE

$$\frac{1}{\sqrt{3}}$$

b
$$\frac{1}{\sqrt{7}}$$

$$\frac{2}{\sqrt{3}}$$

d
$$\frac{\sqrt{6}}{\sqrt{3}}$$

e
$$\frac{12}{\sqrt{3}}$$

f
$$\frac{3\sqrt{5}}{\sqrt{3}}$$

$$\frac{9\sqrt{12}}{2\sqrt{18}}$$

h
$$\frac{1}{2-\sqrt{3}}$$

2 Rationalise

a
$$\frac{\sqrt{6}}{\sqrt{3}+\sqrt{6}}$$

b
$$\frac{2+\sqrt{3}}{2-\sqrt{3}}$$

c
$$\frac{\sqrt{2} - \sqrt{3}}{\sqrt{2} + \sqrt{3}}$$

d
$$\frac{4\sqrt{2}-2\sqrt{3}}{\sqrt{2}+\sqrt{3}}$$

e
$$\frac{\sqrt{2} + 2\sqrt{5}}{\sqrt{5} - \sqrt{2}}$$

f
$$\frac{\sqrt{7} + \sqrt{3}}{\sqrt{7} - \sqrt{3}}$$

$$h \ \frac{2\sqrt{5} - 3\sqrt{7}}{5\sqrt{6} + 4\sqrt{2}}$$

i
$$\frac{2 + \sqrt{10}}{\sqrt{2} + \sqrt{5}}$$

j
$$\frac{ab}{a\sqrt{b}-b\sqrt{a}}$$

$$k \frac{a-b}{a\sqrt{b}-b\sqrt{a}}$$

BE FAMILIAR WITH THE FUNCTIONS a^x AND $\log_a x$ AND RECOGNISE THE SHAPES OF THEIR GRAPHS

You need to be familiar with functions in the form $y = a^x$ where a > 0

Look at a table of values for $y = 2^x$

| \boldsymbol{x} | -3 | -2 | -1 | 0 | 1 | 2 | 3 |
|------------------|-----|---------------|-----|---|---|---|---|
| y | 1/8 | $\frac{1}{4}$ | 1/2 | 1 | 2 | 4 | 8 |


Note: $2^0 = 1$

In fact a^0 is always equal to 1 if a is positive and

$$2^{-3}=\frac{1}{2^3}=\frac{1}{8}$$

a negative index turns the number into its reciprocal

The graph of $y = 2^x$ looks like this:



Note: the *x*-axis is an **asymptote** to the curve.

Other graphs of the type $y = a^x$ have similar shapes, always passing through (0, 1).

EXAMPLE 3

a On the same axes, **sketch** the graphs of $y = 3^x$, $y = 2^x$ and $y = 1.5^x$

SKILLS

b On another set of axes, sketch the graphs of $y = \left(\frac{1}{2}\right)^x$ and $y = 2^x$

ANALYSIS


a For all three graphs, y = 1 when x = 0

When
$$x > 0$$
, $3^x > 2^x > 1.5^x$

$$a^0 = 1$$

When
$$x < 0$$
, $3^x < 2^x < 1.5^x$

Work out the relative positions of the graphs




b
$$\frac{1}{2} = 2^{-1}$$

so,
$$y = \left(\frac{1}{2}\right)^x$$
 is the same as $y = (2^{-1})^x = 2^{-x}$


$$(a^m)^n=a^{mn}$$


Therefore the graph of $y=\left(\frac{1}{2}\right)^x$ is a reflection in the y-axis of the graph of $y=2^x$




EXAMPLE 4

If you compare the graphs of $y = 2^x$ and $y = \log_2 x$ you see the following relationship:








EXAMPLE 5

On the same set of axes sketch the graphs $y = \log_2 x$ and $y = \log_5 x$



Note:

For both graphs y = 0 when x = 1, since $\log_a 1 = 0$ for every value of a.

$$\log_2 2 = 1$$
 so $y = \log_2 x$ passes through (2, 1)

and
$$log_5 5 = 1$$
 so $y = log_5 x$ passes through (5, 1)

EXERCISE 3

1 > On the same set of axes sketch the graphs of

SKILLS

ANALYSIS
REASONING

- **a** $y = 5^x$ **b** $y = 7^x$
- $y = \left(\frac{1}{3}\right)^x$

- 2 Don the same set of axes sketch the graphs of
 - $y = \log_5 x$
- $y = \log_7 x$
- c Write down the coordinates of the point of intersection of these two graphs.
- 3 > On the same set of axes sketch the graphs of
 - **a** $y = 3^x$
- $\mathbf{b} \quad y = \log_3 x$
- 4 > On the same set of axes sketch the graphs of
 - $y = \log_3 x$
- $\mathbf{b} \quad y = \log_5 x$
- $y = \log_{0.5} x$
- **d** $y = \log_{0.25} x$

BE FAMILIAR WITH EXPRESSIONS OF THE TYPE e^x and use them in graphs

Consider this example: Zainab opens an account with \$1.00. The account pays 100% interest per year. If the interest is credited once, at the end of the year, her account will contain \$2.00. How much will it contain after a year if the interest is calculated and credited more frequently? Let us investigate this more thoroughly.



| HOW OFTEN INTEREST IS CREDITED INTO THE ACCOUNT | VALUE OF ACCOUNT AFTER 1 YEAR (\$) |
|---|---|
| Yearly | $\left(1+\frac{1}{1}\right)^1=2$ |
| Semi-annually | $\left(1 + \frac{1}{2}\right)^2 = 2.25$ |
| Quarterly | $\left(1+\frac{1}{4}\right)^4=2.441406$ |
| Monthly | $\left(1 + \frac{1}{12}\right)^{12} = 2.61303529$ |
| Weekly | $\left(1 + \frac{1}{52}\right)^{52} = 2.69259695$ |
| Daily | $\left(1 + \frac{1}{365}\right)^{365} = 2.71456748$ |
| Hourly | $\left(1 + \frac{1}{8760}\right)^{8760} = 2.71812669$ |
| Every minute | $\left(1 + \frac{1}{525600}\right)^{525600} = 2.7182154$ |
| Every second | $\left(1 + \frac{1}{31536000}\right)^{31536000} = 2.71828247$ |

The amount in her account gets bigger and bigger the more often the interest is compounded, but the rate of growth slows. As the number of compounds increases, the calculated value appears to be approaching a fixed value. This value gets closer and closer to a fixed value of 2.71828247254......This number is called 'e'.

The number e is called a natural exponential because it arises naturally in mathematics and has numerous real life applications.





EXAMPLE 6


Draw the graphs of e^x and e^{-x}

SKILLS ANALYSIS INTERPRETATION

| \boldsymbol{x} | -2 | -1 | 0 | 1 | 2 | 3 | 4 |
|------------------|------|------|---|-----|-----|----|----|
| e^x | 0.14 | 0.37 | 1 | 2.7 | 7.4 | 20 | 55 |








EXAMPLE 7


Draw the graphs of these exponential functions.

- $y = e^{2x}$
- **b** $y = 10e^{-x}$
- $y = 3 + 4e^{\frac{1}{2}x}$


| a | x | -2 | -1 | 0 | 1 | 2 |
|---|-----------------|------|-----|---|-----|----|
| | e ^{2x} | 0.02 | 0.1 | 1 | 7.4 | 55 |







| C | \boldsymbol{x} | -2 | -1 | 0 | 1 | 2 |
|---|-------------------------|-----|-----|---|-----|------|
| | $3 + 4e^{\frac{1}{2}x}$ | 4.5 | 5.4 | 7 | 8.9 | 13.4 |



On pages 7–9 you saw the connection between $y = \log_a x$ and and $y = a^x$. The function $y = \log_e x$ is particularly important in mathematics and so it has a special notation:

$$\log_{\rm e} x \equiv \ln x$$

Your calculator should have a special button for evaluating $\ln x$.



EXAMPLE 8

Solve these equations.

a
$$e^x = 3$$

$$\ln x = 4$$

a When
$$e^x = 3$$

b When
$$\ln x = 4$$

$$x = \ln 3$$

$$x = e^4$$

As you can see, the inverse of e^x is $\ln x$ (and vice versa)

EXAMPLE 9


Sketch these graphs on the same set of axes.

$$y = \ln x$$

b
$$y = \ln(3 - x)$$

$$y = 3 + \ln(2x)$$

а-с



EXERCISE 4

Sketch these graphs.

SKILLS ANALYSIS REASONING

a
$$y = e^x + 1$$

b
$$y = 4e^{-2x}$$

$$y = 2e^x - 3$$

d
$$y = 6 + 10^{1x}$$

$$y = 100e^{-x} + 10$$

2 > Sketch these graphs, stating any asymptotes and intersections with the axes.

a
$$y = \ln(x + 1)$$

b
$$y = 2 \ln x$$

$$y = \ln(2x)$$

d
$$y = \ln(4 - x)$$


$$y = 4 + \ln(x + 2)$$

BE ABLE TO USE GRAPHS OF FUNCTIONS TO SOLVE EQUATIONS

EXAMPLE 10

- a Complete the table of values for: $y = e^{\frac{1}{2}x} 2$ Giving your answers to two decimal places where appropriate.
- **b** Draw the graph of $y = e^{\frac{1}{2}x} 2$ for $0 \le x \le 5$
- Use your graph to estimate, to 2 significant figures, the solution of the equation $e^{\frac{1}{2}x} = 8$ Show your method clearly.
- d By drawing a suitable line on your graph, estimate to 2 significant figures the solution to the equation $x = 2 \ln(7 2x)$

| а | x | -1 | 0 | 1 | 2 | 3 | 4 | 5 |
|---|---|-------|----|-------|------|------|------|-------|
| | y | -1.39 | -1 | -0.35 | 0.72 | 2.48 | 5.39 | 10.18 |



y = 5 - 2x

HINT

Make the LHS = $e^{\frac{1}{2}x} - 2$ i.e. the equation of the graph. To do this, you need to subtract 2 from 8 and draw the line y = 6 (as shown in the diagram).

HINT

Make the **LHS** equal to the given equation i.e $e^{\frac{1}{2}x} - 2$. Draw the line y = 5 - 2x (as shown in the diagram) on your graph and find points of intersection.

$$e^{\frac{1}{2}x} = 8$$

$$e^{\frac{1}{2}x} - 2 = 8 - 2$$

$$e^{\frac{1}{2}x} = 6$$

So the solution is the intersection of the curve $y=e^{\frac{1}{2}x}$ and y=6 $x\approx 4.15$ (In the exam you will be allowed a range of values.)

d
$$x = 2 \ln(7 - 2x)$$

 $\frac{x}{2} = \ln(7 - 2x)$

$$e^{\frac{1}{2}x} = 7 - 2x$$

$$e^{\frac{1}{2}x} - 2 = 7 - 2x - 2$$

$$e^{\frac{1}{2}x} = 5 - 2x$$

So, the solution is the intersection of the curve and $y=\mathrm{e}^{\frac{1}{2}x}$ and the line $y=5-2x,\,x\approx2.1$


EXAMPLE 11

a Complete the table below of values of $y = 2 + \ln x$, giving your values of y to decimal places.

| \boldsymbol{x} | 0.1 | 0.5 | 1 | 1.5 | 2 | 3 | 4 |
|------------------|------|------|---|------|------|------|---|
| y | -0.3 | 1.31 | | 2.41 | 2.69 | 3.10 | |

- **b** Draw the graph of $y = 2 + \ln x$ for $0.1 \le x \le 4$
- Use your graph to estimate, to 2 significant figures, the solution of the equation $\ln x = 0.5$
- **d** By drawing a suitable line on your graph estimate, to 1 significant figure, the solution of the equation $x = e^{x-2}$


| а | \boldsymbol{x} | 0.1 | 0.5 | 1 | 1.5 | 2 | 3 | 4 |
|---|------------------|------|------|---|------|------|------|------|
| | y | -0.3 | 1.31 | 2 | 2.41 | 2.69 | 3.10 | 3.39 |



c $2 + \ln x = 0.5$ Make the LHS = $2 + \ln x$ i.e the equation of the graph. Therefore you need to add

In x = 2.5 + 2 from 0.5 and draw the line y = 2.5

$$y = 2.5$$



The solution is the intersection of the curve and the line y = 2.5. From the graph this is approximately 1.6. In the exam you will be given a small range of answers.


 $x = e^{x-2}$

Using the properties of logs

$$\ln x = x - 2$$

 $\ln x + 2 = x - 2 + 2$

Make the LHS equal to the given equation i.e. $\ln x + 2$.



- a Draw the graph $y = 3 + 2e^{-\frac{1}{2}x}$ for $0 \le x \le 6$
 - b Use your graph to estimate, to 2 significant figures, the solution to the equation $e^{-\frac{1}{2}x} = 0.5$, showing your method clearly.
 - By drawing a suitable line, estimate, to 2 significant figures, the solution of the equation $x = -2 \ln\left(\frac{x-2}{2}\right)$
- **2 a** Draw the graph $y = 2 + \frac{1}{3}e^x$ for $-1 \le x \le 3$
 - b Use your graph to estimate, to 2 significant figures, the solution to the equation $e^x = 12$ showing your method clearly.
 - By drawing a suitable line, estimate, to 2 significant figures, the solution to the equation $x = \ln(6 - 6x)$
- 3 \triangleright a Complete the table below of values of $y = 5 \sin 2x 2 \cos x$, giving your values of y to 2 decimal places.

| \boldsymbol{x} | 0 | 15 | 30 | 45 | 60 | 75 | 90 |
|------------------|----|------|----|------|------|----|----|
| y | -2 | 0.57 | | 3.59 | 3.33 | | 0 |

- **b** Draw the graph of $y = 5 \sin 2x 2 \cos x$ for $\theta \le x \le 90^{\circ}$
- Use your graph to estimate, to 2 significant figures, the solution of the equation $2(1 + \cos x) = 5 \sin 2x$ showing your method clearly.

HINT

In the exam log₁₀ will be written as lg. This textbook uses Ig for log₁₀

WRITING AN EXPRESSION AS A LOGARITHM

 $\log_a n = x$ means that $a^x = n$, where a is called the base of the logarithm.

- $\log_a 1 = 0$ (a > 0), because $a^0 = 1$
- $\log_a a = 1$ (a > 0), because $a^1 = a$

EXAMPLE 12

Write as a logarithm $2^5 = 32$

SKILLS ADAPTABILITY

$$2^5 = 32$$

So
$$\log_2 32 = 5$$

Here
$$a = 2, x = 5, n = 32$$

Here 2 is the base, 5 is the logarithm. In words, you would say '2 to the power of 5 equals 32'. You would also say 'the logarithm of 32, to base 2, is 5'.

EXAMPLE 13

Rewrite using a logarithm

$$10^3 = 1000$$

$$5^4 = 625$$

$$c$$
 $2^{10} = 1024$

a
$$lg 1000 = 3$$

$$\log_5 625 = 4$$

$$\log_2 1024 = 10$$

EXAMPLE 14

Find the value of

- $a \log_3 81$
- $\log_4 0.25$
- $c \log_{0.5} 4$
- d $\log_a(a^5)$

$$\log_3 81 = 4$$

$$3^4 = 81$$

b
$$\log_4 0.25 = -1$$

$$4^{-1} = \frac{1}{4} = 0.25$$

$$\log_{0.5} 4 = -2$$

$$0.5^{-2} = \left(\frac{1}{2}\right)^{-2} = 2^2 = 4$$

d
$$\log_a(a^5) = 5$$

$$a^5 = a^5$$

EXERCISE 6

SKILLS CRITICAL THINKING

Rewrite these exponentials as logarithms.

$$4^3 = 64$$

b
$$5^{-2} = \frac{1}{25}$$

$$86 = 262144$$

d
$$3^x = 9$$

$$8^x = 1$$

f
$$2^x = \frac{1}{4}$$

2 Write these logarithms in exponential form.

$$\log_3 81 = 4$$

$$\log_3 729 = 6$$

d
$$\log_{16} 4 = \frac{1}{2}$$

e
$$\log_3\left(\frac{1}{27}\right) = -3$$
 f $\lg 1000 \ 0.01 = -2$

f
$$lg1000 0.01 = -2$$

3 Without a calculator find the value of

$$a \log_2 4$$

e
$$\log_5(\frac{1}{125})$$
 f $\lg\sqrt{10}$ g $\log_3\sqrt{27}$ h $\log_3\sqrt[5]{3}$

g
$$\log_3\sqrt{27}$$

$$\log_3 \sqrt[5]{3}$$

4 Find the value of x for which

$$\log_3 x = 4$$

b
$$\log_6 x = 3$$

$$\log_x 64 = 3$$

d
$$\log_x 16 = \frac{4}{3}$$

d
$$\log_x 16 = \frac{4}{3}$$
 e $\log_x 64 = \frac{2}{3}$

5 Find using your calculator

- a lg 20
- **b** lg 14
- c lg 0.25
- d lg 0.3
- e lg 54.6

UNDERSTAND AND USE THE LAWS OF LOGARITHMS

 $2^5 = 32$ and $\log_2 32 = 5$

The rules of logarithms follow the rules of indices.

| EXPONENT (POWERS) | LOGARITHMS | LAW | | |
|----------------------------|--|--------------------|--|--|
| $c^x \times c^x = c^{x+y}$ | $\log_c xy = \log_c x + \log_c y$ | Multiplication Law | | |
| $c^x \div c^y = c^{x-y}$ | $\log_c \frac{x}{y} = \log_c x - \log_c y$ | Division Law | | |
| $(c)^q$ | $\log_c(x^q) = q \log_c x$ | Power Law | | |
| $\frac{1}{c} = c^{-1}$ | $\log_c\left(\frac{1}{x}\right) = -\log_c x$ | | | |
| $c^1 = c$ | $\log_c(c) = 1$ | | | |
| $c^0 = 1$ | $\log_c(1) = 0$ | | | |

EXAMPLE 15

Write as a single logarithm

SKILLS

DECISION MAKING

- $a \log_3 6 + \log_3 7$
- **b** $\log_2 15 \log_2 3$ **c** $2 \log_5 3 + 3 \log_5 2$ **d** $\lg 3 4 \lg \left(\frac{1}{2}\right)$

- a $log_3(6 \times 7)$ $= log_3(42)$
- **b** $\log_2(15 \div 3)$ $= log_2 5$
- $c 2 \log_5 3 = +3 \log_5 2$ $= \log_5(3^2) = +\log_5(2^3)$ $= \log_5 9 + \log_5 8$ $= log_5 72$
- **d** $\lg 3 4 \lg \left(\frac{1}{2}\right)$ $= \lg 3 - \lg \left(\frac{1}{2}\right)^4$ $= \lg \left(3 \div \frac{1}{16}\right)$ = lg 48

Use the multiplication law

Use the division law

Apply the power law to both expressions

Use the multiplication law

Use the power law

Use the division law

EXAMPLE 16

Find the value of

Write in terms of $\log_a x$, $\log_a y$ and $\log_a z$

- a $\log_a(x^2yz^3)$
- **b** $\log_a\left(\frac{x}{u^3}\right)$
- $\operatorname{col}_a\left(\frac{x\sqrt{y}}{z}\right)$ $\operatorname{dol}_a\left(\frac{x}{a^4}\right)$

- a $\log_a(x^2yz^3)$
 - $= \log_a(x^2) + \log_a(y) + \log_a(z^3)$
 - $= 2\log_a(x) + \log_a(y) + 3\log_a(z)$
- **b** $\log_a\left(\frac{x}{u^3}\right)$
 - $= \log_a(x) \log_a(y^3)$
 - $= \log_a(x) 3\log_a(y)$
- $\log_a\left(\frac{x\sqrt{y}}{z}\right)$
 - $= \log_a(x\sqrt{y}) \log_a(z)$
 - $= \log_a(x) + \log_a(\sqrt{y}) \log_a(z)$
 - $= \log_a(x) + \frac{1}{2}\log_a(y) \log_a(z)$

Use the power law $\sqrt{y} = y^{\frac{1}{2}}$

- d $\log_a\left(\frac{x}{a^4}\right)$
 - $= \log_a(x) \log_a(a^4)$

$$= \log_a(x) - 4 \log_a(a)$$
$$= \log_a(x) - 4$$

$$\log_a a = 1$$

EXERCISE 7

Write as a single logarithm

SKILLS CRITICAL THINKING

$$a \log_4 8 + \log_4 8$$

$$\log_5 27 + \log_5 3$$

$$e 2 \log_6 9 - 10 \log_6 81$$

$$g \log_8 25 + \log_8 10 - 3\log_8 5$$

$$12 \log 20 - (\log 5 + \log 8)$$

$$\log_9 3 + \log_9 2$$

d
$$\log_4 24 + \log_4 15 - \log_5 3$$

$$\frac{1}{2}\log_2 25 + 2\log_2 3$$

h
$$2\log_{12}3 + 4\log_{12}2$$

2 Write in terms of $\log_a x$, $\log_a y$, $\log_a z$

a
$$\log_a x^4 y^3 z$$
 b $\log_a \frac{x^6}{y^3}$

$$b \log_a \frac{x^6}{y^3}$$

$$\log_a((xz)^2)$$

d
$$\log_a \frac{1}{xyz}$$

$$\log_a \sqrt{xy}$$

e
$$\log_a \sqrt{xy}$$
 f $\log_a \sqrt{x^4y^2z^3}$

g
$$\log_a \frac{\sqrt{x^3y^7}}{z^3}$$

CHANGE THE BASE OF A LOGARITHM

Working in base a, suppose that $\log_a x = m$

 $a^m = x$ Writing this as a power

Taking logs to a different base b $\log_b(a^m) = \log_b(x)$

Using the power law $m\log_b a = \log_b x$

 $\log_b x = \log_a x \times \log_b a$ Writing m as $\log_a x$

 $\log_a x = \frac{\log_b x}{\log_b a}$ This can be written as

Using this rule, notice in particular that $\log_a b = \frac{\log_b b}{\log_b a}$

but
$$\log_b b = 1$$

so,
$$\log_a b = \frac{1}{\log_b a}$$

EXAMPLE 17

Find, to 3 significant figures, the value of log₈(11)

SKILLS EXECUTIVE

One method is to use the change of base rule

$$\log_8 11 = \frac{\lg 11}{\lg 8}$$
$$= 1.15$$

Another method is to solve $8^x = 11$

$$Let x = log_8(11)$$

$$8^x = 11$$

$$\lg (8^x) = \lg 11$$

Take logs to base 10 of each side

$$x \log 8 = \log 11$$

Use the power law

$$x = \frac{\lg 11}{\lg 8}$$

Divide by Ig8

EXAMPLE 18

Solve the equation $\log_5 x + 6\log_x 5 = 5$

$$\log_5 x + \frac{6}{\log_5 x} = 5$$

Use change of base rule, special case

Let $\log_5(x) = y$

x = 1.15 (3 s.f.)

$$y+\frac{6}{y}=5$$

$$y^2 + 6 = 5y$$

Multiply by y

$$y^2 - 5y + 6 = 0$$

$$(y-3)(y-2)=0$$

So
$$y = 3$$
 or $y = 2$

$$\log_5 x = 3 \text{ or } \log_5 x = 2$$

$$x = 5^3 \text{ or } x = 5^2$$

Write as powers

$$x = 125 \text{ or } x = 25$$

EXERCISE 8

SKILLS

EXECUTIVE FUNCTION

Find, to 3 significant figures

a log₈785

 $\log_5 15$

 $c \log_6 32$

d log₁₂4

 $\log_{15} \frac{1}{7}$

Solve, giving your answer to 3 significant figures

a $6^x = 15$

b $9^x = 751$

 $c 15^x = 3$

 $3^x = 17.3$

 $3^{2x} = 25$

 $4^{3x} = 64$

$$\mathbf{g} \quad 7^{3x} = 152$$

HINT

If no base is given in a question, you should assume base 10.

Solve, giving your answer to 3 significant figures

 $\log_2 x = 8 + 9 \log_x 2$

 $\log_6 x + 3 \log_x 6 = 4$

 $\log x + 5 \log_x 10 = -6$

 $\log_2 x + \log_4 x = 2$

SOLVE EQUATIONS OF THE FORM $a^x = b$

You need to be able to solve equations of the form $a^x = b$

EXAMPLE 19

Solve the equation $3^x = 20$, giving your answer to 3 significant figures.

SKILLS

 $3^x = 20$

PROBLEM SOLVING **ANALYSIS**

 $\lg (3^x) = \lg 20$

Take logs to base 10 on each side

Use your calculator for logs to base 10

Use the power law

Divide by la 3

$$x \log 3 = \log 20$$

$$x = \frac{\lg 20}{\lg 3}$$

$$x = \frac{1.3010...}{0.4771...}$$

Or, a simpler version

$$3^x = 20$$

$$x = \log_3 20$$

$$x = 2.73$$

$$x = \log_3 20$$

$$x = 2.73$$

EXAMPLE 20

Solve the equation $7^{x+1} = 3^{x+2}$

$$(x + 1) \lg 7 = (x + 2) \lg 3$$

$$x \lg 7 + \lg 7 = x \lg 3 + 2 \lg 3$$

$$x \lg 7 - x \lg 3 = 2 \lg 3 - \lg 7$$

$$x(\lg 7 - \lg 3) = 2 \lg 3 - \lg 7$$

$$x = \frac{2\lg 3 - \lg 7}{\lg 7 - \lg 3}$$

$$x = 0.297$$
 (3 s.f.)

Use the power law

Multiply out

Collect *x* terms on left and numerical terms on right

Factorise

Divide by $\lg 7 - \lg 3$

EXAMPLE 21

Solve the equation $5^{2x} + 7(5^x) - 30 = 0$, giving your answer to 2 decimal places.

Let
$$y = 5^x$$

$$y^2 + 7y - 30 = 0$$

So
$$(y + 10)(y - 3) = 0$$

So
$$y = -10$$
 or $y = 3$

If
$$y = -10$$
, $5^x = -10$, has no solution

If
$$y = 3$$
, $5^x = 3$

$$lg(5^x) = lg 3$$

$$x \log (5) = \log 3$$

$$x = \frac{\lg 3}{\lg 5}$$

$$x = 0.683$$
 (3 s.f.)

Use
$$5^{2x} = (5^x)^2 = y^2$$

5^x cannot be negative

Solve as in previous examples

EXAM PRACTICE: CHAPTER 1

- Simplify $\sqrt{32} + \sqrt{18}$, giving your answer in the form $p\sqrt{2}$, where p is an integer. [2]
- Simplify $\frac{\sqrt{32} + \sqrt{18}}{3 + \sqrt{2}}$ giving your answer in the form $a\sqrt{2} + b$, where a and b are integers. [3]
- **a** Expand and simplify $(7 + \sqrt{5})(3 \sqrt{5})$ [2]
 - **b** Express $\frac{7+\sqrt{5}}{2+\sqrt{5}}$ in the form $a+b\sqrt{5}$, where a and b are integers. [2]
- Write $\sqrt{75} \sqrt{27}$ in the form $k\sqrt{x}$, where k and x are integers. [2]
- A rectangle A has a length of $(1 + \sqrt{5})$ cm and an area of $\sqrt{80}$ cm². Calculate the width of A in cm, giving your answer in the form $a + b\sqrt{5}$, where a and b are integers to be found. [3]
- Sketch the graph of $y = 8^x$, showing the coordinates of any points at which the graph crosses the axes. [3]
- Solve the equation $8^{2x} 4(8^x) = 3$, giving your answer to 3 significant figures. [3]
- a Given that $y = 6x^2$, Show that $\log_6 y = 1 + 2\log_6 x$ [2]
 - **b** Hence, or otherwise, solve the equation $1 + 2\log_3 x = \log_3(28x 9)$, giving your answer to 3 significant figures. [3]
- Find the values of x such that: $2 \log_3 x \log_3 (x-2) 2 = 0$ [2]
- Find the values of y such that: $\frac{\log_2 32 + \log_2 16}{\log_2 y} \log_2 y = 0$ [3]
- Given that $\log_b y + 3 \log_b 2 = 5$, express y in terms of b in its simplest form. [2]
- Solve $5^{2x} = 12(5^x) 35$ [4]
- Find, giving your answer to 3 significant figures where appropriate, the value of x for which $5^x = 10$ [3]
- Given that $\log_3(3b+1) \log_3(\alpha-2) = -1$, $\alpha > 2$, express b in terms of α . [2]
- Solve $3^{3x-2} = \sqrt[3]{9}$ [4]
- Solve $25^t + 5^{t+1} = 24$, giving your answer to 3 significant figures. [3]

c Hence, or otherwise, solve

- Given that $\log_2 x = p$, find, in terms of p, the simplest form of
 - a $\log_2(16x)$, [1]
 - b $\log_2\left(\frac{x^4}{2}\right)$. [1]
 - $\log_2(16x) \log_2\left(\frac{x^4}{2}\right) = \frac{1}{2}$

Give your answer in its simplest surd form. [3]

- Solve $\log_3 t + \log_3 5 = \log_3 (2t + 3)$ [3]
- a Draw the graph $y = 2 + \ln x$ for $0.1 \le x \le 4$ [2]
 - **b** Use your graph to estimate, to 2 significant figures, the solution to $\ln x = 0.5$, showing clearly your method. [2]
 - c By drawing a suitable line, estimate to 2 significant figures, the solution of the equation $x = e^{x-2}$ [2]

CHAPTER SUMMARY: CHAPTER 1

You can simplify expressions by using the power (indices) laws.

| $c^x \times c^x = c^{x+y}$ |
|----------------------------|
| $c^x \div c^y = c^{x-y}$ |
| $(c^p)^q = c^{p \times q}$ |
| $\frac{1}{c} = c^{-1}$ |
| $c^1 = c$ |
| $c^0 = 1$ |

- You can manipulate surds using these rules:

 - The rules for rationalising surds are:
 - If you have a fraction in the form $\frac{1}{\sqrt{a}}$ then multiply top and bottom by \sqrt{a}
 - If you have a fraction in the form $\frac{1}{1+\sqrt{a}}$ then multiply top and bottom by $(1-\sqrt{a})$
 - If you have a fraction in the form $\frac{1}{1-\sqrt{a}}$ then multiply top and bottom by $(1+\sqrt{a})$
- $\log_a n = x$ can be rewritten as $a^x = n$ where a is the base of the logarithm.
- The laws of logarithms are:

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \frac{x}{y} = \log_a x - \log_a y$$


$$\log_a (x^q) = q \log_a x$$

$$\log_a \left(\frac{1}{x}\right) = -\log_a x$$

$$\log_a (c) = 1$$

$$\log_a (1) = 0$$

- The change of base rule for logarithms can be written as $\log_a x = \frac{\log_b x}{\log_b a}$
- From the change of base you can derive $\log_a b = \frac{1}{\log_b a}$
- The natural logarithm is defined as: $\log_e x \equiv \ln x$
- The graph of $y = e^x$ is shown below.
 - $y = e^{x}$ $y = e^{x}$ y =
- The graph of $y = \ln x$ is shown below.



25

2 THE QUADRATIC FUNCTION

The path followed by a bottlenose dolphin jumping out of the water is called a parabola. A parabola is a visual realisation of the quadratic function $y = -x^2 + k$. Using this formula, scientists can calculate the height of a dolphin's jump (on the y-axis) and the distance travelled (on the x-axis).

There is no scientific agreement about why dolphins jump. Some scientists believe it is because they are trying to conserve energy. some believe it is to help them find food, and others believe they do it just for fun.

The parabola is a beautiful and elegant shape, commonly seen in nature. It is also seen in many man-made structures such as bridges and buildings.



LEARNING OBJECTIVES

- Factorise quadratic expressions where the coefficient of x^2 is greater than 1
- Complete the square and use this to solve quadratic equations
- Solve quadratic equations using the quadratic formula
- Understand and use the discriminant to identify whether the roots are (i) equal and real, (ii) unequal and real or (iii) not real
- Understand the roots a and β and know how to use them

 \circ 9 $qm^2 - 27m$

 $x^2 - 2x - 3$

STARTER ACTIVITIES

1 Factorise

$$6x^2 + 9x$$

d
$$9xy^2 + 36x^2y$$

$$b 2b^2 + 8b$$

$$= 24x - 64x^2$$

$$x^2 + 9x + 18$$

d
$$x^2 + 15x + 36$$

$$x^2 - 7x + 12$$

$$r^2 + 12r + 27$$

a
$$x^2 - 9$$

$$9x^2 - 16$$

$$r^2 - 7r + 1$$

$$x^2 + 12x + 27$$

All the parts in question 3 are examples of the difference of two squares.

b
$$x^2 - 25$$

d
$$25x^2 - 16$$

FACTORISE QUADRATIC EXPRESSIONS WHERE THE COEFFICIENT OF x^2 **IS GREATER THAN 1**

EXAMPLE 1

Factorise $3x^2 + 5x - 2$

SKILLS

DECISION MAKING CRITICAL THINKING You need to find two numbers:

They must **add** together to make +5 (the coefficient of x), and they must **multiply** together to give -6 (the coefficient of $x^2 \times$ the **constant** term, in this case 3×-2)

$$6 \times -1 = -6$$
 and $6 + (-1) = 5$

These numbers are then used to split the 5x into two terms, 6x and -1x

$$3x^2 + 5x - 2 = 3x^2 + 6x - x - 2$$
 The $5x$ term has been split into $6x$ and $-x$

$$= 3x(x + 2) - (x + 2)$$

$$= (x + 2)(3x - 1)$$
 $3x^2 + 6x$ and $-x - 2$ have both been factor

 $3x^2 + 6x$ and -x - 2 have both been factorised

EXAMPLE 2

Factorise $6x^2 - 5x - 4$

SKILLS

CRITICAL THINKING DECISION MAKING

Find two numbers that add together to make -5 and multiply together to give -24(the coefficient of x^2 multiplied by the constant term, in this case 6 \times -4)

$$3 \times -8 = -24$$
 and $3 + -8 = -5$
 $6x^2 - 5x - 4 = 6x^2 + 3x - 8x - 4$ The $-5x$ term has been split into $3x$ and $-8x$
 $= 3x(2x + 1) - 4(2x + 1)$
 $= (2x + 1)(3x - 4)$ $6x^2 + 3x$ and $-8x - 4$ have both been factorised

EXERCISE 1

SKILLS

CRITICAL THINKING DECISION MAKING

1 Factorise

a
$$3x^2 - 7x - 6$$

b
$$2x^2 + 11x + 5$$

c
$$2x^2 - 7x - 4$$

d
$$5x^2 - 16x + 3$$

$$6x^2 + x - 12$$

$$9x^2 - 6x + 1$$

$$9x^2 - 18x - 7$$

$$2x^2 + 5x + 2 = 0$$

b
$$4x^2 + 17x + 4 = 0$$

$$3x^2 - 13x = -4$$

$$6x^2 - 10x + 4 = 0$$

$$(x^2-2)=-17x$$

$$15x^2 + 42x - 9 = 0$$

$$3x(3x-4)=-4$$

COMPLETE THE SQUARE AND USE THIS TO SOLVE QUADRATIC EQUATIONS

A perfect square quadratic is in the form:

$$x^2 + 2bx + b^2 = (x + b)^2$$

$$x^2 - 2bx + b^2 = (x - b)^2$$

To turn a non-perfect square into a perfect square you need to manipulate the expression.

To complete the square of the function $x^2 + 2bx$ you need a further term b^2 .

So the completed square form is:

$$x^2 + 2bx = (x + b)^2 - b^2$$

Similarly

$$x^2 - 2bx = (x - b)^2 - b^2$$

Completing the square:
$$x^2 + bx = \left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2$$

EXAMPLE 3

Complete the square for

SKILLS

a
$$x^2 + 12x$$

b
$$2x^2 - 10x$$

REASONING EXECUTIVE FUNCTION

$$x^2 + 12x$$

$$= (x + 6)^2 - 6^2$$

$$= (x + 6)^2 - 36$$

b
$$2x^2 - 10x$$

$$=2(x^2-5x)$$

$$=2\left[\left(x-\frac{5}{2}\right)^2-\left(\frac{5}{2}\right)^2\right]$$

$$=2\left(x-\frac{5}{2}\right)^2-\frac{25}{4}$$

2b = 12 so b = 6

Here the coefficient of x^2 is 2

So take out the coefficient of x^2

Complete the square on $(x^2 - 5x)$

Any quadratic equation can be solved by completing the square.

EXAMPLE 4

Complete the square to solve

$$x^2 + 8x + 10 = 0$$

$$2x^2 - 8x + 7 = 0$$

$$x^2 + 8x + 10 = 0$$

$$x^2 + 8x = -10$$

$$(x + 4)^2 - 4^2 = -10$$

$$(x + 4)^2 = -10 + 16$$

$$(x + 4)^2 = 6$$

$$(x + 4) = \pm \sqrt{6}$$

$$x = -4 \pm \sqrt{6}$$

Check coefficient of $x^2 = 1$

Subtract 10 to get the LHS in the form $ax^2 + bx$

Complete the square for $(x^2 + 8x)$

Add 4² to both sides

Square root both sides

Subtract 4 from both sides

Then the solutions of $x^2 + 8x + 10 = 0$ are either

$$x = -4 + \sqrt{6}$$
 or $x = -4 - \sqrt{6}$

Leave your answer in surd form

b
$$2x^2 - 8x + 7 = 0$$

$$x^2 - 4x + \frac{7}{2} = 0$$

$$x^2 - 4x = -\frac{7}{2}$$

$$(x-2)^2-2^2=-\frac{7}{2}$$

$$(x-2)^2 = -\frac{7}{2} + 4$$

$$(x-2)^2 = \frac{1}{2}$$

$$(x-2)=\pm\sqrt{\frac{1}{2}}$$

$$x = 2 \pm \frac{1}{\sqrt{2}}$$

So the roots are either

$$x = 2 + \frac{1}{\sqrt{2}}$$
 or $x = 2 - \frac{1}{\sqrt{2}}$

The coefficient of
$$x^2$$
 is 2

So divide by 2

Subtract $\frac{7}{2}$ from both sides

Complete the square for $x^2 - 4x$

Add 2² to both sides

Combine the RHS

Square root both sides

Add 2 to both sides

SKILLS

PROBLEM

SOLVING

Complete the square for these expressions.

$$x^2 + 4x$$

b
$$x^2 - 16x$$

$$3x^2 - 24x$$

d
$$x^2 - x - 12$$

b
$$x^2 - 16x$$

e $x^2 + x - 1$

$$5x^2 - 6x + 1$$

$$\mathbf{g} \ 2x^2 + 3x - 1$$

h
$$4x^2 + 6x - 1$$

Solve these quadratic equations by completing the square.

Leave your answer in surd form where appropriate.

a
$$6x^2 - 11x - 10 = 0$$

$$x^2 + 2x - 2 = 0$$

$$2r^2 - 6r + 1 = 0$$

$$\frac{1}{2}$$
 $2r^2 + 3r - 6 = 0$

b
$$x^2 + 2x - 2 = 0$$
 c $2x^2 - 6x + 1 = 0$ **e** $4x^2 - 59x - 15 = 0$ **f** $4x^2 + 8x - 9 = 0$

$$1x^2 + 8x - 9 = 0$$

d
$$2x^2 + 3x - 6 = 0$$

e $4x^2 - 59x - 15$
g $15 - 6x - 2x^2 = 0$
h $4x^2 - x - 8 = 0$

h
$$4x^2 - x - 8 = 0$$

3 Now by completing the square that the solutions to $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2}$

SOLVE QUADRATIC EQUATIONS USING THE QUADRATIC FORMULA

The quadratic formula $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2}$

can be used to solve any quadratic equation of the form $ax^2 + bx + c = 0$

EXAMPLE 5

Use the quadratic formula to solve $4x^2 - 3x - 2 = 0$

SKILLS

PROBLEM SOLVING

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(4)(-2)}}{2 \times 4}$$

$$x = \frac{3 \pm \sqrt{(9 + 32)}}{8}$$

$$x = \frac{3 \pm \sqrt{41}}{8}$$

Then
$$x = \frac{3 + \sqrt{41}}{8}$$
 or $x = \frac{3 - \sqrt{41}}{8}$

$$-4 \times 4 \times -2 = 32$$

Leave your answer in surd form

Use $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, where a = 4, b = -3, c = -2

EXERCISE 3

SKILLS EXECUTIVE FUNCTION

1 Solve these equations using the quadratic formula. Leave your answer in surd form where appropriate.

$$x^2 = 6x$$

$$2x^2 + 50x = 0$$

b
$$2x^2 + 50x = 0$$
 c $x^2 - x - 6 = 0$

$$p^2 - 4p + 2 = 0$$

$$m^2 + 2m - 2 = 0$$

d
$$p^2 - 4p + 2 = 0$$
 e $m^2 + 2m - 2 = 0$ **f** $x^2 + 7x + 10 = 0$

$$t^2 - 5t - 6 = 0$$

h
$$x^2 + 2x - 35 = 0$$
 i $n^2 - 4n + 4 = 0$

$$n^2 - 4n + 4 = 0$$

$$x^2 + 6x + 6 = 0$$

2 Solve these equations, leaving your answer in surd form.

$$9x^2 - 6x - 25 = 0$$

b
$$3x^2 - 6x + 2 = 0$$

c
$$6x^2 - 5x - 6 = 0$$

$$2x^2 + 3x - 1 = 0$$

$$2x^2 + 7x + 3 = 0$$

$$3x^2 + 7x + 1 = 0$$

$$4x^2 - 16x + 15 = 0$$

h
$$7x^2 + 5x - 3 = 0$$


$$10x^2 - 15x - 8 = 0$$

$$x^2 + 3x - 6 = 0$$

$$(x-7)^2=63$$

$$4x^2 + 15x + 13 = 0$$

3 The diagram shows the floor plan of a bedroom. The total area is $35.5 \,\mathrm{m}^2$ Find the value of x.



UNDERSTAND AND USE THE DISCRIMINANT TO IDENTIFY WHETHER THE ROOTS ARE (i) EQUAL AND REAL, (ii) UNEQUAL AND REAL OR (iii) NOT REAL

The equation $ax^2 + bx + c = 0$ has two solutions

$$x = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$
 and $x = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$

These two solutions may be the same or different. real numbers or not real numbers.

The nature of the roots (solutions) of the equation will clearly depend on the expression $b^2 - 4ac$. This expression $b^2 - 4ac$ is called the *discriminant*, as it allows us to identify (discriminate) whether the roots of a particular equation are equal and real, unequal and real or not real at all.

lf

- $b^2 4ac > 0$ the roots of the equation are **unequal and real (two** roots)
- $b^2 4ac = 0$ the roots of the equation are equal and real (one root)
- $b^2 4ac < 0$ there are **no real roots** of the equation (**no** real roots)

EXAMPLE 6

What can you deduce from the values of the discriminants of these equations? Find the roots where possible.

SKILLS REASONING

$$2x^2 - 2x + 5 = 0$$

$$2x^2 - x - 1 - 0$$

a
$$2x^2 - 3x + 5 = 0$$
 b $3x^2 - x - 1 = 0$ **c** $4x^2 - 12x + 9 = 0$

$$2x^2 - 3x + 5 = 0$$

$$a = 2, b = -3, c = 5$$

$$b^{2} - 4ac$$

$$(-3)^2 - 4 \times 2 \times 5 = -31$$

Therefore there are no real roots.

b
$$3x^2 - x - 1 = 0$$

$$a = 3$$
, $b = -1$, $c = -1$

$$b^2 - 4ac = (-1)^2 - 4 \times 3 \times (-1) = 13$$

Therefore there are two unequal roots.

$$x = \frac{-(-1) \pm \sqrt{13}}{2 \times 3}$$
 so

$$x = 0.768, -0.434$$

$$4x^2 - 12x + 9 = 0$$

$$a = 4, b = -12, c = 9$$

$$(-12)^2 - 4 \times 4 \times 9 = 0$$

Calculate the discriminant

Therefore the roots are real and equal.

$$x = \frac{-(-12)}{2 \times 4}$$
 and $x = \frac{3}{2}$

When the discriminant is 0 you can always factorise. In this case $4x^2 - 12x + 9 = (2x - 3)^2 = 0$

Use the quadratic formula to find solutions

EXAMPLE 7

The equation $kx^2 - 2x - 8 = 0$ has two real roots.

What can you deduce about the value of the constant k?

Since the equation has two real roots, you know that the discriminant $b^2 - 4ac$ must be greater than zero.

You substitute a = k, b = -2 and c = -8 into the inequality $b^2 - 4ac > 0$, giving

$$(-2)^2 - 4 \times k \times -8 > 0$$

$$4 + 32k > 0$$

$$k > -\frac{4}{32}$$

$$k > -\frac{1}{8}$$

EXERCISE 4

SKILLS

REASONING

1 b Use the discriminant to **determine** whether these equations have one root, two roots or no roots.

a
$$x^2 - 2x + 1 = 0$$
 b $x^2 - 3x - 2 = 0$

$$r^2 - 3r - 2 = 0$$

$$2x^2-3x-4=0$$

$$2x^2 - 4x + 5 = 0$$

$$2r^2 - 4r + 2 = 0$$

e
$$2x^2 - 4x + 2 = 0$$
 f $2x^2 - 7x + 3 = 0$

$$3x^2 - 6x + 5 = 0$$

$$7x^2 - 144x + 57 = 0$$

$$16x^2 - 2x + 3 = 0$$

$$x^2 + 22x + 121 = 0$$

$$\mathbf{k} \ 5x^2 - 4x + 81 = 0$$

- The equation $px^2 2x 7 = 0$ has two real roots. What can you deduce about the value of p?
- **3** The equation $3x^2 + 2x + m = 0$ has equal roots. Find the value of m.

UNDERSTAND THE ROOTS a AND β AND HOW TO USE THEM

If α and β are the **roots** of the equation $ax^2 + bx + c = 0$ then you deduce that

$$(x-a)(x-\beta)=0$$

You can rewrite this as $x^2 - x(\alpha + \beta) + \alpha\beta = 0$

Comparing this with $ax^2 + bx + c = 0$ you can see that

$$\alpha + \beta = \frac{-b}{a}$$
 and $\alpha\beta = \frac{c}{a}$

KEY POINTS

For the equation $ax^2 + bx + c = 0$

- The sum of roots, $\alpha + \beta = \frac{-b}{a}$
- The product of the roots, $a\beta = \frac{c}{a}$

EXAMPLE 8

1 The roots of the equation $3x^2 + x - 6 = 0$ are α and β .

SKILLS

REASONING CRITICAL THINKING

- **a** Find an expression for $\alpha + \beta$ and an expression for $\alpha\beta$.
- **b** Hence find an expression for $a^2 + \beta^2$ and an expression for $a^2\beta^2$.
- Find a quadratic equation with roots α^2 and β^2 .

$$3x^2 + x - 6 = 0$$

$$x^2 + \frac{1}{3}x - 2 = 0$$

Therefore sum of the roots $\alpha + \beta = -\frac{1}{3}$

Note: Sometimes you will need to manipulate the expressions to help you solve the questions.

Product of the roots $a\beta = -2$

b
$$(a + \beta)^2 = a^2 + 2a\beta + \beta^2$$

Therefore
$$a^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$$

Substituting the results from part a, gives

$$a^2 + \beta^2 = \left(-\frac{1}{3}\right)^2 - 2(-2)$$

$$=\frac{37}{9}$$

$$\alpha^2 \beta^2 = (\alpha \beta)^2 = (-2)^2 = 4$$

Divide the equation by 3 to obtain an equation where the coefficient of x^2 is 1

The sum of roots $a + \beta = -\frac{b}{a}$

The product of the roots $a\beta = \frac{c}{a}$

c Let the equation be $x^2 + px + q = 0$

$$p = (\alpha^2 + \beta^2) = -\frac{37}{9}$$

$$q = \alpha^2 \beta^2 = 4$$

So the equation is: $x^2 - \frac{37}{9}x + 4 = 0$

OR
$$9x^2 - 37x + 36 = 0$$

Use p = -sum of the roots

q =product of the roots

Simplify the equations in order for the all the coefficients to be integers

Note: This question can be answered without finding α and β .

EXAMPLE 9

The roots of the equation $x^2 - 3x - 2 = 0$ are α and β .

Without finding the value of α and β , find the equations with the roots

b
$$\frac{1}{a}$$
, $\frac{1}{\beta}$

$$c$$
 a^2, β^2

If α and β are the roots of $x^2 - 3x - 2 = 0$ then $\alpha + \beta = 3$ and $\alpha\beta = -2$

a If the roots are 3α and 3β then

Sum of the roots =
$$3(\alpha + \beta) = 3 \times 3 = 9$$

Using
$$\alpha + \beta = 3$$
 and $\alpha\beta = -2$

Using $\alpha + \beta = 3$ and $\alpha\beta = -2$

Product of the roots =
$$3\alpha \times 3\beta = 9\alpha\beta = -18$$

Equation is
$$x^2 - 9x - 18 = 0$$

b If the roots are $\frac{1}{a}$, $\frac{1}{8}$

Sum of roots
$$\frac{1}{a} + \frac{1}{\beta} = \frac{\beta + \alpha}{\alpha\beta}$$
$$= -\frac{3}{2}$$

Product of roots:
$$\frac{1}{\alpha} \times \frac{1}{\beta} = \frac{1}{\alpha\beta}$$

$$=-\frac{1}{2}$$

Equation is:
$$x^2 + \frac{3}{2}x - \frac{1}{2} = 0$$

so
$$2x^2 + 3x - 1 = 0$$

c If the roots are α^2 , β^2 then

Sum of roots =
$$a^2 + \beta^2 = (a + \beta)^2 - 2a\beta$$

= $3^2 - 2(-2) = 13$

Product of roots =
$$\alpha^2 \beta^2 = \alpha \beta^2 = (-2)^2 = 4$$

Therefore equation is
$$x^2 - 13x + 4 = 0$$

EXERCISE 5

SKILLS

REASONING CRITICAL THINKING

- 1 The roots of the equation $x^2 + 5x + 2 = 0$ are α and β . Find an equation whose roots are
 - a $2\alpha + 1$ and $2\beta + 1$
 - **b** $\alpha\beta$ and $\alpha^2\beta^2$
- **2** The roots of the equation $x^2 + 6x + 1 = 0$ are α and β . Find an equation whose roots are
 - a $\alpha + 3$ and $\beta + 3$
 - **b** $\frac{a}{\beta}$ and $\frac{\beta}{a}$
- **3** The roots of the equation $x^2 x 1 = 0$ are α and β . Find an equation whose roots are
 - a $\frac{1}{a}$ and $\frac{1}{\beta}$
 - **b** $\frac{\alpha}{\alpha + \beta}$ and $\frac{\beta}{\alpha + \beta}$

EXAM PRACTICE: CHAPTER 2

- $f(x) = 0 = 3x^2 10x 2$
 - a Without solving the equation f(x)=0, form an equation, with integer coefficients
 - i roots $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$
 - ii roots $2\alpha + \beta$ and $\alpha = 2\beta$ [6]

[4]

[2]

- **b** Solve f(x) = 0 using completing the square.
- The roots of a quadratic equation are α and β where $\alpha + \beta = -\frac{9}{5}$ and $\alpha\beta = -3$. Find a quadratic equation, with integer coefficients, which has roots α and β [4]
- Given that $\alpha + \beta = 7$ and $\alpha^2 + \beta^2 = 25$
 - a Show that $\alpha\beta = 12$. [2]
 - b Hence, or otherwise, form a quadratic equation with the integer coefficients, which has roots α and β . [3]
 - Form a quadratic equation, with integer coefficients, which ahas roots $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$. [5]
- The equation $x^2 + (p-3)x + (3-2p) = 0$, where p is a constant, has two distinct
 - a Show that p satisfies $p^2 + 2p 3 > 0$ [1]
 - **b** Find the possible values of *p*.
- a Show that $x^2 + 6x + 11$ can be written as $(x + a)^2 + b$. [2]
 - **b** Find the value of the discriminant. [2]
- Factorise completely
 - $5x^2 + 16x + 3$ [3]
 - **b** $3x^2 7x + 4$ [3]
- Solve these equations by completing the square.
 - $p^2 + 3p + 2 = 0$ [3]
 - **b** $3x^2 + 13x 10 = 0$ [3]
- Solve these equations by using the quadratic formula.
 - $5x^2 + 3x 1 = 0$ [2]
 - **b** $(2x-5)^2=7$ [3]
- $4x 5 x^2 = b (x + a)^2$ where a and b are integers.
 - **a** Find the value of a and b. [2]
- b Calculate the discriminant. [2] Solve $\frac{4}{2x+1} - 3 = -\frac{1}{4x^2-1}$ [3]

CHAPTER SUMMARY: CHAPTER 2

- $x^2 y^2 = (x y)(x + y)$ is known as the difference of two squares.
- Quadratic equations can be solved by
 - factorisation
 - \blacksquare completing the square: $x^2 + bx = \left(x + \frac{b}{2}\right)^2 \left(\frac{b}{2}\right)^2$
 - using the quadratic formula: $x = \frac{-b \pm \sqrt{b^2 4ac}}{2a}$
- The *discriminant* of a quadratic expression is $b^2 4ac$
- If α and β are the roots of the equation $\alpha x^2 + bx + c = 0$ then
 - $\blacksquare \ \alpha + \beta = -\frac{b}{a}$
 - \blacksquare $\alpha\beta = \frac{c}{a}$

CHAPTER 3



CHAPTER 3


3 INEQUALITIES AND IDENTITIES

The only value at which Fahrenheit and Celsius are the same temperature is -40.

If you substitute F = C into the **linear** equation F = 1.8C + 32you get: C = F = -40

In this chapter you will learn about simultaneous equations, inequalities and linear programming (a form of mathematical modelling).

All of these mathematical tools are vital for research into global warming. Mathematical modelling enables us to predict the effects of global warming, from how it will affect human health to the effect it has on global food production. Scientists. businesses, economists and politicians all use mathematical models to plan ways of limiting the consequences of global warming.



LEARNING OBJECTIVES

- Solve two simultaneous equations, one linear and one quadratic
- Solve linear inequalities
- Solve quadratic inequalities
- **Graph** linear inequalities in two variables

- Use inequalities to solve linear programming problems
- Divide a **polynomial** by $(x \pm p)$
- Factorise a polynomial by using the factor theorem
- Use the remainder theorem to find the remainder when a polynomial is divided by (ax - b)

STARTER ACTIVITIES

1 Using substitution, solve these simultaneous equations.

a
$$x + y = 15$$
 $x - y = 3$

$$x - y = 3$$

b
$$2x - 3y = 2$$
 $x + 2y = 8$

$$x + 2u = 8$$

$$x + y = 7$$
 $2x - 3y = 9$

$$2x - 3y = 9$$

d
$$11y + 15x = -23$$
 $7y - 2x = 20$

$$7u - 2x = 20$$

2 > Sketch these equations on one graph.

a
$$y = x^2$$

b
$$y = 2x^2$$

$$y = -x^2$$

$$y = 5x^2 + 2$$

$$y = -3x^2 - 4$$

3 Use long division to solve these equations.

SOLVE SIMULTANEOUS EQUATIONS, ONE LINEAR AND ONE QUADRATIC

EXAMPLE 1

Solve

a
$$x + 2y = 3$$

 $x^2 + 3xy = 10$

b
$$3x - 2y = 1$$

 $x^2 + y^2 = 25$

$$x = 3 - 2y$$

$$(3-2y)^2+3y(3-2y)=10$$

$$9 - 12y + 4y^2 + 9y - 6y^2 = 10$$
$$-2y^2 - 3y - 1 = 0$$

$$2y^2 + 3y + 1 = 0$$

$$(2y + 1)(y + 1) = 0$$

Rearrange the linear equation to get $x=\dots$ or $y=\dots$

Substitute this into the quadratic equation, in place of x

$$(3-2y)^2 = (3-2y) \times (3-2y)$$

Solve for *y* using factorisation

$$y = -\frac{1}{2} \text{ or } y = -1$$

So
$$x = 4$$
 or $x = 5$

Solutions are
$$x = 4$$
, $y = -\frac{1}{2}$ and $x = 5$, $y = -1$.

There are two solution pairs. The graph of the linear equation (straight line) intersects the graph of the quadratic (curve) at two points.

b
$$3x - 2y = 1$$

 $x^2 + y^2 = 25$

$$2y = 3x - 1$$
$$y = \frac{3x - 1}{2}$$

$$x^2 + \left(\frac{3x-1}{2}\right)^2 = 25$$

$$x^{2} + \left(\frac{9x^{2} - 6x + 1}{4}\right) = 25$$
$$4x^{2} + 9x^{2} - 6x + 1 = 100$$

$$13x^2 - 6x - 99 = 0$$

$$(13x + 33)(x - 3) = 0$$

So
$$x = -\frac{33}{13}$$
 or $x = 3$

$$y = -\frac{56}{13}$$
 or $y = 4$

Find y = ... from linear equation

Substitute $y = \frac{3x-1}{2}$ into the quadratic equation to form an equation in x

Now multiply by 4

Substitute *x*-values into
$$y = \frac{3x - 1}{2}$$

The solutions are
$$x = 3$$
, $y = 4$ and $x = -\frac{33}{13}$, $y = -\frac{56}{13}$

b $2x + 1 = y, x^2 - x + 3 - y = 0$

d $y = x^2 + 2x - 3$, y = 2x + 1

 $x = 3 + 2y, x^2 + 2y^2 = 27$

h 2y + 4x = 6, $2x^2 - 3xy = 14$

EXERCISE 1

SKILLS

PROBLEM SOLVING 1 Solve these simultaneous equations.

$$3 - x^2 = y, x + 1 = y$$

$$x^2 + y^2 = 5, x + y = 3$$

$$y = x^2 - 3x + 1, y = 2x - 5$$

$$2x^2 + xy + y^2 = 22, x + y = 1$$

$$2x + 4y = 6$$
, $x^2 - 2y + 4y^2 = 18$

$$xy = 14, 4x + 3y = 29$$

$$x^2 + 4y^2 + 3x + 2y - 56 = 0, 5x - 2y + 7 = 0$$

$$2x + 4y = 9, 4x^2 + 16y^2 - 20x - 4y + 19 = 0$$

m Find the coordinates of the point where the line 2x + 2y = 18 meets the circle $(x - 2)^2 + (y - 3)^2 = 16$

2 Find the coordinates of the intersection of the given curves in each case.

$$y = 4x^2 + 11x$$

$$y = x^3 + 3x^2 - x$$

b
$$y = x^3 + 3x$$

$$y=x^2+5x$$

$$y + 8x^2 = 2x^3$$

$$y + x^2 + 3x = 0$$

d
$$y = 2x^2 - x$$

$$y = 2x^3 + 3x^2 - 7x$$

SOLVE LINEAR INEQUALITIES

You can use the skills learnt in solving linear equations to solve linear inequalities.

EXAMPLE 2

Find the set of values of x for which

a
$$2x - 5 < 7$$

b
$$5x + 9 > x + 20$$

c
$$12 - 3x < 27$$

d
$$3(x-5) > 5-2(x-8)$$

a
$$2x - 5 < 7$$

Add 5 to both sides

Divide both sides by 2

b
$$5x + 9 > x + 20$$

$$4x + 9 > 20$$

/ 20

Subtract *x* from both sides
Subtract 9 from both sides

$$x > \frac{11}{4}$$

Divide both sides by 4

c
$$12 - 3x < 27$$

$$-3x < 15$$

Subtract 12 from both sides

$$x > -5$$

Divide both sides by -3. You therefore need to reverse the inequality sign

Alternative approach

$$12 - 3x < 27$$

$$12 < 27 + 3x$$

Add 3x to both sides

$$-15 < 3x$$

Subtract 27 from both sides

$$-5 < x$$

Divide both sides by 3

$$x > -5$$

Rewrite with x on the LHS

d
$$3(x-5) > 5-2(x-8)$$

$$3x - 15 > 5 - 2x + 16$$

Expand

$$5x > 5 + 16 + 15$$

Add 2x + 15 to both sides

$$x > \frac{36}{5}$$

Divide both sides by 5

EXAMPLE 3

Find the set of values of x for which

$$3x - 5 < x + 8$$
 and $5x > x - 8$

$$3x - 5 < x + 8$$
 gives

$$2x - 5 < 8$$

$$x < \frac{13}{2}$$

$$5x > x - 8$$
 gives

$$4x > -8$$

$$x > -2$$

Draw a number line to illustrate the two inequalities.

So the required set of values is $-2 < x < \frac{13}{2}$

Note: Draw a number line.

EXAMPLE 4

Find the set of values of x for which 4x + 7 > 3 and 17 < 11 + 2x

$$4x + 7 > 3$$
 gives


$$4x > -4$$

$$x > -1$$

$$17 < 11 + 2x$$
 gives

Note: Draw a number line.

So the required set of values is x > 3



EXERCISE 2

SKILLS REASONING

1 Find the set of values of x for which

$$5x - 8 < 6 - 2x$$

c
$$15x + 3 \le 27x - 3$$

$$= 85x + 15 > 7 + 3x$$

$$9x + 71 \le 16 + 8x$$

$$2(3x + 5) \ge 4x + 3$$

$$4(x+7) > 3(x+2)$$

$$e$$
 $4(4x - 3) \le 6(7x - 3)$

$$16(4-2x) \le 7(3-x)$$

b
$$7x - 17 > 7 - 7x$$

d
$$16x - 3 \ge 3 - 2x$$

$$\mathbf{f}$$
 16x + 25 < 27 + 3x

h
$$2x + 69 \ge -3x + 52$$

b
$$3(5x + 15) < 6x$$

d
$$5(3x-6) \ge 7(5x+2)$$

f
$$22(3x + 8) > 3(5 - x)$$

$$-7(3x + 3) > -6(4 - x)$$

3
$$\triangleright$$
 Find the set of values of x for which

a
$$3(x-2) > x-4$$
 and $4x+12 > 2x+17$

b
$$4(x + 3) > 2$$
 and $5x + 5 > 6$

d
$$3(3+2x) < 2(4x+1)$$
 and $6(2x+3) < 3(x-4)$

$$9(2x-3) \ge 4$$
 and $15(6x+3) \ge 5$

f
$$4(2-x) > 3(3x+3)$$
 and $6(x-2) < 4(2x+2)$

SOLVE QUADRATIC INEQUALITIES

To solve quadratic inequalities you need to

- Solve the quadratic equation
- ii Sketch the corresponding quadratic function
- iii Use your sketch to find the required set of values.

HINT

To solve the quadratic inequality, you may have to rearrange it so that all the terms are on one side.

EXAMPLE 5

Find the set of values of x for which $x^2 - 4x - 5 < 0$ and draw a sketch to show this.


SKILLS

INTERPRETATION THINKING

$$x^2 - 4x - 5 = 0$$

$$(x + 1)(x - 5) = 0$$

$$x = -1 \text{ or } x = 5$$



Quadratic equation

Factorise

-1 and 5 are called critical values

Your sketch does not need to be accurate. All you really need to know is that the graph is 'U-shaped' and crosses the x-axis at -1 and 5

$$x^2 - 4x - 5 < 0$$
 ($y < 0$) for the part of the graph below the x-axis.

So the required set of values is -1 < x < 5

KEY POINTS

Be careful how you write your answers

-1 < x < 5 is fine, showing that x is between -1 and 5.

But it is wrong to write something like 5 < x < -1 or -1 > x > 5 because x cannot be less than −1 and greater than 5 at the same time.

This type of solution needs to be written in two separate parts, x < -1, x > 5.

EXAMPLE 6

Find the set of values of x for which $3-5x-2x^2<0$ and sketch the graph of $y=3-5x-2x^2$

$$3 - 5x - 2x^2 = 0$$

$$2x^2 + 5x - 3 = 0$$


$$(2x-1)(x+3)=0$$

$$x = \frac{1}{2} \text{ or } x = -3$$

Quadratic equation

Multiply by -1, so it's easier to factorise

$$\frac{1}{2}$$
 and -3 are the critical values



Since the coefficient of x^2 is negative, the graph is 'upside-down U- shaped', and crosses the x-axis at -3 and $\frac{1}{2}$

 $3-5x-2x^2<0$ (y<0) for the outer parts of the graph, below the x-axis

So the required set of vales is x < -3 or $x > \frac{1}{2}$

EXAMPLE 7

Find the set of values of x for which $12 + 4x > x^2$

Method 1: sketch graph

$$12 + 4x > x^2$$

$$12 + 4x - x^2 > 0$$

$$x^2 - 4x - 12 = 0$$


$$(x + 2)(x - 6) = 0$$

$$x = -2 \text{ or } x = 6$$

Sketch of
$$y = 12 + 4x - x^2$$

$$12 + 4x - x^2 > 0$$

Therefore -2 < x < 6



Method 2: table

$$12 + 4x > x^2$$

$$12 + 4x - x^2 < 0$$

$$x^2 - 4x - 12 < 0$$

$$x^2 - 4x - 12 = 0$$

$$(x + 2)(x - 6) = 0$$

$$x = -2 \text{ or } x = 6$$

| | x < -2 | -2 < x < 6 | <i>x</i> > 6 | 4 |
|----------------|----------|------------|--------------|---|
| x + 2 | _ | + | + | |
| x-6 | _ | _ | + | - |
| (x + 2)(x - 6) | + | _ | + | |
| | A | | | |

Use the critical values to split the real number line into sets

For each set, check whether the set of values makes the value of the bracket positive or negative

For example, if x < -2, x + 2 is negative, x - 6 is negative, and (x + 2)(x - 6) is positive

$$x^2 - 4x - 12 < 0$$

$$(x + 2)(x - 6) < 0$$

$$(x + 2)(x - 6)$$
 is negative for $-2 < x < 6$

Therefore -2 < x < 6

EXERCISE 3

SKILLS INTERPRETATION

CRITICAL

1 > By sketching graphs, solve these inequalities.

$$(x-2)(x+3)<0$$

b
$$(x + 3)(x - 4) \le 0$$

$$(2x + 3)(x + 4) < 0$$

d
$$(7x + 8)(4 - 6x) > 0$$

$$(6x + 3)(4x - 3) \ge 0$$

f
$$(4x-3)(5x-7) \le 0$$

2 Find the set of values of x which satisfy these inequalities.

$$x^2 - 7x - 18 < 0$$

b
$$x^2 + 6x + 9 \le 0$$

$$x^2 + 13x + 12 \ge 0$$

d
$$x^2 - 14x + 40 < 0$$

$$x^2 + 7x + 12 > 0$$

$$f 7x^2 - 31x - 20 \le 0$$

$$2x^2 + 17x + 21 > 0$$

h
$$5x^2 - x - 18 < 0$$

$$6x^2 + 14x - 12 > 0$$

$$x^2 - 25 \le 0$$

$$x^2 - 16 \ge 0$$

HINT

If the line is to be included in the region you use \geq or \leq in place of > or \leq .


So points in region A, including the line, would be represented by: $ax + by \ge c$ or $y \ge c$


mx + c

To decide which side of the line does not satisfy the inequality, choose a convenient point in each region and substitute its coordinates into the inequality to see whether (or not) it satisfies the inequality. Shade the region containing the point that does not satisfy the inequality.

GRAPH LINEAR INEQUALITIES IN TWO VARIABLES

Consider the two diagrams below.





All the points that lie on the line are represented by

$$ax + by = c \text{ or } y = mx + c$$


- All the points that lie above the line, in region A, are represented by ax + by > c or y > mx + c
- All the points that lie below the line, in region B, are represented by ax + by < c or y < mx + c

When illustrating an inequality, first draw a straight line and then use shading.

- Inequalities using > or < are represented by a **dashed** (or **dotted**) **line**. This indicates the the **line itself** is *not* **included in the region**.
- Inequalities using ≥ or ≤ are represented by a solid line to show that the line is included in the region.


The convention is that the region that does *not* satisfy the inequality is shaded.

The feasible region satisfying several inequalities will therefore be the one left unshaded.




EXAMPLE 8

SKILLS ANALYSIS Write down the inequalities shown by regions A, B, C, D, E, F, G, H, I and J in the diagrams below.



Note: x = 3y - 6 could be written in many different forms, such as 3y = x + 6 or 3y - x = 6

You must use a coordinate to check the direction of the inequality since the region, I, is given by $x \le 3y - 6$ or equivalent.



EXAMPLE 9


Illustrate on a diagram the region R, for which:

$$x \ge 1$$
 $4x + 3y < 12$ $2y \le x$ $x, y \ge 0$ Label the region R .

There are five inequalities here and three lines to add to the axes.

$$x \ge 1$$

Draw x = 1




$$4x + 3y < 12$$

Draw 4x + 3y = 12 as a broken line.


To draw it, note that when x = 0, y = 4

When
$$y = 0$$
, $x = 3$




$$2y \le x$$


To draw the line 2y = x, note that it passes through (0, 0) and (2, 1)



Also draw x, y > 0



Combine all these on one diagram:



47

Note: Colour has been used here for clarity. Do not use colour in the examination!

EXAMPLE 10

SKILLS

CRITICAL THINKING REASONING ANALYSIS

Note: This is an example of Linear Programming.

A food cart manager sells cheeseburgers and chicken burgers. The cart is open only during the lunch break of 12-2 pm. The manager can cook a maximum of 40 cheeseburgers and a maximum of 70 chicken burgers. He cannot cook more than 90 burgers in total. The profit on a cheeseburger is 33 pence and the profit on a chicken burger is 21 pence. How many of each kind of burger should the manager sell to maximise profit?



Step 1

Formulate the problems into linear equations.

Let x = number of cheeseburgers

Let y = number of chicken burgers

x > 0 and y > 0

You cannot have a negative number of burgers

Number of each burger

 $x \le 40$ and $y \le 70$

These equations are created from information in the question

Total number of burgers

$$x + y \leq 90$$

Step 2


Obtain a profit inequality

0.33(x) + 0.21(y) = profit

Convert pence to pounds

Step 3

Now draw these equations on a graph.



The region shaded in red is the only region that satisfies all the inequalities. Any value in this region will work but the maximum profit value will be close to one of the vertices of the regions marked A, B, C, D.

Step 4

Find the coordinates of the vertices:

The coordinates of A are (0, 70).

The coordinates of B are (20, 70).

The coordinates of C are (40, 50).

The coordinates of D are (40, 0).

Step 5

Input these coordinates into profit inequality found in step 2 (0.33(x) + 0.21(y) = profit):

For A
$$0.33(0) + 0.21(70) = 14.70$$

For B
$$0.33(20) + 0.21(70) = 21.30$$

For C
$$0.33(40) + 0.21(50) = 23.70$$

For **D**
$$0.33(40) + 0.21(0) = 13.20$$


Therefore the maximum possible profit is £23.70.


For this the manager will need to sell 40 cheeseburgers and 50 chicken burgers.

EXERCISE 4


For questions 1-2 describe the shaded region.


SKILLS CRITICAL THINKING REASONING **ANALYSIS**








For questions 3–8, define the region labelled R.











For questions 9–12, represent the region defined by each set of inequalities, by drawing lines and shading.

- 8 $> y \ge 2x, x + 2y \le 4, y + 2x > 1$
- 9 \triangleright 2y > x, y + 2x \leq 4, y > 2x + 2
- **10** $x \ge 0, y \ge 0, y < \frac{x}{2} + 4, y \le 6 2x$
- **11** $x > 0, y \ge 0, 3x + 4y \le 12, 5x + 2y \le 10$
- Li is buying some chickens and lambs for his farm. He buys c chickens at Y120 each. He buys l lambs at Y200 each. He wants at least 10 animals in total. He wants more lambs than chickens. He has a maximum of Y1800 to spend.
 - a Write down three inequalities involving c and l.
 - Li can sell the produce of each chicken for Y150 and the produce of each lamb for Y450.
 How many of each should he buy to maximise his profit? Use a graphical approach.



- Tala is a semi-professional photographer in her spare time. As a wedding photographer she can earn 10 Jordanian Dinars (JOD). As a studio photographer she can earn 8 JOD an hour. She wants to spend at least 6 hours as a studio photographer. She can only work for a maximum of 20 hours a week. Obtain inequalities and graph your solution to find her maximum possible weekly earnings.
- 14 A firm is planning to produce two types of light fittings, type A and type B.

Market research suggests that, each week, at least 50 type A light fittings should be produced.

The number of type A light fittings should be between 20% and 40% of the total number of light fittings produced.

Each type A light fitting requires 3 light bulbs and each type B light fitting requires 2 light bulbs. The firm can only buy 200 light bulbs each week.



The profit on each type A light fitting is £15. The profit on each type B light fitting is £12.

The firm wants to maximise its weekly profit.

Formulate this situation as a linear programming problem and find the maximum profit.

Q14 HINT

Use the information given in the question to set up linear inequalities.

DIVIDE A POLYNOMIAL BY $x \pm p$

KEY POINTS

A quotient in mathematics is the result of the division of two numbers.

EXAMPLE 11

Divide $x^3 + 2x^2 - 17x + 6$ by (x - 3)

SKILLS

ADAPTIVE LEARNING

Start by dividing the first term of the polynomial by
$$x$$
, so that $x^3 \div x = x^2$

$$x - 3)x^3 + 2x^2 - 17x + 6$$

$$x^3 \div x = x^2$$
Next multiply $(x - 3)$ by x^2 , so that $x^2 \times (x - 3) = x^3 - 3x^2$

$$5x^2 - 17x$$
Now subtract, so that $(x^3 + 2x^2) - (x^3 - 3x^2) = 5x^2$
Then copy $-17x$

Step 2:

Repeat the method. Divide
$$5x^2$$
 by x , so that $5x^2 \div x = 5x$

$$\frac{x^2 + 5x}{x^3 - 3x^2} = \frac{x^3 - 3x^2}{5x^2 - 15x} = \frac{5x^2 - 15x}{-2x + 6}$$
Multiply $(x - 3)$ by $5x$, so that $5x \times (x - 3) = 5x^2 - 15x$
Subtract, so that $(5x^2 - 17x) - (5x^2 - 15x) = -2x$

Step 3:

Repeat the method. Divide
$$-2x$$
 by x , so that $-2x \div x = -2$
$$x - 3)x^3 + 2x^2 - 17x + 6$$

$$x^3 - 3x^2$$

$$5x^2 - 17x$$

$$5x^2 - 15x$$

$$-2x + 6$$


$$-2x + 6$$

$$0$$
Multiply $(x - 3)$ by -2 , so that $-2 \times (x - 3) = -2x + 6$
Subtract, so that $(-2x + 6) - (-2x + 6) = 0$
No numbers left to copy, so you have finished. The **remainder** is 0

So
$$x^3 + 2x^2 - 17x + 6 \div (x - 3) = x^2 + 5x - 2$$
 $x^2 + 5x - 2$ is called the **quotient**.

EXAMPLE 12

Divide
$$-3x^4 + 8x^3 - 8x^2 + 13x - 10$$
 by $(x - 2)$.



Step 2:

Step 2:

$$\begin{array}{r}
-3x^3 + 2x^2 \\
x - 2) - 3x^4 + 8x^3 - 8x^2 + 13x - 10 \\
\underline{-3x^4 + 6x^3} \\
2x^3 - 8x^2 \\
\underline{2x^3 - 4x^2} \\
-4x^2 + 13x
\end{array}$$

Repeat the method. Divide $2x^3$ by x, so that $2x^3 \div x = 2x^2$

Multiply
$$(x - 2)$$
 by $2x^2$, so that $2x^2 \times (x - 2) = 2x^3 - 4x^2$

Subtract, so that $(2x^3 - 8x^2) - (2x^3 - 4x^2) = -4x^2$

Copy 13x

Step 3:

Step 4:

5x - 10Multiply (x-2) by 5, so that $5 \times (x-2) = 5x - 10$ Subtract, so that (5x - 10) - (5x - 10) = 0

So
$$(-3x^4 + 8x^3 - 8x^2 + 13x - 10) \div (x - 2) = -3x^3 + 2x^2 - 4x + 5$$

The quotient is $-3x^3 + 2x^2 - 4x + 5$. The remainder is 0.

EXERCISE 5

1 Divide

SKILLS ADAPTIVE LEARNING

a
$$x^3 + 2x^2 - 23x - 60$$
 by $x + 4$

a
$$x^3 + 2x^2 - 23x - 60$$
 by $x + 4$

c
$$x^3 + 6x^2 + 8x + 3$$
 by $x + 1$

e
$$x^3 - x^2 + x + 14$$
 by $x + 2$

g
$$x^3 - x^2 + x - 1$$
 by $x - 1$

b
$$x^3 + 2x^2 - 21x + 18$$
 by $x - 3$

d
$$x^3 + 3x^2 - 18x - 40$$
 by $x - 4$

$$x^3 - 5x^2 + 8x - 4$$
 by $x - 2$

h
$$x^3 + 5x^2 + 2x + 1$$
 by $x + 5$

2 Divide

$$-2x^3 - 5x^2 + 3x + 10$$
 by $x + 2$

c
$$6x^3 + 27x^2 + 14x + 8$$
 by $x + 4$

b
$$4x^3 + 20x^2 - 8x - 96$$
 by $x + 3$

d
$$3x^2 - 10x^2 - 10x + 8$$
 by $x - 4$

e
$$-3x^3 + 15x^2 + 108x - 540$$
 by $x - 6$ **f** $-5x^3 - 27x^2 + 23x + 30$ by $x + 6$

g
$$2x^3 - 3x^2 - 3x + 2$$
 by $x - 2$

3 Divide

a
$$2x^3 + 5x^2 - 5x + 1$$
 by $(2x - 1)$ **b** $6x^3 - 7x^2 - 21x - 12$ by $(3x + 4)$

c
$$3x^3 - 10x^2 - 27x + 10$$
 by $(3x - 1)$ **d** $2x^3 - 3x^2 + 11x + 6$ by $(2x - 1)$

e
$$6x^3 + x^2 - 7x + 2$$
 by $(3x - 1)$ **f** $2x^3 + 7x^2 + 7x + 2$ by $(2x + 1)$

FACTORISE A POLYNOMIAL BY USING THE FACTOR THEOREM

You can factorise a polynomial by using the factor theorem:

If f(x) is a polynomial and f(p) = 0, then x - p is a factor of f(x).

EXAMPLE 13

Show that (x-2) is a factor of $x^3 + x^2 - 4x - 4$ by

SKILLS **ADAPTIVE**

LEARNING

a Algebraic division

b The factor theorem.

b
$$f(x) = x^3 + x^2 - 4x - 4$$
 Write the polynomial as a function $f(2) = 2^3 + 2^2 - 4 \times 2 - 4$ Substitute $x = 2$

$$= 8 + 4 - 8 - 4 = 0$$
 Use the factor theorem:
$$f(x) = x^3 + x^2 - 4x - 4$$
 Substitute $x = 2$

$$= 8 + 4 - 8 - 4 = 0$$
 Use the factor theorem:
$$f(x) = x^3 + x^2 - 4x - 4$$
 Substitute $x = 2$

$$= 8 + 4 - 8 - 4 = 0$$
 Use the factor of $(x) = 0$, then $(x - p)$ is a factor of $(x) = 0$. Here $(x) = 0$, then $(x) = 0$ is a factor of $(x) = 0$.

EXAMPLE 14

Factorise $2x^3 + x^2 - 18x - 9$.

$$f(x) = 2x^{3} + x^{2} - 18x - 9$$
Write the polynomial as a function
$$f(-1) = 2(-1)^{3} + (-1)^{2} - 18(-1) - 9 = 8$$
Try values of x that are a factor of 9,
$$f(1) = 2(1^{3}) + (1)^{2} - 18(1) - 9 = -24$$

$$f(2) = 2(2)^{3} + (2)^{2} - 18(2) - 9 = -25$$

$$f(3) = 2(3)^{2} + (3)^{2} - 18(3) - 9 = 0$$
So $(x - 3)$ is a factor of $2x^{3} + x^{2} - 18x - 9$
Use the factor theorem: If $f(p) = 0$, then $(x - p)$ is a factor of $f(x)$. Here $f(x) = 0$

$$x - 3) 2x^{2} + 7x + 3$$

$$x - 3) 2x^{3} + x^{2} - 18x - 9$$

$$2x^{3} - 6x$$

$$7x^{2} - 18x$$

$$7x^{2} - 21x$$

$$3x - 9$$

$$3x - 9$$

$$0$$

$$2x^{3} + x^{2} - 18x - 9 + x^{2} - 18x - 9$$

$$0$$
You can check your division here: $(x - 3)$ is a factor of $2x^{3} + x^{2} - 18x - 9$, so the remainder must be 0

$$2x^{3} + x^{2} - 18x - 9 = (x - 3)(2x^{2} + 7x + 3)$$

$$= (x - 3)(2x + 1)(x + 3)$$

EXAMPLE 15

Given that (x + 1) is a factor of $4x^4 - 3x^2 + a$, find the value of a.

$$f(x) = 4x^4 - 3x^2 + a$$
Write the polynomial as a function
$$f(-1) = 0$$
Use the factor theorem the other way around: $(x - p)$
is a factor of $f(x)$, so $f(p) = 0$. Here $p = -1$

$$4 - 3 + a = 0$$

$$a = -1$$
Substitute $x = -1$ and solve the equation for a .
Note that $(-1)^4 = (-1)^2 = +1$

EXAMPLE 16

Show that if (x - p) is a factor of f(x) then f(p) = 0

If (x - p) is a factor of f(x) then

 $f(x) = (x - p) \times g(x)$

where g(x) is a polynomial. So

$$f(p) = (p - p) \times g(p)$$

$$= 0 \times g(p)$$

$$= 0$$

So f(p) = 0 as required.

Write the polynomial as a function

Remember that $0 \times \text{anything} = 0$

EXERCISE 6

SKILLS

ANALYSIS

1 b Use the factor theorem to show that

a (x + 2) is a factor of $2x^3 - 3x^2 - 12x + 4$

b (x + 1) is a factor of $2x^4 + 3x^3 - x^2 + 2$

(x + 3) is a factor of $2x^4 + 5x^3 - 8x^2 - 17x - 6$

d (x-2) is a factor of $x^3 - 5x^2 + 2x + 8$

(x + 4) is a factor of $x^7 + 10x^6 + 27x^5 - 57x^3 - 30x^2 + 29x + 20$.

Show that $x^4 - 3x^2 + 2x + 4$ is exactly **divisible** by (x + 1) but not by (x - 2).

3 Now that (x-2) is a factor of and factorise $x^3 - 3x^2 - 2x + 8$.

4 Show that (x-3) is a factor of and factorise $x^3 - 2x^2 - 5x + 6$.

5 Now that (x + 4) is a factor of and factorise $2x^3 + 3x^2 - 18x + 8$.

- 6 Given that (x-2) is a factor of $3x^3 x^2 12x + a$ find the value of a.
- \triangleright Given that (3x + 2) is a factor of $3x^3 + bx^2 3x 2$
 - a find the value of b

b hence factorise completely.

Q8 HINT

Solve simultaneous equations.

- 8 Given that (x-1) and (2x-1) are factors of $px^3 + qx^2 + 9x 2$. find the value of p and q.
- - a find the value of a and b

b factorise f(x) completely.

USING THE REMAINDER THEOREM, FIND THE REMAINDER WHEN A POLYNOMIAL IS DIVIDED BY ax - b

By using the remainder theorem, you can find the remainder when a polynomial is divided by (ax - b). If a polynomial f(x) is divided by (ax - b) then the remainder is $f(\frac{b}{a})$

KEY POINTS

A remainder in mathematics is what is left over in a division problem.

EXAMPLE 17

Find the remainder when $x^3 - 20x + 3$ is divided by (x - 4) using

ADAPTIVE **LEARNING**

- Algebraic division **b** The remainder theorem.
- $\begin{array}{r}
 x^2 + 4x 4 \\
 x 4 \overline{\smash)x^3 + 0x^2 20x + 3} \\
 \underline{x^3 4x^2}
 \end{array}$ Divide $x^3 - 20x + 3$ by (x - 4)Remember to use $0x^2$ $4x^2 - 20x$ $4x^2 - 16x$ -4x + 3-4x + 16-13

The remainder is -13.

b $f(x) = x^3 - 20x + 3$ Write the polynomial as a function

$$f(4) = 4^{3} - 20 \times 4 + 3$$

$$= 64 - 80 + 3$$

$$= -13$$

The remainder is -13.

Use the remainder theorem: If f(x) is divided by (ax - b), then the remainder is $f(\frac{b}{a})$. Compare (x-4) to (ax-b): Here a=1, b=4, and the remainder is $f\left(\frac{4}{1}\right)=f(4)$. So substitute x=4

EXAMPLE 18

When $8x^4 - 4x^3 + ax^2 - 1$ is divided by (2x + 1) the remainder is 3. Find the value of a.

$$f(x) = 8x^4 - 4x^3 + ax^2 - 1$$

$$f\left(-\frac{1}{2}\right) = 3$$

$$8\left(-\frac{1}{2}\right)^4 - 4\left(-\frac{1}{2}\right)^3 + a\left(-\frac{1}{2}\right)^2 - 1 = 3$$

$$8\left(\frac{1}{16}\right) - 4\left(-\frac{1}{8}\right) + a\left(\frac{1}{4}\right) - 1 = 3$$

$$\frac{1}{2} + \frac{1}{2} + \frac{1}{4}\alpha - 1 = 3$$

$$\frac{1}{4}a = 3$$

$$a = 12$$

Use the remainder theorem: If f(x) is divided by (ax - b), then the remainder if $f(\frac{b}{a})$. Compare (2x + 1) to (ax - b): here a = 2, b = -1, and the remainder is 3.

So substitute $x = -\frac{1}{2}$, use the fact that the remainder is 3, and solve the equation for a

EXERCISE 7

REASONING

In each of these questions, find the remainder using the remainder theorem.

a
$$4x^3 - 5x + 4$$
 divided by $(2x - 1)$

b
$$4x^3 - 5x^2 + 7x + 1$$
 divided by $(x - 2)$

c
$$6x^3 + 7x^2 - 15x + 4$$
 divided by $(x - 1)$

c
$$6x^3 + 7x^2 - 15x + 4$$
 divided by $(x - 1)$ **d** $2x^5 - 32x^3 + x - 10$ divided by $(x - 4)$

2 In each of these questions, find the remainder using algebraic division.

a
$$x^3 + 3x^2 + 3x + 1$$
 divided by $(x + 2)$

b
$$x^3 + 2x^2 - x - 1$$
 divided by $(x - 1)$

c
$$3x^3 - x^2 - x - 1$$
 divided by $(x - 4)$

- 3 \triangleright f(x) = $3x^3 5x^2 58x + 40$. Find the remainder when f(x) is divided by (x 4).
- When $5 + 6x + bx^2 x^3$ is divided by (x 1) the remainder is 17. Find b.
- 5 The expression $3x^3 + px^2 3x + 4$ is divided by (x + 4). State the remainder of this expression in terms of p.
- 6 When $16x^3 bx^2 + 30x 8$ is divided by (2x 1) the remainder is 1. Find b.
- 7 When $ax^3 bx + 5$ is divided by (x 2) the remainder is 5 and when it is divided by (x + 1)the remainder is 6. Find the values of a and b.
- 8 When $ax^3 x^2 + bx + 2$ is divided by (x + 3) the remainder is -73 and when it is divided by (x-2) the remainder is 22.
 - **a** Find the values of a and b.
 - **b** Find the remainder when the expression is divided by (3x + 1).
- **9** The expression $2x^3 x^2 + ax + b$ gives a remainder of 14 when divided by (x 2) and a remainder of -86 when divided by (x + 3). Find the values of a and b.
- 10 The expression $3x^3 + 2x^2 px + q$ is divisible by (x 1) but leaves a remainder of 10 when divided by (x + 1). Find the values of p and q.

EXAM PRACTICE: CHAPTER 3

Solve these simultaneous equations:

$$x + y = 2$$

$$4y^2 - x^2 = 11$$

Solve these simultaneous equations:

$$y-3x+2=0$$

$$y^2 - x = 6x^2$$

The line y = x + 2 meets the curve $x^2 + 4y^2 - 2x = 35$ at the points p and q.

Find the coordinates of p and the coordinates of q.

[3]

Find the set of values of x for which:

[3]

a
$$4x - 5 > 15 - x$$

b
$$x(x-4) > 12$$

Find the set of values of x for which:

$$(2x-7)(x+1)<0$$

b
$$3(x-2)-8+2x<0$$

[2]

c Both
$$(2x-7)(x+1) < 0$$
 and $3(x-2)-8+2x < 0$

[3]

Find the set of values of x for which:

$$2x^2 - 5x - 12 > 0$$

[3]

Solve these simultaneous equations:

[4]

$$x-6y-1=0$$

$$x=\frac{1}{y}+4y$$

Find the set of values of *x* for which:

a
$$2p^2 + 9p > 5$$

[2]

b
$$n(n + 9) < 2(n - 5)$$

[3]

$$f(x) = 2x^3 - 7x^2 - 10x + 24$$

[4]

a Show that
$$(x + 2)$$
 is a factor of $f(x)$.

b Hence factorise
$$f(x)$$
.

 $f(x) = x^3 + 8x^2 + 17x + 16$

[4]

a Show that
$$f(x) \div (x + 5)$$
 has a remainder of 6.

b Find the quotient of $f(x) \div (x + 5)$.

$$f(x) = -3x^3 + 13x^2 - 6x + 8$$

[4]

a Show that
$$(x - 4)$$
 is a factor of $f(x)$.

b Hence factorise f(x) completely.

Factorise completely $2x^3 - 7x^2 - 5x + 4$

[3]

13
$$f(x) = x^4 + 5x^3 + px + q$$

p and q are integers.

[4]

[5]

- The remainder when f(x) is divided by (x-2) is equal to the remainder when f(x) is divided by (x+1). Find the value of p.
- **b** Given that (x + 3) is a factor of f(x), find the value of q.

$$f(x) = x^3 - 2x + 21$$
 [4]

- a Show that (x + 3) is a factor of f(x).
- **b** Solve f(x) = 0.

15
$$f(x) = px^3 + 6x^2 + 12x + q$$
. [4]

Given that the remainder when f(x) is divided by (x - 1) is equal to the remainder when f(x) is divided by (2x + 1),

- a find the value of p.
- 16 $f(x) = x^3 x^2 7x + a$, where a is a constant.

Given that f(4) = 0,

- a find the value of a. [1]
- b factorise f(x) as the product of a linear factor and a quadratic factor. [2]
- a Factorise completely $g(x) = 2x^3 x^2 4x + 3$
 - **b** Solve g(x) = 0. [1]
- Illustrate, on a graph, the region that is represented by $x \le \frac{y}{2} + 3$, y > 3x 4 and 2x + y > -4.
- A company buys two types of diary to send to its customers, a desk top diary and a pocket diary. They will need to place a minimum order of 200 desk top and 80 pocket diaries.

They need at least twice as many pocket diaries as desk top diaries.

They will need a total of at least 400 diaries.

Each desk top diary costs \$6 and each pocket diary costs \$3.

The company wishes to minimize the cost of buying these diaries.


- a Write down the inequalities of this problem.
- b Represent these graphically. [4]

CHAPTER SUMMARY: CHAPTER 3

- When you multiply or divide an inequality by a negative number you need to reverse the inequality sign.
- The steps for solving a quadratic inequality are:
 - Solve the corresponding quadratic equation.
 - Sketch the graph of the quadratic function.
 - Use the sketch to find the required set of values.
- If f(x) is a polynomial and f(a) = 0, then (x a) is a factor of f(x). This is known as the *factor* theorem.
- If f(x) is a polynomial and $f\left(\frac{b}{a}\right) = 0$ then (ax b) is a factor of f(x). This is also known as the factor theorem.
- If a polynomial f(x) is divided by (ax b) then the remainder is $f(\frac{b}{a})$. This is known as the *remainder* theorem.

CHAPTER 3

CHAPTER 4



4 SKETCHING POLYNOMIALS

Graphs are a vital tool for physicists. They help explain relationships between physical things, for example the volume of gas is inversely proportional to the pressure acting on it. If this relationship is modelled as a graph it produces the graph $y=\frac{k}{r}$. You will learn about this graph, and others, in this chapter.

An inflated balloon also obeys the equation above. If it is placed in a sealed chamber and the air pressure is doubled, it shrinks to half its original volume. If the pressure is decreased to one-tenth of the original, the balloon will expand to ten times its original volume - if it does not burst first.



LEARNING OBJECTIVES

- Sketch cubic curves of the form $y = ax^3 + bx^2 + cx + d$ or y = (x + a)(x + b)(x + c)
- Sketch and interpret graphs of cubic functions of the form $y = x^3$
- Sketch the **reciprocal function** $y = \frac{k}{r}$ where k is a constant
- Sketch curves of functions to show points of intersection and solutions to equations
- Apply transformations to more complicated curves

STARTER ACTIVITIES

Sketch these curves.

1 **a**
$$y = x^2$$

$$y = x^2 + 1$$

$$y = 2x^2$$

d
$$u = 2(x^2 + 1)$$

$$u = (x + 1)^2$$

$$f \quad u = \Delta x^2$$

$$u = v^2 - A$$

h
$$y = \frac{1}{2}x^2$$

a
$$y = x^2$$
 b $y = x^2 + 1$ **c** $y = 2x^2$ **d** $y = 2(x^2 + 1)$ **e** $y = (x + 1)^2$ **f** $y = 4x^2$ **g** $y = x^2 - 4$ **h** $y = \frac{1}{2}x^2$ **i** $y = \frac{1}{2}x^2 - 2$ **j** $y = \sqrt{x}$ **k** $y = \frac{1}{x}$ **l** $y = x^3$ **m** $y = x^3 + 1$

$$u = \sqrt{x}$$

$$\mathbf{k} \quad y = \frac{1}{x}$$

$$y = x^3$$

$$y = x^3 + 1$$


SKETCH CUBIC CURVES OF THE FORM $y = ax^3 + bx^2 + cx + d$ OR y = (x + a)(x + b)(x + c)


You need to be able to sketch equations of the form

$$y = ax^3 + bx^2 + cx + d$$
 or $y = (x + a)(x + b)(x + c)$

This involves finding the points where the curve crosses the x- and y-axes.

These types of curves are known as *cubic* curves and the general shape is as follows:





 $y = -x^3$ is a reflection in the x-axis, as all the y values are now negative

EXAMPLE 1

Sketch the curve with the equation y = (x - 2)(x - 1)(x + 1).

Put y = 0 and solve for x to find the roots of the equation. (Where the curve crosses the x-axis)

If
$$y = 0$$

$$0 = (x - 2)(x - 1)(x + 1)$$

So
$$x = 2, 1 \text{ or } -1$$

$$(-1, 0)$$
 (1, 0) and (2, 0)

Put x = 0 to find out where the curve crosses the *y*-axis


If
$$x = 0$$
, $y = (0 - 2)(0 - 1)(0 + 1)$

So
$$y = 2$$


(0, 2)

Now plot these coordinates on a Cartesian graph.

y = (x-2)(x-1)(x+1) crosses the y-axis at (0, -2) and the x-axis at (-1, 0) (1, 0) (2, 0).



Next draw a curve through all these points.



Note: You will learn more about maximum and minimum points in Chapter 9.

EXAMPLE 2

Sketch the curve of y = (x - 2)(1 - x)(1 + x) and show the points where it crosses the coordinate axes.

Put y = 0 and solve for x to find the roots of the equation (where the curve crosses the x-axis).

If
$$y = 0$$
, $0 = (x - 2)(1 - x)(1 + x)$

So
$$x = 2, 1 \text{ or } -1$$

$$(-1, 0)$$
 (1, 0) and (2, 0)

Put x = 0 to find out where the curve crosses the *y*-axis.


If
$$x = 0$$
, $y = (0 - 2)(1 - 0)(1 + 0)$

So
$$y = -2$$


$$(0, -2)$$

Now plot these coordinates on a Cartesian graph.

The curve crosses the y-axis at (0, -2) and the x-axis at (-1, 0) (1, 0) (2, 0).



Now draw the curve through all these points.



EXAMPLE 3

Sketch these curves.

$$y = (x-1)^2(x+1)$$

b
$$y = x^3 - 2x^2 - 3x$$

a
$$y = (x - 1)^2(x + 1)$$

If $y = 0$

$$0 = (x - 1)^2(x + 1)$$

So
$$x = 1$$
 or $x = -1$

Points are (-1, 0) and (1, 0).


If
$$x = 0$$

$$y = (0 - 1)^2(0 + 1)$$

So
$$y = 1$$

Point is (0, 1).

Plot the points and sketch the curve.



Repeated root from $(x - 1)^2$

b
$$y = x^3 - 2x^2 - 3x$$

$$y = x(x^2 - 2x - 3)$$

 $y = x(x - 3)(x + 1)$

Factorise

If
$$u = 0$$

$$0 = x(x-3)(x+1)$$

So
$$x = 0.3$$
 or -1

Points are (0, 0), (3, 0) and (-1, 0).


If
$$x = 0$$

$$y = 0(0 - 3)^2(0 + 1)$$

So
$$y = 0$$

Point is (0, 0).

Plot the points and sketch the curve



SKILLS

ADAPTIVE LEARNING Sketch these curves and indicate clearly the points of intersection with the axes.

a
$$y = (x + 3)(x - 2)(x + 1)$$

a
$$y = (x+3)(x-2)(x+1)$$
 b $y = (x+1)(x-2)(x-3)$ **c** $y = (x+4)(x-3)(x+2)$

$$y = (x + 4)(x - 3)(x + 2)$$

d
$$y = x(x-1)(x+1)$$

d
$$y = x(x-1)(x+1)$$
 e $y = 3x(x-1)(2x+1)$ **f** $y = x(2x+1)(x-3)$

$$y = x(2x + 1)(x - 3)$$

g
$$y = 4x(2x - 1)(x + 2)$$
 h $y = (x + 1)^2(3 + x)$ i $y = (x - 2)^2(x - 4)$

$$y = (x + 1)^2(3 + x)$$

$$u = (x-2)^2(x-4)^2$$

$$y = (x + 2)^2(5 + x)$$

$$y = x^2(3 + x)$$

$$y = (x-4)^2(x+5)$$

Sketch these functions and indicate the points of intersection with the axes.

$$y = x^3 + x^2 - 2x$$

b
$$y = x^3 + 5x^2 + 4x$$
 c $y = x - x^3$

$$u = x - x$$

d
$$y = 3x + 2x^2 - x^3$$


$$y = 12x^3 - 3x$$

$$y = x^3 - 9x$$

$$y = x^3 - 9x^2$$

SKETCH AND INTERPRET GRAPHS OF CUBIC FUNCTIONS OF THE FORM $u=x^3$

Below is a sketch of the graph $y = x^3$



EXAMPLE 4

Sketch the curves of these equations and show their relative positions to $y = x^3$.


a
$$y = -x^3$$

b
$$y = (x + 1)^3$$


$$y = (3 - x)^3$$

HINT

You do not need to plot any points. It is quicker to realise that the curve $y = -x^3$ is a reflection of the curve $y = x^3$



b





HINT

You do not need to plot any points. Instead realise that this is a translation of the curve $y = x^3$. When a value 'a' is added to a cubic, inside brackets, it creates a horizontal shift of '-a'

When
$$x = 0$$
: $y = (0 + 1)^3$, so that $y = 1$

C





When x = 0: $y = (3 - 0)^3$, so that y = 27

EXERCISE 2

Sketch these curves and show their relative positions in relation to $y = x^3$

SKILLS

ADAPTIVE LEARNING

- **a** $y = (x-2)^3$ **b** $y = (2-x)^3$ **c** $y = (x-3)^3$ **d** $y = (x+4)^3$

- **e** $y = -(x-2)^3$ **f** $y = -(x-4)^3$ **g** $y = (x+\frac{1}{2})^3$ **h** $y = (1-x)^3$

SKETCH THE RECIPROCAL FUNCTION $y = \frac{k}{x}$ where k is a constant


The curves with the equations $y = \frac{k}{r}$ fall into two categories.

Type 1: $y = \frac{k}{x}$ where k > 0

The curve does not cross the axes.

The curve tends towards the x-axis when x is large and positive or large and negative.

The curve tends towards the y-axis when x is large and positive or large and negative.



When x = 0, y is undefined. When y = 0, x is undefined.

$$x \to +\infty$$
, $y \to 0$


$$-x \to +\infty$$
, $y \to 0$

$$y \to +\infty, x \to 0$$

$$-y \to +\infty, x \to 0$$

These are the horizontal and vertical asymptotes (where the graph never reaches the x- and y-axes).


Type 2: $y = \frac{k}{x}$ where k < 0



The curves behave in the same way as in Type 1, but are confined to two different quadrants


EXAMPLE 5

- a Sketch the graph $y = \frac{3}{x}$
- **b** Sketch the graph $y = -\frac{1}{x}$
- Sketch the graph of the function $y = \frac{3}{x}$.



The curve will be the same as $y = \frac{1}{x}$, but further out from the origin. Its asymptotes are, however, the same.

b Sketch the graph of the function $y = -\frac{1}{x}$.



The curve will be the same as $y = \frac{1}{x}$, but reflected in the *x*-axis.

Sketch these pairs of equations on the same axes.

SKILLS INTERPRETATION

a
$$y = \frac{1}{x}, y = \frac{3}{x}$$

b
$$y = \frac{4}{x}, y = \frac{8}{x}$$

b
$$y = \frac{4}{x}, y = \frac{8}{x}$$
 c $y = \frac{3}{x}, y = -\frac{3}{x}$

d
$$y = -\frac{4}{x}, y = -\frac{6}{x}$$
 e $y = \frac{8}{x}, y = -\frac{7}{x}$ **f** $xy = 6, xy = -5$


$$y = \frac{8}{x}, y = -\frac{7}{x}$$

f
$$xy = 6, xy = -5$$

SKETCH CURVES OF DIFFERENT FUNCTIONS TO SHOW POINTS OF INTERSECTION AND SOLUTIONS TO EQUATIONS

EXAMPLE 6

- a On the same diagram sketch the curves with the equations y = x(x 3) and $y = x^2(1 x)$.
- **b** Find the coordinates of the points of intersection.



b
$$y = x(x - 3)$$
 and $y = x^2(1 - x)$

The coordinates of the points of intersection are the points where the two functions have the same value.

$$x(x-3)=x^2(1-x)$$

 $x^2-3x=x^2-x^3$ Expand brackets
 $x^3-3x=0$ Collect like terms and equate to 0
 $x(x^2-3)=0$ Factorise
 $x=0 \text{ or } x=\pm\sqrt{3}$

You can use the equation y = x(x - 3) to find the *y*-coordinates:

If
$$x = -\sqrt{3}$$
, $y = -\sqrt{3}(-\sqrt{3} - 3)$ $y = 3 + 3\sqrt{3}$

If $x = 0$, $y = 0(0 - 3)$ $y = 0$


If $x = \sqrt{3}$, $y = \sqrt{3}(\sqrt{3} - 3)$ $y = 3 - 3\sqrt{3}$

The coordinates of the points of intersection are $(-\sqrt{3}, 3 + 3\sqrt{3})$, (0, 0) and $(\sqrt{3}, 3 - \sqrt{3})$.

EXAMPLE 7

a On the same diagram sketch the curves with equations $y = x^2(x-1)$ and $y = \frac{2}{x}$.

b Explain why your sketch shows there are two roots to the equation $x^2(x-1) = \frac{2}{x^2}$



HINT

 $y=x^2(x-1)$

The cubic 'positive' shape crosses the x-axis at x = 0 and x = 1.

$$y = \frac{2}{x}$$

The reciprocal 'positive' shape does not cross any axes.

b From the sketch there are only two points of intersection of the curves. This means there are only two values of x where

$$x^2(x-1)=\frac{2}{x}$$

So this equation has two roots.

Note: You would not be expected to solve this equation.

EXERCISE 4

SKILLS PROBLEM SOLVING 1 Consider the pairs of equations, a-f, below. In each case

i sketch the two curves on the same axes

ii state the number of points of intersection, if any

write down a suitable equation which would give the *x*-coordinates of these points. You do not need to solve the equation.

a
$$y = x^2, y = x^3$$

b
$$y = x(x + 2), y = -\frac{3}{x}$$

$$y = x^2(1-x), y = -\frac{2}{x}$$

d
$$y = x(x-4), y = (x-2)^3$$

$$y = -x^3, y = -\frac{2}{x}$$

$$y = -x^3, y = x^2$$

2 a On the same axes sketch the curves given by $y = x^2(x-4)$ and y = x(4-x).

b Find the coordinates of the point of intersection.

3 **a** On the same axes sketch the curves given by $y = (x - 1)^3$ and $y = x^2 - 1$.

b Find the coordinates of the point of intersection.

4 Non the same axes sketch $y = \frac{2}{x}$ and $y = -x(x-1)^2$ and use your sketch to explain why there is no solution to $\frac{2}{x} = -x(x-1)^2$.

Sketch the curves of $y = 1 - 4x^2$ and $y = x(x - 2)^2$ and use your sketch to explain how many solutions there are to the equation $1 - 4x^2 = x(x - 2)^2$.

Sketch the curves $y = x^3 - 2x^2 - x + 2$ and the line y = 14x + 2 and find the points of intersection.

69

APPLY TRANSFORMATIONS TO CURVES

Applying transformations to curves is beyond the scope of the specification and won't be examined. However this exercise can be set as an extension activity. Polynomial curves can be transformed in four basic ways. This is explained in more detail below.

| NAME | f(x) | DESCRIPTION |
|---|----------|--|
| Horizontal translation of $-a$ | f(x + a) | The value of a is subtracted from all the x -coordinates, but the y -coordinates stay unchanged. In other words the curve moves a units to the left. |
| Vertical translation of $+a$ | f(x) + a | The value of a is added to all the y -coordinates, but the x -coordinates stay unchanged. In other words the curve moves a units up. |
| Horizontal stretch of scale factor $\frac{1}{a}$ | f(ax) | All the x coordinates are multiplied by $\frac{1}{a}$, but the y -coordinates stay unchanged. In other words the curve is squashed in a horizontal direction. |
| Vertical stretch of scale factor α | af(x) | All the y -coordinates are multiplied by a , but the x -coordinates stay unchanged. In other words the curve is stretched in a vertical direction. |

EXAMPLE 8


Sketch these functions.

a
$$f(x) = x^2$$


b
$$g(x) = (x + 3)^2$$

c
$$h(x) = x^2 + 3$$


 $f(x) = x^2$ is a standard curve:



 $g(x) = (x + 3)^2$ is the same moved 3 units to the left:




 $h(x) = x^2 + 3$ is the same curve, moved 3 units up:




EXAMPLE 9


Sketch the graph of $y = (x - 2)^2 + 3$



Step 1: Sketch the graph of $f(x) = x^2$




Step 2: Sketch the graph of $f(x) = (x - 2)^2$ Horizontal translation of +2





Step 3: Sketch the graph of $f(x) = (x - 2)^2 + 3$ Vertical translation of +3


EXAMPLE 10

Sketch the graph of $y = \frac{2}{x+5} - 3$









Step 1: Start with the graph $y = \frac{1}{r}$

$$f(x)=\frac{1}{x}$$

Asymptotes are: x = 0 and y = 0

Step 2:
$$y = \frac{1}{x+5}$$

Horizontal shift of -5

Or:

$$f(x + 5)$$

and

Asymptotes are: x = -5 and y = 0

Step 3:
$$y = \frac{2}{x+5}$$

Vertical stretch, scale factor 2

Or:

$$2f(x + 5)$$

and

Asymptotes are: x = -5 and y = 0

Step 4:
$$y = \frac{2}{x+5} + 3$$

Horizontal translation of -3

Or:

$$2f(x + 5) + 3$$

Asymptotes are: x = -5 and y = 3

EXERCISE 5

Sketch these curves.



$$y = 3x^2 - 4$$

b
$$y = \frac{3}{x} - 2$$

$$y = 3(3x + 1)^2$$

a
$$y = 3x^2 - 4$$
 b $y = \frac{3}{x} - 2$ **c** $y = 3(3x + 1)^2$ **d** $y = \frac{2}{x - 2}$

$$y = \frac{1}{x - 1} + 3$$

e
$$y = \frac{1}{x-1} + 3$$
 f $y = \frac{1}{x-2} - 1$ **g** $y = \frac{3}{2-x} - 5$ **h** $y = 3x^2 + 2$

$$y = \frac{3}{2-x} - \xi$$


h
$$y = 3x^2 + 2$$

[4]

EXAM PRACTICE: CHAPTER 4

- The curve C has equation $y = \frac{3}{x}$ and the line l has equation y = 2x + 5
 - a Sketch the graphs of *C* and *l*, indicating clearly the coordinates of any intersections with the axes.
 - **b** Find the coordinates of the points of intersection of C and I.
- The curve *C* has the equation $y = (x + \frac{1}{2})(x 2)(x + 1)$. Sketch the curve, indicating where the curve crosses the *x*-axis. [3]
- Sketch the graphs of y = x(x + 2)(3 x) and $y = -\frac{2}{x}$, showing clearly the coordinates of all the points where the curves cross the coordinate axes. [4]
 - b Using your sketch, state the number of real solutions to the equation $x(x+2)(3-x)=-\frac{2}{x}$ and give a reason for your answer. [1]
- Sketch the graphs of y = x(4 x) and $y = x^2(6 x)$ showing clearly the coordinates of the point of intersection. [3]
- a Factorise completely $x^3 3x$. [1]
 - **b** Sketch the curve C with equation $y = x^3 3x$. [2]
- The diagram below shows a sketch of $y = \frac{1}{x}$.


 Copy the diagram. On the same diagram sketch the graph $y = \frac{2}{x+1} + 3$ [2]



The diagram below shows a sketch of the curve with equation

$$y = f(x)$$
 where $f(x) = \frac{x}{x-2}$.

Copy the diagram. On the same diagram sketch the curve with equation y = f(x - 1) and state the equations of the asymptotes of this curve. [3]



- a $x^3 4x^2 + x + 6 = (x + 1)(ax^2 bx + 6)$. Find a and b by factorising.
 - **b** Hence sketch the curve of $y = x^3 4x^2 + x + 6$. [2]


[3]

[2]

- Sketch these curves on the same axes.
 - **a** $y = x^3$
 - **b** $y = -\left(x \frac{1}{3}\right)^3$
 - $y = -2\left(x \frac{1}{3}\right)^3$ [4]
- The diagram below shows a sketch of a curve y = f(x).
 - a On the same axes sketch the curves with the equations

$$y = f(x + 3)$$

$$y = -f(-x)$$
 [2]




- a On the same axes sketch the curves given by $y = \frac{1}{x}$ and $y = -x(x-2)^2$. [4]
 - **b** Explain how your sketch shows that there are no solutions to the equation $1 + x^2(x - 1)^2 = 0.$ [1]

73


CHAPTER SUMMARY: CHAPTER 4

You need to know the shapes of these basic curves.


$$y = x^2$$




$$y = (x - a)(x - b)(x - c)$$



$$y = x^3$$




$$y = \frac{1}{x}$$



You also need to know the basic rules of transformations.

| NAME | f(x) | DESCRIPTION |
|--------------------------------|----------|---|
| Horizontal translation of $-a$ | f(x + a) | The value of a is subtracted from all the x -coordinates, |
| | | but the y -coordinates stay unchanged. In other words |
| | | the curve moves a units to the left. |
| Vertical translation of $+a$ | f(x) + a | The value of a is added to all the y -coordinates, but |
| | | the <i>x</i> -coordinates stay unchanged. In other words the |
| | | curve moves a units up. |
| Horizontal stretch of scale | f(ax) | All the x-coordinates are multiplied by $\frac{1}{a}$, but the |
| factor $\frac{1}{a}$ | | y-coordinates stay unchanged. In other words the |
| a | | curve is squashed in a horizontal direction. |
| Vertical stretch of scale | af(x) | All the y -coordinates are multiplied by a , but the |
| factor a | | x-coordinates stay unchanged. In other words the |
| | | curve is stretched in a vertical direction. |

CHAPTER 5



5 SEQUENCES AND SERIES

A recurrence relationship is a sequence where the next **term** of the sequence is derived from the preceding terms. The best-known is the Fibonacci sequence, named after the Italian mathematician who wanted to find out how fast rabbits could breed.

Each term is found by adding up the two previous terms. The sequence is: 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

The sequence is found throughout nature, from leaves to pineapple scales. Natural growth often follows this pattern, for example in animal cells, grains and even **beehives**.



Fibonacci introduced the sequence to Europe in 1202, but it was known to the Arabs and the Indians before that.

LEARNING OBJECTIVES

- Identify an arithmetic sequence
- Find the common difference, first term and nth term of an arithmetic series
- Find the sum of an arithmetic series and be able to use Σ (sigma) notation
- Identify a geometric sequence
- Find the **common ratio**, first term and *n*th term of the sequence
- Find the sum of a geometric series
- Find the sum to infinity of a convergent geometric series

STARTER ACTIVITIES

The two key concepts in this chapter are sequences and series. In the English language these terms are interchangeable. In mathematics, however, the definition of a sequence and a series differ greatly, therefore it is important to know the difference.

A sequence is a list of numbers: a_1 , a_2 , a_3 , ...

A series is the expression you get when you insert 'addition' symbols: $a_1 + a_2 + a_3 + ...$

In a sequence the order in which the numbers are listed is important, for instance:

1, 2, 3, 4, 5, ... is one sequence, and

2, 1, 4, 3, 6, 5, ... is an entirely different sequence.

A series is a sum of numbers. For example,

$$1 + 3 + 5 + 7 + 9 + \dots$$

or

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots$$

IDENTIFY AN ARITHMETIC SEQUENCE

Consider these three common sequences

2, 5, 8, 11, ...

0, 5, 10, 15, ...

 $5, 2, -1, -4, \dots$

To get to the next term, you add a fixed value to the previous term. Hence the difference between the terms is fixed. Note that you can also subtract a common difference to obtain a sequence.

A sequence that increases or decreases by a constant amount each time is a called an *arithmetic* sequence.

Here are some examples of arithmetic sequences

3, 7, 11, 15, 19, ... because you add 4 each time

2, 7, 12, 17, 22, ... because you add 5 each time

17, 14, 11, 8, ... because you subtract 3 each time

 $a, a + d, a + 2d, a + 3d, \dots$ because you add d each time

KEY POINTS

- A recurrence relationship is a sequence where there is a relationship between two consecutive terms.
- A recurrence relationship of the form $U_{k+1}=U_k+c,\,k\geq 1,\,c\in\mathbb{R}$ is called an arithmetic sequence

EXAMPLE 1

Find the a 10th, b nth and c 50th terms of the arithmetic sequence 3, 7, 11, 15, 19, ...

SKILLS

ADAPTIVE LEARNING PROBLEM SOLVING a First term = 3 The sequence goes up in fours, starting at 3. The first term is $3 + 0 \times 4$

Second term = 3 + 4

The second term is $3 + 1 \times 4$

Third term = 3 + 4 + 4

The third term is $3 + 2 \times 4$

Fourth term = 3 + 4 + 4 + 4

The fourth term is $3 + 3 \times 4$

Therefore the 10th term is $3 + 9 \times 4 = 3 + 36 = 39$

b The *n*th term is:

$$3 + (n-1) \times 4 = 4n-1$$

nth term = first term + $(n - 1) \times$ difference

c The 50th term is $3 + (50 - 1) \times 4 = 3 + 196 = 199$

EXAMPLE 2

A tree that is 6 metres tall is planted in a garden. If it grows 1.5 metres a year

- a how tall will it be after it has been in the garden for 8 years?
- **b** after how many years will it be 24 metres tall?
- **a** $6 + 8 \times 1.5$

It starts at 6 metres and it has 8 years' growth at 1.5 metres a year

- = 6 + 12
- = 18 metres

b 24 - 6 = 18 metres

Find out how much it has grown in total



So number of years = $\frac{18}{1.5}$ = 12 years.

It grows 1.5 metres a year

EXAMPLE 3

Find the number of terms in the arithmetic sequence 7, 11, 15, ..., 143

The sequence goes up in fours.

It goes from 7 to 143, a difference of 136.

$$\frac{136}{4} = 34 \text{ jumps}$$

Work out how to get from one term to the next

Work out the difference between the last number and the first number

There is one more term than the number of jumps, so 34 jumps means 35 terms.

EXERCISE

SKILLS ADAPTIVE

LEARNING **PROBLEM**

SOLVING

Which of these sequences are arithmetic?

a 9, 11, 12, 14

b 8, 10, 12, 14, 16 **c** $\frac{1}{3}$, $\frac{1}{2}$, $\frac{2}{3}$

d 16, 8, 4, 2, 1

e 7, 10, 13, 16, 19 **f** 2.5, 4, 5.5, 7

g -3, -6, -8, -10, -12 **h** 2y, 4y, 6y, 8y **i** 4 + d, 6 + 2d, 8 + 3d, 10 + 3d

2 Find the 10th term in this arithmetic progression.

a 3, 5, 7, 9

b 7, 7.5, 8, 8.5

69, 73, 77, 81

 \mathbf{d} x, 2x, 3x, 4x

e 4 + a, 6 + 2a, 8 + 3a

3 Nadine puts \$4000 into her account. Every month after that, she pays in another \$200. How much money in total will she have invested at the start of



b the *n*th month?

Note: At the start of the 6th month she will have only made 5 payments.



Calculate the number of terms in these arithmetic sequences

a 5, 9, 13, ... 133

b 11, 14, 17, ... 59 **c** 136, 125, 114, ... -84

d 671, 663.5, 656, ... 581 **e** y, 3y, 5y, ... 35y **f** m, m + p, m + 2p, ... m(n - 1)d, ...

FIND THE COMMON DIFFERENCE, FIRST TERM AND nth TERM OF AN **ARITHMETIC SERIES**

Arithmetic series are formed by adding together terms of an arithmetic sequence, $u_1 + u_2 + u_3 \dots + u_n$ In an arithmetic series, the next term is found by adding (or subtracting) a constant number.

This number is called the common difference and is often represented by d.

The first term is often represented by a.

All arithmetic series can be written in the form:

$$a$$
 + $(a+d)$ + $(a+2d)$ + $(a+3d)$ + $(a+4d)$ + ...
First term Second term Third term Fourth term Fifth term

Look at the relationship between the number of the term and coefficient of d. You should be able to see that the coefficient of d is one less than the number of the term.

You can use this fact to produce a formula for the nth term of an arithmetic series:

KEY POINTS

The *n*th term of an arithmetic series $u_n = a + (n-1)d$, where a is the first term and d is the common difference. Please note that this formula is not given in the exam.

EXAMPLE 4

Find i the 20th and ii the 50th terms of these series.

SKILLS

PROBLEM SOLVING

a
$$a=4$$
 and $d=3$

$$d = 7 - 4$$

20th term

Use the formula $u_n = a + (n - 1)d$ with n = 20

$$u_{20} = 4 + (20 - 1) \times 3$$

= 4 + 19 × 3
= 61

ii 50th term

Use the formula
$$u_n = a + (n-1)d$$
 with $n = 50$

$$u_{50} = 4 + (50 - 1) \times 3$$

= 4 + 49 × 3
= 151

b
$$a = 100$$
 and $d = -7$

$$d = 93 - 100 = -7$$

20th term

$$u_{20} = 100 + (20 - 1) \times (-7)$$

= 100 + 19 × (-7)
= -33

ii 50th term

$$u_{50} = 100 + (50 - 1) \times (-7)$$

= 100 + 49 × (-7)
= -243

EXAMPLE 5

For the arithmetic series 5 + 9 + 13 + 17 + 21 + ... + 805:

- a Find the number of terms in the series.
- **b** Which term of the series would be 129?

The *n*th term $u_n = a + (n - 1)d$, so replace

 u_n with 805 and solve for n

a The series is 5 + 9 + 13 + 17 + 21 + ... + 805

With
$$a = 5$$
 and $d = 4$

$$805 = 5 + (n - 1) \times 4$$

$$805 = 5 + 4n - 4$$

$$805 = 4n + 1$$

$$804 = 4n$$

$$n = 201$$

There are 201 terms in this series.

b
$$129 = 5 + (n-1)d$$

$$129 = 4n + 1$$

$$128 = 4n$$

$$n = 32$$

The 32nd term is 129.

The nth term $u_n = a + (n - 1)d$, so replace u_n with 129 and solve for n

EXAMPLE 6

Given that the 3rd term of an arithmetic series is 20 and the 7th term is 12

a find the first term

b find the 20th term.

Note: These are very popular questions and involve setting up and solving simultaneous equations.

a 3rd term is 20 so a + 2d = 20 (1)

7th term is 12 so a + 6d = 12 (2)

Subtracting (1) from (2):

$$4d = -8$$

$$d = -2$$

The common difference is -2.

$$a + 2 \times (-2) = 20$$

$$a - 4 = 20$$

$$a = 24$$

The first term is 24.

b The 20th term is:

$$u_{20} = a + 19d$$

$$= 24 + 19 \times (-2)$$

$$= 24 - 38$$

$$= -14$$

The 20th term is -14.

Use that the nth term is $u_n = a + (n - 1)d$, with n = 3 and n = 7

Substitute d = -2 back into equation (1)

EXERCISE :

Find i the 20th term and ii the nth term of these arithmetic series.

e
$$\frac{6}{5} + \frac{23}{15} + \frac{28}{15} + \frac{11}{15} + \dots$$

$$f q + 3q + 5q + 7q + ...$$

$$5k + k + (-3k) + (-7k) + \dots$$

2 Find the number of terms in these arithmetic series.

b
$$-14 + (-8) + (-2) + ... + 700$$

d
$$x + 5x + 9x + ... + 201x$$

- The 1st term of an arithmetic series is 6 and the 4th term is 18.
 Find the common difference.
- 4 ▶ The 3rd term of an arithmetic series is 12 and the 10th term is -93.
 - a Find the first term and the common difference.
 - **b** Find the term in which the series turns negative.
- 5 The 20th term of an arithmetic series is 18.82 and the 35th term is 15.52. Find the 5th term
- The first three terms of an arithmetic series are 5p, 20 and 3p. Find the value of p and hence the values of the three terms.
- 7 For which values of n would the expressions -8, n^2 and 17n form the first three terms of an arithmetic series?

FIND THE SUM OF AN ARITHMETIC SERIES AND BE ABLE TO USE Σ NOTATION

The method of finding the sum of an arithmetic series is **attributed** to the famous mathematician Gauss. He is said to have solved the problem at junior school.

The formula he proved for the sum of the arithmetic series is

KEY POINTS

The formula for the sum of an arithmetic series is

$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$
 or $S_n = \frac{n}{2}(\alpha + L)$

where a is the first term, d is the common difference, n is the number of terms and L is the last term in the series.



EXAMPLE 7

Find the sum of the first 100 odd numbers.

$$S_n = 1 + 3 + 5 + 7 + ...$$

= $\frac{100}{2}[2 \times 1 + (100 - 1)2]$
= $50(2 + 198)$
= 50×200
= 10000

Use
$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$

Alternatively, find L and use $S_n = \frac{n}{2}(\alpha + L)$

Note: This is a very useful formula that is worth remembering.

$$L = \alpha + (n - 1)d$$

= 1 + 99 × 2
= 199

$$S_n = \frac{n}{2}(a + L)$$

$$= \frac{100}{2}(1 + 199)$$

$$= 10000$$

EXAMPLE 8

Find the minimum number of terms required for the sum 4 + 9 + 14 + 19 + ... to exceed 2000.

$$4 + 9 + 14 + 19 + ... > 2000$$

Always establish what you are given in a question. As you are adding on positive terms, it is easier to solve the equality $S_n = 2000$

Using
$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$

You know
$$a = 4$$
, $d = 5$ and $S_n = 2000$, you need to find n

$$2000 = \frac{n}{2}[2 \times 4 + (n-1) \times 5]$$

$$4000 = n[2 \times 4 + (n-1)5]$$

$$4000 = n(8 + 5n - 5)$$

$$4000 = n(5n + 3)$$

$$4000 = 5n^2 + 3n$$

$$5n^2 + 3n - 4000 = 0$$

$$n = \frac{-3 \pm \sqrt{9 + 80000}}{10}$$

$$n = 28.0 \text{ or } n = -28.6$$

28 terms are needed

Solve using the quadratic formula

Accept positive answer

EXAMPLE 9

Robert starts his new job on a salary of £15000. He is promised a salary increase of £1000 each year, at the end of every year, until he reaches his maximum salary of £25000. Find his total earnings (since appointed) after

a 8 years

b 14 years.

a Total earnings =
$$£15000 + £16000 + ...$$

$$a = 15\,000, d = 1000 \text{ and } n = 8$$

$$S_8 = \frac{8}{2}[30\,000 + 7 \times 1000]$$

Use
$$S_n = \frac{n}{2}[2a + (n-1)d]$$

 $S_8 = £148000$

b Total earnings = £15000 + £16000 + ... + 25000 + 25000 + 25000 + 25000

This time there are 10 years of increases, taking him to the end of his 11th year, and 3 years of the same salary

$$a = 15\,000$$
, $d = 1000$ and $n = 11$

For the first 11 years

$$S_{11} = \frac{11}{2} [30\,000 + 10 \times 1000]$$

Use
$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$

$$S_{11} = £220\,000$$

After that 3 years at £25 000 so add another £75 000

Total earnings = $£220\,000 + £75\,000 = £295\,000$

EXAMPLE 10

Show that the sum of the first *n* natural numbers is $\frac{1}{2}n(n+1)$.

This is an arithmetic series with a = 1, d = 1, n = n In the sequence of natural numbers, the first

term is 1, the difference between terms is 1 and the number of terms in the sequence is n

$$S_n = \frac{n}{2}[2 \times 1 + (n-1) \times 1]$$

$$S_n = \frac{n}{2}[2 + n - 1]$$

Use
$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$

$$S_n = \frac{n}{2}(n+1)$$

$$S_n = \frac{1}{2}n(n+1)$$

You can use Σ to mean 'the sum of'.

For example:

$$\sum_{n=1}^{10} 2n$$
 means the sum of $2n$ from $n=1$ to $n=10$

$$\sum_{n=1}^{10} 2n = 2 + 4 + 6 + 8 + 10 + 12 + 14 + 16 + 18 + 20$$

$$\sum_{n=1}^{10} U_n = U_1 + U_2 + U_3 + U_4 + U_5 + U_6 + U_7 + U_8 + U_9 + U_{10}$$

$$\sum_{r=0}^{10} (2+3r)$$
 means the sum of 2 + 3r from $r=0$ to $r=10$

EXAMPLE 11

Calculate
$$\sum_{r=1}^{20} 4r + 1$$

$$\sum_{r=1}^{20} 4r + 1$$

$$a = 5$$
, $d = 4$ and $n = 20$

$$S_{20} = \frac{20}{2} [2 \times 5 + (20 - 1) \times 4]$$

$$= 10 \times (10 + 19 \times 4)$$

$$= 10 \times 86$$

$$= 860$$

Substitute r = 1, 2, ... to find the terms in the series

This is an arithmetic series

Use
$$S_n = \frac{n}{2}[2\alpha + (n-1)d]$$

EXERCISE 3

SKILLS

THINKING PROBLEM SOLVING

1 Find the sum of these series.

c
$$150 + 145 + 140 + 135 + \dots$$
 (10 terms) **d** $-15 + (-12.5) + (-10) + \dots$ (50 terms)

$$(x + 1) + (2x + 1) + (3x + 1) + ... + (21x + 11)$$

Find how many terms of these series are needed to make the given sum?

$$a 6 + 16 + 26 + \dots$$
, sum = 12550

b
$$7 + 11 + 15 + 19 + \dots$$
, sum = 900

$$-20 + (-17) + (-14) + (-8) + \dots$$
, sum = 848

d
$$29 + 24 + 19 + 14 + \dots$$
, sum = -247491

Find the sum of the first 30 odd numbers.

▶ Ali starts works on an annual salary of \$20 000. His contract will give him an increase of \$120 every 6 months. If he keeps working for 15 years, how much will Ali earn?

Find the sum of the natural numbers up to 200.

► Kumar earns €35 000 a year. He gets an increase of €123 every year on his salary. He works for 20 years and then retires. His pension entitles him to 5% of his overall salary as a monthly allowance. What is his monthly pension allowance?

7 Find the sum of the even numbers from 26 to 126.

8 A jogger is training for a 5 km run. She starts training with 200 m and increases her run by 50 m a day. How many days does it take her to reach 5 km?



Show that the sum of the first 2n natural numbers is n(2n + 1)

10 The sum of the interior angles of a triangle is 180°, a quadrilateral 360° and a pentagon 540° Assuming the pattern continues, find the sum of the interior angles of a dodecagon (12-sided shape).

11 Rewrite these series using sigma Σ notation.

d The multiples of 4 less than 75

12 Prove that the formula of an arithmetic series is: $S_n = \frac{n}{2}(a + L)$

13 Calculate

a
$$\sum_{r=1}^{15} (4r-1)^{r}$$

b
$$\sum_{r=1}^{21} 3(r-1)$$

$$\sum_{n=0}^{12} (3-6n)^n$$

a
$$\sum_{r=1}^{15} (4r-1)$$
 b $\sum_{r=1}^{21} 3(r-1)$ **c** $\sum_{r=1}^{12} (3-6r)$ **d** $\sum_{r=1}^{7} (15r+13)$

14 For what value of n does $\sum_{r=1}^{n} (2r+3)$ exceed 100?

15 For what value of *n* is $\sum_{r=1}^{n} (100 - 6r) < 0$?

Q14 HINT

You will need to solve a quadratic inequality.

IDENTIFY A GEOMETRIC SEQUENCE

Consider the sequence

3, 9, 27, 81 ...

The pattern here is that each term is multiplied by 3.

Consider another sequence

$$3, -6, 12, -24...$$

The pattern here is that each term is multiplied by -2.

Sequences like these are known as geometric sequences.

These sequences are called geometric sequences.

1, 2, 4, 8, 16, ...

100, 25, 6.25, 1.5625, ...

$$2, -6, 18, -54, 162, \dots$$

To get from one term to the next you multiply by the same number each time. This number is called the common ratio, and often represented by r.

KEY POINTS

Common ratio $r=\dfrac{u_2}{u_1}$, where u_1 and u_2 are the first two elements

EXAMPLE 12

Find the common ratios in these geometric sequences.

a 2, 10, 50, 250, ...

- **b** 90, -30, 10, $-\frac{10}{3}$, ...
- a Common ratio = $\frac{10}{2}$ = 5
- **b** Common ratio = $-\frac{30}{90} = -\frac{1}{3}$

The common ratio can be negative and it can be a fraction

EVEDCISE A

1 Which of these are geometric sequences? For the ones that are, find the common ratio.

SKILLS REASONING

c 211, 105.5, 52.75...

a 2, 10, 50, 250...

d 4, 7, 9, 12...

b 3, 6, 18, 34...

e 7, -7, 7, -7...

f 9, 9, 9, 9...

- ., .,., ...

- . 0, 0, 0, 0...
- **g** 120, -30, 7.5, -1.875...
- **h** 61, 59, 57, 55...
- 2 Find the next three terms of these geometric sequences.
 - **a** 113, -339, 1017...

b 20, 80, 320...

 $\frac{2}{9}, \frac{2}{3}, 2...$

d 3, 15, 75...

e 1, −0.25, 0.0625...

- $f = 1, x, x^2...$
- 3 lf 4, p and 16 are the first three terms of geometric sequence, find
 - a the exact value of p.

b the exact value of the 5th term.

FIND THE COMMON RATIO, FIRST TERM AND nth TERM OF A GEOMETRIC SEQUENCE

You can define a geometric sequence using the first term a and the common ratio r.

a.

ar

 ar^2 ,

 ar^3 ,

 ar^{n-1}

First term

Second term

Third term

Fourth term

nth term

Sometimes a geometric sequence is called a geometric progression.

HINT

Look at the relationship between the position of the term in the sequence and the index of the term. You should be able to see that the index of r is one less than its position in the sequence. So the nth term of a geometric sequence $u_n = \alpha r^{n-1}$.

EXAMPLE 13

Find i the 10th term and ii the *n*th term in these geometric sequences.

a 3, 6, 12, 24, ...

b 40, -20, 10, -5, ...

a 3, 6, 12, 24, ...

i $a = 3, r = \frac{6}{3} = 2$ and n = 10

10th term = $3 \times (2)^9$

Use ar^{n-1}

 $= 3 \times 512$

= 1536

ii a = 3, r = 2 and n = n

nth term = $3 \times (2)^{n-1}$

Use ar^{n-1}

b 40, -20, 10, -5

i $\alpha = 40, r = \frac{-20}{40} = -\frac{1}{2}$ and n = 10.

10th term = $40 \times \left(-\frac{1}{2}\right)^9$

Use ar^{n-1}

 $=-\frac{5}{64}$

ii $40 \times \left(-\frac{1}{2}\right)^{n-1}$

 $=5\times8\times\left(-\frac{1}{2}\right)^{n-1}$

 $= 5 \times 2^3 \times (-1)^{n-1} \times \left(\frac{1}{2}\right)^{n-1}$

 $= 5 \times (-1)^{n-1} \times 2^3 \times (2^{-1})^{n-1}$

 $= 5 \times (-1)^{n-1} \times 2^3 \times 2^{-n+1}$

 $= 5 \times (-1)^{n-1} \times 2^{3-n+1}$

 $= 5 \times (-1)^{n-1} \times 2^{4-n}$

Use law of indices $\frac{1}{x^m} = x^{-m}$ and $(x^m)^n = x^{m \times n}$

EXAMPLE 14

The second term of a geometric sequence is 4, and the 4th term is 8. The common ratio of the sequence is greater than 0. Find the exact values of

- a the common ratio
- b the first term
- the 10th term.

a 2nd term = 4,
$$ar$$
 = 4 (1)

4th term = 8,
$$ar^3$$
 = 8 (2)

using *n*th term= ar^{n-1} with n=2 and 4

(2)
$$\div$$
 (1) gives:

$$r^2=\frac{8}{4}=2$$

$$r = \sqrt{2}$$

So the common ratio is $\sqrt{2}$

b Substitute the common ratio back into (1) to find: $a\sqrt{2} = 4$

$$a=\frac{4}{\sqrt{2}}$$

$$=\frac{4\sqrt{2}}{2}$$

$$=2\sqrt{2}$$

To rationalise $\frac{4}{\sqrt{2}}$, multiply top and bottom by $\sqrt{2}$

So the first term is $2\sqrt{2}$

c The 10th term is:

$$ar^9 = 2\sqrt{2} \times (\sqrt{2})^9$$

$$=2\times (\sqrt{2})^{10}$$

$$= 2 \times 2^{5}$$

$$= 2^{6}$$

$$= 64$$

Substitute the values of a and r back into ar^{n-1} with n = 10

Use
$$(\sqrt{2})^{10} = (2\frac{1}{2})^{10} = 2\frac{1}{2} \times 10 = 2^{5}$$

EXAMPLE 15

The numbers 3, x and x + 6 form the first three terms of a positive geometric sequence.

Find

- a the possible values of x
- b the 10th term of the sequence.
- a The sequence is a geometric sequence only if $\frac{U_2}{U_4} = \frac{U_3}{U_2}$

$$\frac{x}{3} = \frac{x+6}{x}$$

Cross multiply

$$x^2 = 3(x + 6)$$

$$x^2 = 3x + 18$$

$$x^2 - 3x - 18 = 0$$

Factorise

$$(x-6)(x+3)=0$$

The solutions are x = 6, x = -3, but this is a positive sequence so the only answer is x = 6.

b The 10th term is ar^9 . Here a=3 and $r=\frac{U_2}{U_1}=\frac{6}{3}=2$.

Use the formula for the nth term $u_n = \alpha r^{n-1}$ with

$$ar^9 = 3 \times 2^9 = 3 \times 512 = 1536$$

n = 10, a = 3, r = 2

EXERCISE 5

SKILLS

PROBLEM SOLVING

- 1 Find the 5th, 7th and nth term of this sequence:
 - **a** 3, 6, 12...
- **b** 160, 80, 40...
- **c** 2, (-6), 18...
- **d** (-1), (-2), (-4)...
- 2 Find the common ratio and the first term in the geometric sequence where:
 - a The 2nd term is 6 and the 4th term is 24.
 - **b** The 3rd term is 60 and the 6th term is -480.
 - \mathbf{c} The 3rd term is -7.5 and the 6th term is 0.9375.
 - d The 4th term is 4 and the 7th term is 32.
- 3 \blacktriangleright The *n*th term of a geometric sequence is $3(5^{n-1})$
 - a Find the first term.

- **b** Find the 5th term.
- 4 Given that the first term of a geometric sequence is 8 and the third term is 2, find the possible values of for the 6th term.
- 5 \blacktriangleright The expressions p-6, 2p and p^2 form the first three terms of a geometric sequence. Find the possible values of the first term.

Q5 HINT

Calculate different expressions for the common ratio. Then form and solve an equation in *p*.

FIND THE SUM OF A GEOMETRIC SERIES

The general rule for the sum of a geometric series is:

$$S_n = \frac{\alpha(r^n - 1)}{r - 1}$$
 or $S_n = \frac{\alpha(1 - r^n)}{1 - r}$

Note: You will not be asked to prove these formulae.

EXAMPLE 16

Find the sum of these series.

a The series is 2 + 6 + 18 + 54 + ... (for 10 terms)

So
$$a = 2$$
, $r = \frac{6}{2} = 3$, $n = 10$

So
$$S_{10} = \frac{2(3^{10} - 1)}{3 - 1} = 59\,048$$

b The series is 1024 - 512 + 256 - 128 + ... + 1

So
$$a = 1024$$
, $r = -\frac{512}{1024} = -\frac{1}{2}$, and n th term is 1.

Solve $ar^{n-1} = 1$ to find n

$$1024 \times \left(-\frac{1}{2}\right)^{n-1} = 1$$

$$1024 = (-2)^{n-1}$$

1024 is positive so $(-2)^{n-1}$ must be positive, and equal to 2^{n-1}

$$2^{n-1} = 1024$$

$$(n-1)\lg 2 = \lg (1024)$$

$$n-1 = \frac{\lg(1024)}{\lg(2)} = 10$$
 $n = 11$

If you noticed that $1024 = 2^{10}$ you could have solved this without using logarithms

So:

$$S_{11} = \frac{1024 \left[1 - \left(-\frac{1}{2} \right)^{11} \right]}{1 - \left(-\frac{1}{2} \right)}$$

$$= \frac{1024 \times \left(1 + \frac{1}{2048} \right)}{1 + \frac{1}{2}}$$

$$= \frac{\frac{2049}{2}}{\frac{3}{2}} = 683$$

EXAMPLE 17

An investor invests £2000 on January 1st each year in a savings account that guarantees him 4% per annum for life. If interest is calculated on the 31st of December each year, how much will be in the account at the end of the 10th year?

End of year 1, amount = 2000×1.04

Start of year 2, amount = $2000 \times 1.04 + 2000$

End of year 2, amount = $(2000 \times 1.04 + 2000) \times 1.04$

$$= 2000 \times 1.04^2 + 2000 \times 1.04$$

A rate of 4% means × 1.04

Every new year he invests £2000

At the end of every year the total amount in the account is multiplied by 1.04

Start of year 3,

amount =
$$(2000 \times 1.04^2 + 2000 \times 1.04 + 2000) \times 1.04$$

= $2000 \times 1.04^3 + 2000 \times 1.04^2 + 2000 \times 1.04$

So by end of year 10,

Look at the values for the end of year 3 and extend this for 10 years

amount =
$$2000 \times 1.04^{10} + 2000 \times 1.04^{9} + ... + 2000 \times 1.04$$

= $2000 \times (1.04^{10} + 1.04^{9} + ... + 1.04)$ This is a given by the second of the seco

This is a geometric series. Substitute a = 1.04, $r = 1.04, n = 10 \text{ in } S = \frac{a \times (r^n - 1)}{r - 1}$

EXAMPLE 18

Find the least value of n such that the sum of 1 + 2 + 4 + 8 + ... to n terms would exceed 2000000.

Sum to *n* terms is $S_n = 1 \times \frac{2^n - 1}{2 - 1} = 2^n - 1$

Substitute a = 1, r = 2 into $S_n = \frac{a \times (r^n - 1)}{r - 1}$

If this is to exceed 2000000 then:

$$2^n - 1 > 2000000$$

$$2^n > 2000001$$

 $n \lg(2) > \lg(2000001)$

$$n > \frac{\lg(2\,000\,001)}{\lg(2)}$$

Use laws of logarithms: $\lg(a^n) = n \lg a$

Round up to the nearest integer

It needs 21 terms to exceed 2000000

Alternative method: $2^n - 1 > 2000000$

 $2^n > 2000001$

HINT

 $n \log_2 2 > \log_2 2 000 001$ $\log_2 2 = 1$

 $n > \log_2 2000001$

n > 20.9

It needs 21 terms to exceed 2 000 000

EXAMPLE 19

Find
$$\sum_{r=1}^{10} (3 \times 2^r)$$

$$S_{10} = \sum_{r=1}^{10} (3 \times 2^r)$$

$$= 3 \times 2^1 + 3 \times 2^2 + 3 \times 2^3 + \dots + 3 \times 2^{10}$$

$$= 3[2^1 + 2^2 + 2^3 + \dots + 2^{10}]$$

$$= 3 \times 2(\frac{2^{10} - 1}{2 - 1})$$

So $S_{10} = 6138$

The quantity in parentheses is a geometric series with a = 2, r = 2, n = 10

Use
$$S_n = \frac{a \times (r^n - 1)}{r - 1}$$

SKILLS

CRITICAL THINKING 1 Find the sum of these geometric series:

a 1 + 3 + 9 + 27 + ... (7 terms)

e
$$\sum_{r=1}^{7} 5^r$$

g
$$\sum_{r=1}^{11} 6 \times \left(\frac{1}{0.75}\right)^r$$


b
$$50 + (-25) + 12.5 + (-6.25) + ... (10 terms)$$

$$\int_{r=1}^{1} (2 \times 2^r)$$

h
$$\sum_{r=1}^{6} 3(-7^r)$$

- 2 Legend has it that the inventor of chess was asked to name his reward for the invention by the ruling king of the time. He asked for 1 grain of rice on the first square of the chess board, 2 grains of rice of the second square, 4 grains of rice on the third square and so on until all the 64 squares were covered in rice. How many grains of rice could the inventor claim?
- 3 A savings scheme is offering a rate of interest of 3.5% per annum for the lifetime of the plan. Ji Won wants to save up to \$20000. She works out that she can afford to save \$500 every year. How many years will it be until she has saved up \$20 000?
- The rate of tax in India operates on a flat rate system. If you earn more than 50 000 rupees then you pay a flat rate of tax of 20%. If you earn less than 50000 rupees then you pay a flat tax rate of 15%. You earn 3750 rupees in the first month, and for each month after that your salary increases by 3% of the previous month's salary. How much tax would you pay that year?
- 5 You are offered a job that lasts for 5 weeks and you can choose your salary. You have two options:
 - You get paid \$100 a day for the first day, \$200 for the second day, \$300 for the third day and so on. Each day your salary increases by \$100.

- b You get paid 1 cent for the first day, 2 cents for the second day, 4 cents for the third day and so on. Each day you are paid double of what you were paid for the previous day. Which salary would you choose?
- 6 Abishek is sponsored to cycle from Cape Town to Durban (1600 kilometres away) over a number of days. Abishek cycles 15 km on day 1 and increases his distance by 15% each day. How long will it take him to complete the challenge?





8 The first and last terms of a geometric series are 2 and 2048 respectively. The sum of the series is 2730. Find the number of terms and the common ratio.

FIND THE SUM TO INFINITY OF A CONVERGENT GEOMETRIC SERIES

Consider the series S = 3 + 1.5 + 0.75 + 0.375 + 0.1875 + ...

No matter how many terms of the series you take, the sum never exceeds a certain number.

This number is called the limit of the sum, or more often, the sum to infinity.

You can find out what this limit is.

As
$$a = 3$$
 and $r = \frac{1}{2}$, then $S = \frac{1(1 - r^n)}{1 - r} = \frac{3(1 - (\frac{1}{2})^n)}{1 - \frac{1}{2}} = 6(1 - (\frac{1}{2})^n)$

If you replace n with certain values to find the sum you find that:

when
$$n = 3$$
, $S_3 = 5.25$

when
$$n = 5$$
, $S_5 = 5.8125$

when
$$n = 10$$
, $S_{10} = 5.9994$

when
$$n = 20$$
, $S_{20} = 5.999994$

You can see that as n gets larger, S becomes closer and closer to 6.

You say that this infinite series is convergent, and has a sum to infinity of 6. Convergent means that the series tends towards a specific value as more terms are added.

Not all series converge. The reason that this one does is that the terms of the sequence are getting smaller.

This happens because -1 < r < 1. The sum to infinity of a geometric series exists only if -1 < r < 1.

$$S_n = \frac{a(1-r^n)}{1-r}$$

If
$$-1 < r < 1$$
, $r^n \to 0$ as $n \to \infty$, so that:

$$S_{\infty} = \frac{a(1-0)}{1-r} = \frac{a}{1-r}$$

HINT

|r| < 1 means all numbers whose actual size, irrespective of sign, is less than 1.

HINT

You can write 'the sum to infinity' as S∞.

> The sum to infinity of a geometric series is $\frac{a}{1-r}$ if |r| < 1**KEY POINTS**

EXAMPLE 20

Find the sum to infinity of these series.

SKILLS

CRITICAL

a 40 + 10 + 2.5 + 0.625 + ...

b
$$1 + \frac{1}{p} + \frac{1}{p^2} + \frac{1}{p^3} + \dots$$
, where p is a positive number.

a
$$40 + 10 + 2.5 + 0.625 + ...$$

In this series $a = 40$, $r = \frac{10}{40} = \frac{1}{4}$

Always write down the values of
$$a$$
 and r , using $\frac{U_2}{U_1}$ for r

Use the formula $S_n = \frac{a(1-r^n)}{1-r}$, with n=4

Solve the two equations simultaneously

$$-1 < r < 1$$
 , so S_∞ exists:
$$S_\infty = \frac{a}{1-r} = \frac{40}{1-\frac{1}{1}} = \frac{40}{\frac{3}{2}} = \frac{160}{3}$$

Substitute
$$a=40$$
, $r=\frac{10}{40}=\frac{1}{4}$ into $S_{\infty}=\frac{a}{1-r}$

b
$$1 + \frac{1}{p} + \frac{1}{p^2} + \frac{1}{p^3} + \dots$$

In this series
$$a = 1$$
, $r = \frac{U_2}{U_1} = \frac{\frac{1}{p}}{1} = \frac{1}{p}$

$$S$$
 will exist if $\left|\frac{1}{p}\right| < 1$, so S_{∞} exists only if $p > 1$

If
$$p > 1$$
, $S_{\infty} = \frac{1}{1 - \frac{1}{p}} = \frac{p}{p - 1}$

Multiply top and bottom by p

EXAMPLE 21

The sum to 4 terms of a geometric series is 15 and the sum to infinity is 16.

- Find the possible values of r.
- Given that the terms are all positive, find the first term in the series.

a
$$S_4 = 15$$

$$\frac{a(1-r^4)}{1-r} = 15 \quad (1)$$

$$\frac{a}{1-r} = 16$$
 (2)

$$16(1-r^4)=15$$

$$16 - 16r^4 = 15$$

$$16r^4 = 1$$

$$r^4 = \frac{1}{16}$$

$$r=\pm\frac{1}{2}$$

b As all terms are positive, r is $+\frac{1}{2}$

Substitute $r = \frac{1}{2}$ back into equation (2) to find a:

$$\frac{a}{1-\frac{1}{2}}=16$$

$$16(1-\frac{1}{2})=a$$

$$a = 8$$

The first term in the series is 8.

SKILLS CRITICAL THINKING

Find the sum to infinity, if it exists, of these series.

d
$$1-2y+4y^2-8y^3+...$$

2 The first three terms of a geometric series are 4 + 2 + 1

What is the common ratio of this series?

b Find the sum to infinity.

3 The first three terms of a geometric series are 100 + 90 + 81

What is the common ratio of this series?

b Find the sum to infinity.

Find $\sum_{n=0}^{\infty} 3(0.5)^n$

5 ▶ Find the sum to infinity of the geometric series having a second term of -9 and a fifth term of $\frac{1}{2}$

6 A ball is dropped from a height of 6 metres. Suppose the ball rebounds $\frac{2}{3}$ of the height from which it falls. Find the total distance of travelled by the ball.



7 > The lengths of the radii of circles form a geometric series. The radius of the first circle is 15cm. The radius decreases by $\frac{4}{5}$ of the length of the previous circle. Determine the total area of all the circles in the series. Give your answer in terms of π .

8 A factory contributes £1 million into a small city. It is estimated that 75% of that money is respent back into the community. Economists assume that the money is re-spent again and again at a rate of 75%. Find an approximation for the total money re-spent from this original £1 million.

The sum to the three terms of a geometric series is 9 and its sum to infinity is 8. What can you deduce about the common ratio and why? Find the first term and common ratio.

[2]

[4]

- The third and fifth terms of an arithmetic series are 67 and 121.
 - a Find the first term.
 - b Find the common difference. [2]
 - Find the sum of the first 25 terms of the series. [2]
- The first three terms of an arithmetic series are 1, (2q-5) and 8.6
 - a Find the value of q. [2]
 - b Find the 16th term of the series. [4]
- 3 Calculate $\sum_{r=3}^{20} (5r-1)$
- The fifth and twentieth terms of an arithmetic series are 18 and 63.
 - a Find the first term and common difference.
 - **b** Given that the sum of the n terms of the series is 270, find the value of n.
- The sixth and fifteenth terms of an arithmetic series are 277 and 214.
 - a Find the common difference and first term. [3]
 - a Find the possible values of n. [4]
 - a Find the maximum value of S_n , the sum of the first n terms of the series. [3]
- A geometric series has a common ratio of -2 and the first term of 3.
 - a Find the sum of the first ten terms of the series [2]
 - b Show that the sum of the first eight positive terms of the series is 65 535. [4]
- 7 Evaluate $\sum_{r=3}^{12} r^2$
- The second term of a geometric series S is $\sqrt{2}-2$ and the third term is $3-\sqrt{2}-4$
 - a Find the exact value of the common ratio and the first term of the series. [5]
 - **b** Find the sum to infinity of *S* and give a reason why the series is convergent. [4]
- The second term of a geometric series is 80 and the fifth term is 5.12
 - a Show that the common ratio of the series is 0.4

Calculate:

- b The first term of the series. [2]
- The sum to infinity of the series, giving your answer as an exact fraction. [2]
- d The difference between the sum to infinity of the series and the sum of the first 14 terms of the series,
 - giving your answer in the form $a \times 10^n$ where $1 \le a \le 10$ and n is an integer.

10 A geometric series is $a + ar + ar^2 + ...$

Prove that the sum of the first n terms of this series is given by $S_n = \frac{a(1-r^n)}{1-r}$. [4]

- The first three terms of a geometric series are (p + 4), p and (2p 15) respectively, where p is a positive constant.
 - a Show that $p^2 7p 60 = 0$ [3]
 - **b** Hence show that p = 12
 - c Find the common ratio. [2]
 - d Find the sum to infinity. [2]

CHAPTER SUMMARY: CHAPTER 5

All arithmetic series can be written in the form:

$$a + (a+d) + (a+2d) + (a+3d) + ... + (a+(n-1)d) + ...$$

First term Second term Third term Fourth term nth term

- The nth term of an arithmetic sequence is a + (n 1)d, where a is the first term and d is the common difference.
- The sum of an arithmetic series is:

$$S_n = \frac{n}{2}[2\alpha + (n-1)d] \text{ or } S_n = (\alpha + L)$$

where a is the first term, d is the common difference, n is the number of terms and L is the last term in the sequence.

- You can use the symbol Σ to indicate 'sum of'. Σ is used to write a series in a quick and concise way. For example, $\sum_{r=1}^{500} (2r+50) = 52+54+56+...+1050$
- In a geometric sequence you can get from one term to another by multiplying by a constant called the common ratio.
- The formula for the *n*th term is ar^{n-1} where a is the first term and r is the common ratio.
- The formula for the sum of the n terms of a geometric series is

$$S_n = \frac{\alpha(1 - r^n)}{1 - r}$$
 or $S_n = \frac{\alpha(r^n - 1)}{r - 1}$

The sum to infinity exists if -1 < r < 1 and $S_{\infty} = \frac{a}{1-r}$.

CHAPTER 6



6 THE BINOMIAL SERIES

The **binomial series** has a long history that stretches round the entire world.

Both the ancient Greeks and the ancient Hindus were fascinated by the different ways in which you can select objects from a group. The binomial theorem itself was first proved by Persian mathematician Al-Karaji in the 11th century, who also described its triangular pattern. Another Persian, Omar Khayyam, took the formula to higher powers. The binomial expansion of small degrees was known in the 13th century to Yang Hui of China.

In 1544, the German mathematician Michael Stifel introduced the term 'binomial coefficient' and showed how to express $(1 + a)^n$ and $(1 + a)^{n-1}$ using **Pascal's triangle**. Sir Isaac Newton of England wrote down the generalised binomial theorem, valid for any rational exponent.



Sir Isaac Newton (1642–1727)

LEARNING OBJECTIVES

- Use $\binom{n}{r}$ to work out the coefficients in the binomial expansion
- Use the binomial expansion to expand $(1 + x)^n$
- Determine the range of values for which x is true and valid for the expansion.

STARTER ACTIVITIES

Consider these mathematical statements.

$$(a + b)^{0} = 1$$

$$(a + b)^{1} = 1a + 1b$$


$$(a + b)^{2} = 1a^{2} + 2ab + 1b^{2}$$

$$(a + b)^{3} = 1a^{3} + 3a^{2}b + 3ab^{2} + 1b^{3}$$

$$(a + b)^{4} = 1a^{4} + 4a^{3}b + 6a^{2}b^{2} + 4ab^{3} + 1b^{4}$$

There is a pattern in the coefficients. Can you spot it?

The coefficients form a pattern known as Pascal's triangle:



Note: To get from one line to another you add the adjacent numbers, as shown in the diagram.

EXAMPLE 1

By using Pascal's triangle find the expansion of

$$(2x + y)^3$$

b
$$(3a - 2b)^5$$

a
$$(2x + y)^3$$
 is in the form $(a + b)^n$ with $a = 2x$, $b = y$ and $n = 3$ so use $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ and substitute in for a and b

These numbers have come from the triangle in the key point above

$$(2x + y)^3 = (2x)^3 + 3(2x)^2(y) + 3(2x)(y)^2 + (y)^3$$
$$= 8x^3 + 12x^2y + 6xy^2 + y^3$$

Simplify

HINT

Watch out for negative powers.

- **b** $(3a-2b)^4$ is in the form $(a+b)^n$ with a=3a, b=-2b and n=4 so use $(a+b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$ and substitute in for a and b $(3a - 2b)^4 = (3a)^4 + 4(3a)^3(-2b) + 6(3a)^2(-2b)^2 + 4(3a)(-2b)^3 + (-2b)^4$ $= 81a^4 - 216a^32b + 216a^2b^2 - 96ab^3 + 16b^4$ Simplify
- 1 Expand these expressions using Pascal's triangle.

$$(3 + x)^3$$

b
$$(5 + 2x)^3$$

$$(2 + x)^4$$

2 Find the coefficient of x^3 in these expansions.

$$(5 + x)^7$$

b
$$(6 + 3x)^5$$

TO WORK OUT THE COEFFICIENTS IN THE BINOMIAL EXPANSION

For a binomial expression with a large power, e.g. $(1 + x)^{32}$, Pascal's triangle contains a huge number of rows. A better method is to use the binomial theorem. This enables you to expand $(a + b)^n$ into increasing powers of b and decreasing powers of a. You will look at expanding expressions of the form $(a + b)^2$, $(a + b)^3$, ..., $(a + b)^{32}$, ... when the power is a positive whole number.

KEY POINTS

is sometimes written as ${}^{n}C_{r}$ where n is a positive integer. This can be evaluated on most calculators. ${}^{n}C_{r}$ is often pronounced 'n choose r'.

The binomial expansion, when n is a positive integer, is a formula for

$$(a + b)^n = (a + b) \times \ldots \times (a + b)$$
 (n times)

$$(a + b)^n = {^n}\mathsf{C}_0 a^n + {^n}\mathsf{C}_1 a^{n-1} b^1 + {^n}\mathsf{C}_2 a^{n-2} b^2 + \ldots + {^n}\mathsf{C}_{n-1} a^1 b^{n-1} + {^n}\mathsf{C}_n b^n$$

$$(a+b)^n = \binom{n}{0}a^n + \binom{n}{1}a^{n-1}b^1 + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{n-1}a^1b^{n-1} + \binom{n}{n}b^n$$

$$\frac{n!}{(n-r)!r!}$$
 (*r* is effectively the 'position' in the expansion)

$$\frac{n!}{(n-1)!1!}$$
 (*n* is the power which the bracket is raised to)

EXAMPLE 2

Use the binomial theorem to find the expansion of $(2x + y)^4$

$$(2x + y)^4 = {4 \choose 0}(2x)^4 + {4 \choose 1}(2x)^3(y) + {4 \choose 2}(2x)^2(y)^2 + {4 \choose 3}(2x)^1(y)^3 + {4 \choose 4}(y)^4 \quad n = 4, \ \alpha = 2x \text{ and } b = y$$

$$= (2x)^4 + (4)(2x)^3(y) + (6)(2x)^2(y)^2 + (4)(2x)^1(y)^3 + (y)^4$$

$$\binom{n}{r} = \frac{n!}{(n-r)!r!} \text{ so } \binom{4}{0} = \frac{4!}{4!0!} = 1, \binom{4}{1} = \frac{4!}{3!1!} = 4, \binom{4}{2} = \frac{4!}{2!2!} = 6, \binom{4}{3} = \frac{4!}{1!3!} = 4, \binom{4}{4} = \frac{4!}{0!4!} = 1$$

$$= 16x^4 + 32x^3y + 24x^2y^2 + 8xy^3 + y^4$$

EXAMPLE 3

Use the binomial theorem to find the expansion of $(3 - 2x)^5$

$$(3-2x)^5 = \binom{5}{0}(3)^5 + \binom{5}{1}(3)^4(-2x) + \binom{5}{2}(3)^3(-2x)^2 + \binom{5}{3}(3)^2(-2x)^3 + \binom{5}{4}(3)(-2x)^4 + \binom{5}{5}(-2x)^5 + \binom{5}{1}(3)(-2x)^4 + \binom{5}{1}(3)(-2x)^4$$

$$n = 5$$
, $a = 3$ and $b = -2x$

$$= (3)^5 + (5)(3)^4(-2x) + (10)(3)^3(-2x)^2 + (10)(3)^2(-2x)^3 + (5)(3)(-2x)^4 + (-2x)^5(-2x)^4 + (-2x)^5(-2x)^5(-2x)^5 + (-2x)^5(-2$$

$${n \choose r} = \frac{n!}{(n-r)!r!} \text{ so } {5 \choose 0} = \frac{5!}{5!0!} = 1, {5 \choose 1} = \frac{5!}{4!1!} = 5, {5 \choose 2} = \frac{5!}{3!2!} = 10,$$
$${5 \choose 3} = \frac{5!}{3!2!} = 10, {5 \choose 4} = \frac{5!}{4!1!} = 5, {5 \choose 5} = \frac{5!}{5!0!} = 1$$

$$= 243 - 810x + 1080x^2 - 720x^3 + 240x^4 - 32x^5$$

EXERCISE 1

- 1 Expand these expressions using the Binomial Expansion.

- **a** $(3x + y)^3$ **b** $(m n)^4$ **c** $(1 + 3x)^3$ **d** $\left(1 + \frac{1}{4}y\right)^4$ **e** $(3y + 4x)^3$
- **2** Find the term in x^4 of these expansions.
- **a** $(3x+6)^6$ **b** $(4+5x)^7$ **c** $(1+x)^{21}$ **d** $(1-x)^{30}$
- 3 b Using the binomial theorem, or otherwise, find the first four terms of these expressions.

 - **a** $(3x+1)^4$ **b** $\left(1-\frac{1}{4}y\right)^7$ **c** $(7-b)^3$ **d** $(3x^2+2)^{10}$
- 4 The coefficient of x^2 in the binomial expansion of $\left(1 + \frac{2}{5}x\right)^n$, where n is a positive integer, is 1.6
 - a Find the value of n.
 - **b** Use your value of n to find the coefficient of the x^4 in the expansion.

USE THE BINOMIAL EXPANSION TO EXPAND $(1 + x)^n$

There is a shortened version of the expansion when one of the terms is 1.

Whatever power 1 is raised to, it will be 1, and can therefore be ignored.

The coefficients give values from Pascal's triangle. ${}^{n}C_{1}$ cancels to n, ${}^{n}C_{2}$ cancels to $\frac{n(n-1)}{2!}$ and so on.

KEY POINTS

$$(1+x)^n = \binom{n}{0} 1^n + \binom{n}{1} 1^{n-1} x^1 + \binom{n}{2} 1^{n-2} x^2 + \ldots + \binom{n}{r} 1^{n-r} x^r + \ldots + \binom{n}{n} x^n$$

Using the expression for the binomial coefficients

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

For example, if n = 4

$$(1+x)^4 = 1 + 4x + 6x^2 + 4x^3 + x^4$$

EXAMPLE 4

Find the first four terms in the binomial expansion of

$$(1 + 3x)^6$$

b
$$(1-2x)^3$$

a
$$(1 + 3x)^6 = 1 + 6(3x) + \frac{6(5)}{2!}(3x)^2 + \frac{6(5)(4)}{3!}(3x)^3 + \dots$$

= $1 + 6(3)x + 15(3x)^2 + 20(3x)^3$
= $1 + 18x + 135x^2 + 540x^3$

Substitute in n = 6 and x = 3x

Simplify

b
$$(1-2x)^3 = 1 + (3)(-2x) + (3)(2)\frac{(-2x)^2}{2} + (3)(2)(1)\frac{(-2x)^3}{6}$$

 $= 1 - 6x + 12x^2 - 8x^3$

Substitute in n = 3 and x = -2x

Simplify

including x parts.

important to put

brackets around the expressions

EXAMPLE 5

HINT

It is very

Find the coefficient of

in the binomial expansion of $\left(1 - \frac{1}{2}x\right)^6$

a
$$\left(1 - \frac{1}{2}x\right)^6 = {6 \choose 3} \left(-\frac{1}{2}x\right)^3$$

= $-\frac{5}{2}x^3$

so the coefficient of x^3 is $-\frac{5}{2}$

b
$$\left(1 - \frac{1}{2}x\right)^6 = {6 \choose 5} \left(-\frac{1}{2}x\right)^5$$

= $-\frac{3}{16}x^5$

$${}^{6}C_{3} = {6 \choose 3} = \frac{6!}{3!3!}$$

You do not need the entire expansion, just the required terms.

HINT

so the coefficient of x^5 is $-\frac{3}{16}$

CHAPTER 6

EXERCISE 2

Use the binomial expansion to find the first four terms of these series.

$$(1+x)^9$$

b
$$(1-3y)^5$$

$$(1 + 4x)^2$$

d
$$(1 + \frac{x}{2})^9$$

$$(1-3x)^7$$

a
$$(1+x)^9$$
 b $(1-3y)^5$ **c** $(1+4x)^7$ **e** $(1-3x)^7$ **f** $\left(1-\frac{1}{3}x\right)^6$ **g** $\left(1-\frac{x}{4}\right)^6$

g
$$\left(1-\frac{x}{4}\right)^6$$

When $(1-5x)^q$ is expanded the coefficient of x is 40. Given than q>0, find

- a the constant q
- b the coefficient of x^2
- the coefficient of x^3 .

3 \triangleright a Find the first four terms of $(1-x)^8$.

b By substituting a suitable value of x into your answer for part a, obtain an estimate for 0.99^8

4 **a** Find the first four terms of $\left(1 + \frac{x}{8}\right)^8$

b By substituting a suitable value of x into your answer for part a, obtain an estimate for 1.026

5 Find the coefficient of x^3 in these expressions

$$(1-x)^6$$

b
$$(1 + x)^9$$

$$(1 + 2x)^7$$

6 When $\left(1-\frac{2}{3}x\right)^p$ is expanded, the coefficient of x is -4. Given than q>0, find

a the constant p

b the coefficient of x^2

• the coefficient of x^3 .

DETERMINE THE RANGE OF VALUES FOR WHICH $oldsymbol{x}$ is true and valid for an **EXPANSION**

The binomial expansion is shown below. When n is a positive integer, the expansion is finite and exact. This is not the case when n is negative or a fraction. When n is not a positive integer the series will be infinite. In this case the expansion $(1 + x)^n$ will be valid only in the range -1 < x < 1. This can also be written using the **modulus** function as |x| < 1

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

EXAMPLE 6

Use the binomial expansion to find the first four terms of $\frac{1}{1+2r}$

$$\frac{1}{1+2x} = (1+2x)^{-1}$$
Rewrite $\frac{1}{1+2x}$ in index form
$$= 1 + (-1)(2x) + \frac{(-1)(-2)}{2}(2x)^2 + \frac{(-1)(-2)(-3)}{6}(2x)^3 + \dots$$

Use the formula $(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3$ with n = -1 and x = 2x

$$= 1 - 2x + 4x^2 - 8x^3 + \dots$$

The expansion is infinite, but the first four terms can be used as an approximation for $\frac{1}{1+2r}$ provided that |2x| < 1. The approximation works well for 2x close to 0, but not as well if 2x is close to 1, and not at all if 2x Is larger than 1.

KEY POINTS

When n is not a positive integer none of the (n-r) terms in the coefficients will be equal to zero and so the series will continue and be infinite. In this case the expansion of $(1 + x)^n$ will only be valid for values of x in the range -1 < x < 1. This is sometimes written using the modulus function as |x| < 1

EXAMPLE 7

Use the binomial expansion to find the first three terms of

$$\sqrt[3]{1-2x}$$

$$\frac{1}{(1+4x)^3}$$

and state the range of values of x for which the expressions are valid.

a
$$\sqrt[3]{1-2x} = 1-2x^{\frac{1}{3}}$$
 Rewrite in index form
$$= 1+(\frac{1}{2})(-2x)+(\frac{1}{2})(-\frac{2}{2})(-\frac{2}{2}x)^{2}+(\frac{1}{2})(-\frac{2}{2})(-\frac{5}{2})(-\frac{5}{2}x)^{3}$$

$$= 1 + \left(\frac{1}{3}\right)(-2x) + \left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\frac{(-2x)^2}{2} + \left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\left(-\frac{5}{3}\right)\frac{(-2x)^3}{6}$$

Use
$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3$$
 with $n = \frac{1}{3}$, $x = -2x$

$$=1-\frac{2}{3}x-\frac{4}{9}x^2-\frac{40}{81}x^3+\dots$$

The range of validity is given by |2x| < 1, so that $|x| < \frac{1}{2}$

In this case the 'x' term is 2x

Rewrite in index form

b
$$\frac{1}{(1+4x)^3} = (1+4x)^{-3}$$

= 1 + (-3)(4x) + (-3)(-4) $\frac{(4x)^2}{2}$ + (-3)(-4)(-5) $\frac{(4x)^3}{6}$

Use
$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3$$
 with $n = -3$, $x = 4x$

$$= 1 - 12x + 96x^2 - 640x^3 + \dots$$

The range of validity is given by |4x| < 1, so that $|x| < \frac{1}{4}$

In this case the 'x' term is 4x

EXAMPLE 8

Find the binomial expansion of $\sqrt{1-2x}$ and by using x=0.01, find and estimate of $\sqrt{2}$.

$$\sqrt{1-2x} = (1-2x)^{\frac{1}{2}}$$

$$= 1 + \left(\frac{1}{2}\right)(-2x) + \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\frac{(-2x)^2}{2} + \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\frac{(-2x)^3}{6}$$

Rewrite in index form

Use
$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3$$
, in this case with $n = \frac{1}{2}$ and $x = 2x$

$$= 1 - x - \frac{1}{2}x^2 - \frac{1}{2}x^3$$

$$\sqrt{0.98} = 1 - 0.01 - 0.00005 - 0.0000005$$

Substitute x = 0.01

$$\sqrt{\frac{98}{100}} = 0.9899495$$

Rewrite 0.98 as a fraction

$$\frac{7\sqrt{2}}{10} = 0.9899495$$

Calculate the square root of the numerator and denominator separately

$$\sqrt{2} = \frac{9.899495 \times 10}{7}$$

Multiply by 10 and divide by 7

= 1.414213571

EXERCISE :

1 \triangleright Expand these expressions up to and including x^3 , and state the values of x for which the expansion is valid where appropriate.

a
$$(1 + 2x)^{-3}$$

$$\frac{1}{(1-3x)}$$

$$\frac{1}{\sqrt{4+6}x}$$

2 Find the coefficient of the x^3 term, and state the values of x for which the expansion is valid.

$$\mathbf{a} \quad \frac{1}{1+x}$$

b
$$\sqrt{1 + 2x}$$

$$\frac{1}{(1-3x)}$$

d
$$\sqrt[3]{1-x}$$

3 \triangleright a Expand $(1 + 3x)^{-1}$, $|x| < \frac{1}{3}$ in ascending powers of x up to and including the term in x^3 .

b Hence, or otherwise, find an expression for $\frac{1-x}{1+3x}$, $|x| < \frac{1}{3}$ in ascending powers of x up to and including the term in x^3 .

4 In the expansion of $(1 + bx)^{-\frac{1}{2}}$ the coefficient of x^2 is 24. Find the possible values of the constant b and the coefficient of x^3 .

5 **a** Find the binomial expansion of $\sqrt[3]{(1+6x)}$ in ascending powers of x up to and including the term in x^3 .

b By substituting x = 0.004 find an approximation to $\sqrt[3]{2}$. By comparing it to the exact value, comment on the accuracy of your approximation.

6 Find the coefficient of the x^3 in $\frac{1+3x}{(1+2x)^3}$

EXAM PRACTICE: CHAPTER 6

- Find the first 4 terms of the binomial expansion, in ascending powers of x, of $\left(1 + \frac{x}{4}\right)^t$ giving each term in its simplest form. [5]
- a Write down in ascending powers of x, up to and including x^3 , the expansion of $(3 + ax)^5$ where a is a non-zero constant. [3]
 - **b** Given that the coefficient of x^2 is double the coefficient of x, find the value of a. [4]
- The coefficient of x^2 in the expansion of $\left(1-\frac{x}{2}\right)^n$, where n is a positive integer, is 9. [3]
- Show that $\sqrt{\frac{1+x}{1-x}} \approx 1 + x + \frac{1}{2}x^2$, |x| < 1. [7]
- f(x) = $\frac{1}{\sqrt{4+x}}$
 - a Use the binomial expansion to expand f(x) in ascending powers of x up to and including x^3 . [4]
 - **b** State for what values of *x* the expansion is valid. [1]
- **a** Find in ascending powers of x, up to and including x^3 , the expansion of $(1 + 3x)^{\frac{1}{2}}$. [3]
 - By substituting x = 0.01 in the expansion, find an approximation to $\sqrt{103}$. [4]
 - **c** By comparing to the exact value, comment on the accuracy. [2]
- Find the coefficient of the x^3 term of $\frac{1+3x}{1+x}$. [3]
- $f(x) = \sqrt{1 8x}$
 - a Use the binomial expansion to expand f(x) in ascending powers of x up to and including x^3 . [4]
 - **b** State for which values of *x* the expansion is valid. [1]
 - **c** By substituting x = 0.01 obtain an approximation to $\sqrt{23}$. [4]

[4]

[4]

- $f(x) = (1 3x)^{\frac{3}{2}}$
 - a Use the binomial expansion to expand f(x) in ascending powers of x up to and including x^3 .
 - **b** By substituting a suitable value of x, find an approximation to $97^{\frac{3}{2}}$.
- 10 a Expand $\frac{5}{\sqrt{1-2x}}$ in ascending powers of x up to and including x^3 and simplify each terms fully. [5]
 - **b** State for which values of x the expansion is valid. [1]
- 11 $f(x) = \frac{4}{\sqrt{1+\frac{2}{9}x}}, -\frac{3}{2} < x < \frac{3}{2}$
 - **a** Expand f(x) in ascending powers of x up to and including x^2 and simplify each term fully. [5]
 - **b** Given that $f(\frac{1}{10}) = \sqrt{15}$. Obtain an approximation of $\sqrt{15}$, giving your answer as a fully simplified fraction. [4]
- Expand $\frac{2}{\sqrt{1+2x}}$ in ascending powers of x up to and including x^3 and simplify each term fully. [4]
 - **b** State for which the values of x the expansion is valid. [2]

CHAPTER SUMMARY: CHAPTER 6

The binomial expansion $(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$

can be used to give an exact expression if n is a positive integer. It can also be used to give an approximation for any **rational** number.

The expansion of $(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$

where n is negative or a fraction, is only valid if |x| < 1.

The first four terms of $(1 + 3x)^6$ are

$$1 + 6(3x) + \frac{6(5)}{2!}(3x)^2 + \frac{6(5)(4)}{3!}(3x)^3 + \dots = 1 + 18x + 135x^2 + 540x^3$$

The expansion is finite and exact.

The first four terms of $(1 - 2x)^{\frac{1}{3}}$ are

$$1 + \left(\frac{1}{3}\right)(-2x) + \left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\frac{(-2x)^2}{2} + \left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)\left(-\frac{5}{3}\right)\frac{(-2x)^3}{6} = 1 - \frac{2}{3}x - \frac{4}{9}x^2 - \frac{40}{81}x^3$$

The expansion is infinite and approximate.

CHAPTER 7



7 SCALAR AND VECTOR QUANTITIES

In the United States, there are around 5000 aeroplanes in the sky at any given time. How do the pilots and air traffic control ensure that these aircraft move around the skies without colliding? They use **vectors** to keep the aircraft on specific flight paths. Vectoring is used to aid navigation and to guide aircraft into their final position so that they can land safely.

Without the use of vectors, commercial aviation would hardly have developed at all.



LEARNING OBJECTIVES

- Use vector notation and draw vector diagrams
- Perform simple vector arithmetic and understand the definition of a unit vector
- Use vectors to describe the position of a point in two dimensions
- Write down and use the Cartesian components of a vector in two dimensions
- Use vectors to demonstrate simple properties of geometrical figures

STARTER ACTIVITIES

A quantity that has both **size** (magnitude) and **direction** is a **vector**.

A quantity that just has size is a scalar.

A scalar quantity can be described by using a single number (its magnitude or size).

A vector quantity has both magnitude and direction.

For example

Scalar: The distance from P to Q is 100 metres.

(Distance is a scalar)

Vector: From P to Q you go 100 metres north.

(This is called the **displacement** from *P* to *Q*. Displacement is a vector.)


Scalar: A ship is sailing at 12 km h^{-1} .

(Speed is a scalar)

Vector: A ship is sailing at 12 km h^{-1} , on a **bearing** of 060° .

(This is called the velocity of the ship. Velocity is a vector.)

The diagram below shows the displacement vector from P to Q where Q is 500 m due north of P.



(for example, 1 cm represents 100 m).

▲ a **←**

Sometimes, instead of using the end points P and Q, a small (lower case) letter is used. In print, the small letter will be in **bold type**. In writing you should underline the small letter to show it is a vector: \mathbf{a} or \mathbf{a}

100 m




12 km h⁻¹

VECTOR NOTATION AND HOW TO DRAW VECTOR DIAGRAMS

You need to be able to write down vectors and draw vector diagrams.


Vectors that are equal have both the same magnitude and the same direction.




Here $\overrightarrow{PQ} = \overrightarrow{RS}$

HINT

Think of displacement vectors. If you travel from P to Q, then from Q to R, the resultant journey is P to $\overrightarrow{R}: \overrightarrow{PO} + \overrightarrow{OR} = \overrightarrow{PR}$




Two vectors are added using the 'triangle law'




When you add the vectors \mathbf{a} and \mathbf{b} , the resultant vector $\mathbf{a} + \mathbf{b}$ goes from 'the start of \mathbf{a} to the finish of \mathbf{b} '. This is sometimes called the triangle law for vector addition

EXAMPLE 1

The diagram shows the vectors \mathbf{a} , \mathbf{b} and \mathbf{c} . Draw another diagram to illustrate the vector $\mathbf{a} + \mathbf{b} + \mathbf{c}$





First use the triangle law for $\mathbf{a} + \mathbf{b}$. Then use it again for $(\mathbf{a} + \mathbf{b}) + \mathbf{c}$

The resultant goes from the start of a to the finish of c

HINT

If you travel from P to Q, then back from Q to P, you are back where you started, so your displacement is zero.

Adding the vectors \overrightarrow{PQ} and \overrightarrow{QP} gives the zero vector $\mathbf{0}$: $\overrightarrow{PQ} + \overrightarrow{QP} = \mathbf{0}$

The zero displacement vector is **0**. It is printed in bold type, or underlined in written work.

You can also write $\overrightarrow{PQ} = -\overrightarrow{QP}$. So $\overrightarrow{PQ} + \overrightarrow{QP} = \mathbf{0}$ or $\overrightarrow{PQ} - \overrightarrow{PQ} = \mathbf{0}$


The modulus, or magnitude, of a vector is written in single vertical bars.

The modulus of the vector \mathbf{a} is written as $|\mathbf{a}|$.

The modulus of the vector \overrightarrow{PQ} is written as $|\overrightarrow{PQ}|$.

EXAMPLE 2

The vector \mathbf{a} is directed due east and $|\mathbf{a}| = 12$. The vector \mathbf{b} is directed due south and $|\mathbf{b}| = 5$. Find $|\mathbf{a} + \mathbf{b}|$.



$$|\mathbf{a} + \mathbf{b}|^2 = 12^2 + 5^2 = 169$$

$$|a + b| = 13$$

Use the triangle law for adding the vectors **a** and **b**

Use Pythagoras' theorem

EXAMPLE 3


In the diagram, $\overrightarrow{QP} = \mathbf{a}$, $\overrightarrow{QR} = \mathbf{b}$, $\overrightarrow{QS} = \mathbf{c}$, and $\overrightarrow{RT} = \mathbf{d}$. Find in terms of \mathbf{a} , \mathbf{b} , \mathbf{c} , and \mathbf{d} .

$$\overrightarrow{PS}$$

$$\overrightarrow{RP}$$

$$\stackrel{\frown}{PT}$$

$$\overrightarrow{d}$$
 \overrightarrow{TS}



$$\overrightarrow{PS} = \overrightarrow{PQ} + \overrightarrow{QS} = -\mathbf{a} + \mathbf{c} = \mathbf{c} - \mathbf{a}$$

$$\overrightarrow{RP} = \overrightarrow{RQ} + \overrightarrow{QP} = -\mathbf{b} + \mathbf{a} = \mathbf{a} - \mathbf{b}$$

$$\overrightarrow{PT} = \overrightarrow{PR} + \overrightarrow{RT} = (\mathbf{b} - \mathbf{a}) + \mathbf{d} = \mathbf{b} + \mathbf{d} - \mathbf{a}$$

$$\overrightarrow{TS} = \overrightarrow{TR} + \overrightarrow{RS} = -\mathbf{d} + (\overrightarrow{RQ} + \overrightarrow{QS})$$

So
$$\overrightarrow{TS} = -\mathbf{d} + (-\mathbf{b} + \mathbf{c}) = \mathbf{c} - \mathbf{b} - \mathbf{d}$$

Add vectors using the triangle $\triangle PQS$

Add vectors using $\triangle RQP$

Add vectors using $\triangle PRT$ and use

$$\overrightarrow{PR} = -\overrightarrow{RP} = -(\mathbf{a} - \mathbf{b}) = (\mathbf{b} - \mathbf{a})$$

Add vectors using $\triangle TRS$ and also $\triangle RQS$

EXERCISE 1

1 Given vector **a** (as seen in the diagram), draw the vectors

SKILLS

SOLVING


REASONING

$$c \frac{1}{4}a$$

$$d -\frac{1}{3}a$$

2 The vectors **a** and **b** are shown in the diagram. Draw


$$\mathbf{a} + \mathbf{b}$$



- 3 \triangleright The vector **a** is directed due east and $|\mathbf{a}| = 32$. The vector **b** is directed due north east and $|\mathbf{b}| = 6$. Find $|\mathbf{a} + \mathbf{b}|$.
- The vector **a** is directed south west and $|\mathbf{a}| = 16$. The vector **b** is directed south east and $|\mathbf{b}| = 20$. Find $|\mathbf{a} + \mathbf{b}|$.
- 5 ln the diagram $\overrightarrow{NP} = \mathbf{a}$, $\overrightarrow{PQ} = \mathbf{b}$, $\overrightarrow{QM} = \mathbf{c}$ and $\overrightarrow{NO} = \mathbf{d}$ Find in terms of a, b, c, d





$$\stackrel{\longrightarrow}{MN}$$



- 6 In the diagram $\overrightarrow{PR} = \mathbf{a}$ and $\overrightarrow{PO} = \mathbf{b}$. M is the midpoint of \overrightarrow{PR} . In terms of a and b find
 - a PM

 - c OM
- 7 \rightarrow In the diagram $\overrightarrow{OP} = \mathbf{a}, \overrightarrow{OR} = \mathbf{b}$ and $\overrightarrow{OS} = \mathbf{c}$. Given that $\overrightarrow{PR} = \overrightarrow{RS}$, prove that $-\mathbf{a} + 2\mathbf{b} = \mathbf{c}$.





PERFORM SIMPLE VECTOR ARITHMETIC AND UNDERSTAND THE **DEFINITION OF A UNIT VECTOR**

KEY POINTS


A unit vector is a vector with a magnitude of 1. Two important unit vectors are commonly used: these are the vectors in the direction of the x and y. The unit vector in the direction of the x-axis is i and the unit vector in the direction of the y-axis is j.

EXAMPLE 4

The diagram shows the vector a.

Draw diagrams to illustrate the vectors $3\mathbf{a}$ and $-2\mathbf{a}$.





Any vector **parallel** to the vector **a** may be written as λ **a**, where λ is a non-zero scalar.

EXAMPLE 5

Show that the vectors $6\mathbf{a} + 8\mathbf{b}$ and $9\mathbf{a} + 12\mathbf{b}$ are parallel.

$$9a + 12b$$


$$=\frac{3}{2}(6\mathbf{a}+8\mathbf{b})$$

$$\lambda = \frac{3}{2}$$

So the vectors are parallel.

Subtracting a vector is equivalent to 'adding a negative vector', so $\mathbf{a} - \mathbf{b}$ is defined to be $\mathbf{a} + (-\mathbf{b})$.





HINT

To subtract **b**, you reverse the direction of **b**, then add.

A unit vector is a vector with modulus 1.

EXAMPLE 6

The vector **a** has magnitude 20 units. Write down a unit vector that is parallel to **a**.

The unit vector is $\frac{\mathbf{a}}{20}$ or $\frac{1}{20}\mathbf{a}$

Divide **a** by its magnitude. In general, the unit vector is $\frac{\mathbf{a}}{|\mathbf{a}|}$

If $\lambda \mathbf{a} + \mu \mathbf{b} = \alpha \mathbf{a} + \beta \mathbf{b}$ and the non-zero vectors \mathbf{a} and \mathbf{b} are not parallel, then $\alpha = \lambda$ and $\beta = \mu$.

The above result can be shown as follows:

 $\lambda \mathbf{a} + \mu \mathbf{b} = \alpha \mathbf{a} + \beta \mathbf{b}$ can be written as $(\lambda - \alpha)\mathbf{a} = (\beta - \mu)\mathbf{b}$, but two vectors cannot be equal unless they are parallel or zero.

Since the vectors **a** and **b** are not parallel or zero, $(\lambda - a) = 0$ and $(\beta - \mu) = 0$, so $\alpha = \lambda$ and $\beta = \mu$.

EXAMPLE 7

Given that $5\mathbf{a} - 4\mathbf{b} = (2s + t)\mathbf{a} + (s - t)\mathbf{b}$, where the non-zero vectors \mathbf{a} and \mathbf{b} are not parallel, find the values of the scalars s, t.

$$2s + t = 5$$

$$s - t = -4$$

$$3s = 1$$

$$s = \frac{1}{3}$$

$$t = s + 4 = 4\frac{1}{3}$$

So
$$s = \frac{1}{3}$$
 and $t = 4\frac{1}{3}$

Equate the coefficients of a and b

Solve simultaneously by adding the equations

HINT

It is useful to

check that the result is correct in special cases.

For example, if

k = 1, SX = 4a.This is correct

because if k = 1,

X and R are the same point

and SR = 4a

EXAMPLE 8


In the diagram, $\overrightarrow{PQ} = 3\mathbf{a}$, $\overrightarrow{QR} = \mathbf{b}$, $\overrightarrow{SR} = 4\mathbf{a}$ and $\overrightarrow{PX} = \overrightarrow{kPR}$. Find, in terms of \mathbf{a} , \mathbf{b} and \mathbf{k} :

$$\overrightarrow{PS}$$

$$\overrightarrow{PX}$$

$$\overrightarrow{SQ}$$

$$\overrightarrow{SX}$$



$$\overrightarrow{PS} = \overrightarrow{PR} + \overrightarrow{RS} = \overrightarrow{PQ} + \overrightarrow{QR} + \overrightarrow{RS} = 3\mathbf{a} + \mathbf{b} - 4\mathbf{a} = \mathbf{b} - \mathbf{a}$$

Use the triangle law

$$\overrightarrow{PR} = \overrightarrow{PQ} + \overrightarrow{QR} = 3\mathbf{a} + \mathbf{b}$$

$$\overrightarrow{PX} = \overrightarrow{kPR} = k(3\mathbf{a} + \mathbf{b})$$

$$\overrightarrow{SQ} = \overrightarrow{SR} + \overrightarrow{RQ} = 4\mathbf{a} + (-\mathbf{b}) = 4\mathbf{a} - \mathbf{b}$$

$$\overrightarrow{SX} = \overrightarrow{SP} + \overrightarrow{PX} = -\overrightarrow{PS} + \overrightarrow{PX} = -(\mathbf{b} - \mathbf{a}) + k(3\mathbf{a} + \mathbf{b}) = (3k + 1)\mathbf{a} + (k - 1)\mathbf{b}$$

SKILLS

REASONING CRITICAL THINKING

1 ln the triangle \overrightarrow{ABC} , $\overrightarrow{AB} = 2\mathbf{a}$, $\overrightarrow{BC} = \mathbf{b}$. The midpoint of \overrightarrow{AC} is M.

Find in terms of a and b


$$\overrightarrow{AC}$$

$$\overrightarrow{AM}$$

$$\stackrel{\smile}{c} \stackrel{\longrightarrow}{CM}$$

- 2 The diagram shows a parallelogram ABCD E is a point on BC such that BE:EC = 1:3Write expressions for these vectors.
 - a DC
- **b** CD
- c AC

- \mathbf{d} AE



3 \triangleright In each part find whether the given vector is parallel to $2\mathbf{a} + 3\mathbf{b}$.

c a +
$$\frac{3}{2}$$
b

4 \triangleright The non-zero vectors **a** and **b** are not parallel. In each part find the value of λ and μ :

a
$$\mathbf{a} + 4\mathbf{b} = 2\lambda\mathbf{a} - \mu\mathbf{b}$$

b
$$(\lambda + 2)$$
a + $(\mu - 3)$ **b** = 0

c
$$5\lambda \mathbf{a} - 4\mathbf{b} - 2\mathbf{a} + \mu \mathbf{b} = 0$$

d
$$(1 + \lambda)a + 3\lambda b = \mu a + 4\mu b$$

e
$$(3\lambda + 5)\mathbf{a} + \mathbf{b} = 2\mathbf{a} + (\lambda - 3)\mathbf{b}$$

f
$$6\mathbf{a} - 5\mathbf{b} = (2\lambda + \mu)\mathbf{a} + (\lambda - \mu)\mathbf{b}$$

g
$$10\mathbf{a} - 8\mathbf{b} = (4\lambda + 2\mu)\mathbf{a} + (\lambda - \mu)\mathbf{b}$$

5 \rightarrow In the triangle PQR, $\overrightarrow{PQ} = \mathbf{a}$, $\overrightarrow{PR} = \mathbf{b}$.


M is the midpoint of PQ and N lies on PR such that PN:PR=1:4. Find in terms of a and b.

$$\overrightarrow{PM}$$

$$\stackrel{\longrightarrow}{PN}$$

$$\overrightarrow{RN}$$

$$\overrightarrow{ON}$$



6 \triangleright *ABCDEF* is a regular **hexagon** with centre O.

$$\overrightarrow{OA} = \mathbf{a} \text{ and } \overrightarrow{AB} = \mathbf{b}$$

Find expressions, in terms of ${\bf a}$ and ${\bf b}$, for


- OB
- \overrightarrow{AC}
- c \overrightarrow{EC}



Vector
$$\overrightarrow{PQ} = 2\mathbf{a}$$
 and vector $\overrightarrow{PS} = 3\mathbf{b}$.

$$\overrightarrow{PR} = 3\overrightarrow{PQ}$$
 and $\overrightarrow{PU} = 2\overrightarrow{PS}$


- a Write the vector \overrightarrow{PV} in terms of **a** and **b**.
- **b** Write the vector \overrightarrow{RU} in terms of **a** and **b**.



2a

HINT


The position vector of P is how you get from O to P



USE VECTORS TO DESCRIBE THE POSITION OF A POINT IN TWO DIMENSIONS

The position vector of a point A is the vector \overrightarrow{OA} , where O is the origin.

 \overrightarrow{OA} is usually written as vector \mathbf{a}




 $\overrightarrow{OA} = \mathbf{a}$

HINT


Use the triangle law to give $\overrightarrow{AB} = \overrightarrow{AO} + \overrightarrow{OB}$ = -a + b So $\overrightarrow{AB} = b - a$

 $\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$, where \mathbf{a} and \mathbf{b} are the position vectors of A and B respectively.



EXAMPLE 9

In the diagram the points A and B have position vectors a and b respectively (referred to the origin O). The point P divides AB in the ratio 1:2.



Find the position vector of P.

$$\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$$

$$\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$$

OP is the position vector of P

$$\overrightarrow{AP} = \frac{1}{3}(\mathbf{b} - \mathbf{a})$$

Use the 1:2 ratio (AP) is one third of AB)


$$\overrightarrow{OP} = \mathbf{a} + \frac{1}{3}(\mathbf{b} - \mathbf{a}) = \frac{2}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}$$

SKILLS ANALYSIS

- ▶ The points A and B have position vectors **a** and **b** respectively (referred to the origin O). The point P divides AB in the ratio 1:5. Find, in terms of **a** and **b**, the position vector of *P*.
- 2 > The points A, B and C have position vectors a, b and c respectively (referred to the origin O). The point P is the midpoint of AB. Find, in terms of **a**, **b** and **c**, the vector *PC*.
- **3** \triangleright OABCDE is a regular hexagon. The points A and B have position vectors **a** and **b** respectively, referred to the origin O. Find, in terms of **a** and **b**, the position vectors of C, D and E.

USE VECTORS TO DEMONSTRATE SIMPLE PROPERTIES OF GEOMETRICAL FIGURES

EXAMPLE 10



ABC is a triangle AB is 2a AC is 3b

Find BC in terms of a and b

$$BC = BA + AC$$
 such that $BD:DC = 2:3$

$$BC = -2a + 3b$$

D is a point on the line BC such that BD:DC = 2:3

b Show that AD is parallel to the vector $\mathbf{a} + \mathbf{b}$

First you need to find AD in terms of **a** and **b**.

$$AD = AB + BD$$

$$AD = AB + \frac{2}{5}BC$$

$$AD = 2\mathbf{a} + \frac{2}{5}(-2\mathbf{a} + 3\mathbf{b})$$

$$AD = 2\mathbf{a} + -\frac{4}{5}\mathbf{a} + \frac{6}{5}\mathbf{b}$$

By expanding the brackets


$$AD = \frac{6}{5}\mathbf{a} + \frac{6}{5}\mathbf{b}$$

Factorise to find if parallel.

$$AD = \frac{6}{5}(\mathbf{a} + \mathbf{b})$$

Therefore line AD is parallel to vector $\mathbf{a} + \mathbf{b}$ because AD is a multiple of vector $\mathbf{a} + \mathbf{b}$

EXAMPLE 11



In the diagram $AB = \mathbf{a}$ is and $AC = \mathbf{b}$

The point *X* divides *AB* in the ratio
$$AX:AB=4:1$$

The point Y divides BC in the ratio

$$BY:YC = 2:3$$

The point D on AC is such that

$$AC:CD = 5:3$$

a Show that
$$\overrightarrow{XY} = \frac{2}{5}\mathbf{b} - \frac{1}{5}\mathbf{a}$$

$$\overrightarrow{XY} = XB + BY$$

$$\overrightarrow{XB} = \frac{1}{5}\mathbf{a}$$

$$\overrightarrow{BC} = -\mathbf{a} + \mathbf{b}$$

Therefore:

$$\overrightarrow{XY} = \frac{1}{5}\mathbf{a} + \frac{2}{5}(-\mathbf{a} + \mathbf{b})$$

$$= \frac{1}{5}\mathbf{a} - \frac{2}{5}\mathbf{a} + \frac{2}{5}\mathbf{b}$$

$$= -\frac{1}{5}\mathbf{a} + \frac{2}{5}\mathbf{b}$$

$$= \frac{2}{5}\mathbf{b} - \frac{1}{5}\mathbf{a}$$

Due to the ratio 2:3

HINT

Co-linear means all the points are on the same line. **b** Show that *X*, *Y* and *D* are co-linear

If two points are parallel and come from the same point then they are co-linear.

$$XD = XA + AD$$

$$XD = XA + AC + CD$$

$$XD = -\frac{4}{5}\mathbf{a} + (\mathbf{b} + \frac{3}{5}\mathbf{b})$$

$$XD = -\frac{4}{5}\mathbf{a} + 1\frac{3}{5}\mathbf{b}$$

$$XD = -\frac{4}{5}\mathbf{a} + \frac{8}{5}\mathbf{b}$$


$$XY = -\frac{1}{5}\mathbf{a} + \frac{2}{5}\mathbf{b}$$

Therefore XD is 4(XY) which shows XD is parallel to XY

Since XD and XY share a point and are parallel then they must be on the same straight line and co-linear

EXERCISE 4

1




In triangle OAB the midpoint on AB is M.

$$\overrightarrow{OA} = \mathbf{x} \quad \overrightarrow{OB} = \mathbf{y} \quad \overrightarrow{OD} = 2\mathbf{x} \text{ and } \overrightarrow{OC} = 2\mathbf{y}$$

- **a** Express \overrightarrow{AB} , \overrightarrow{OM} and \overrightarrow{DC} in terms of **x** and **y**
- **b** Show that *AB* and *DC* are parallel.

2




N is the midpoint of \overrightarrow{OB}

M is the midpoint of \overrightarrow{OA}

$$\overrightarrow{OA} = \mathbf{a} \quad \overrightarrow{OB} = \mathbf{b}$$

Show MN is parallel to AB


3



The ratio of OA:OC = 1:2 and OB:OD = 1:2

- a Find AB, OC, OD and DC
- **b** Show that *AB* is parallel to *DC*

4




OM:MQ = 1:2

$$PR = \frac{1}{3}PQ$$

Show that OA and MN are parallel

5




The diagram shows OABCMN.

$$\overrightarrow{OA} = \mathbf{a}, \overrightarrow{OB} = \mathbf{b}, \overrightarrow{BC} = 3\mathbf{b}$$

$$AM:MB = 2:1$$

$$AN:NC = 1:2$$

- Find \overrightarrow{AB} in terms of **a** and **b**
- **b** Find *OM* in terms of **a** and **b**
- Show that OMN are co-linear



- Show that $AD = 3\mathbf{b} \mathbf{a}$
- Show that A, X, D are co-linear


OACB is a parallelogram

OA is a

OB is **b**

OB to OD is in the ratio 1:3

$$XC$$
 is $\frac{1}{3}BC$



$$OP = \mathbf{a} \quad OR = \mathbf{b}$$

N is the point such that $\overrightarrow{PQ} = \overrightarrow{QN}$

The point M divides PR in the ratio 2:1

- Show that $\overrightarrow{OM} = \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$
- **b** Prove that *OMN* is a straight line


WRITE DOWN AND USE CARTESIAN COMPONENTS OF A VECTOR IN TWO **DIMENSIONS**

The vectors i and j are unit vectors parallel to the x-axis and the y-axis, and in the direction of x increasing and y increasing, respectively.

EXAMPLE 12

The points A and B in the diagrams have coordinates (3, 4) and (11, 2) respectively. Find, in terms of i and j:

- a The position vector of A.
- The position vector of B.
- The vector AB.



HINT

This standard notation is easy to read and also avoids the need to write out lengthy expressions with i and j terms.

$$\mathbf{a} \quad \mathbf{a} = \overrightarrow{OA} = 3\mathbf{i} + 4\mathbf{j}$$

b
$$\mathbf{b} = \overrightarrow{OB} = 11\mathbf{i} + 2\mathbf{j}$$

$$\overrightarrow{AB} = \mathbf{b} - \mathbf{a} = (11\mathbf{i} + 2\mathbf{j}) - (3\mathbf{i} + 4\mathbf{j})$$

= $8\mathbf{i} - 2\mathbf{j}$

(i goes one unit 'across', j goes one unit 'up

You can see from the diagram that the vector ABgoes 8 units 'across' and 2 units 'down'

You can write a vector with Cartesian components $x\mathbf{i} + y\mathbf{j}$ as a **column matrix**: $\begin{pmatrix} x \\ y \end{pmatrix}$

EXAMPLE 13

Given that $\mathbf{a} = 2\mathbf{i} + 5\mathbf{j}$, $\mathbf{b} = 12\mathbf{i} - 10\mathbf{j}$ and $\mathbf{c} = -3\mathbf{i} + 9\mathbf{j}$, find $\mathbf{a} + \mathbf{b} + \mathbf{c}$ using column matrix notation in your working.

$$\mathbf{a} + \mathbf{b} + \mathbf{c} = \begin{pmatrix} 2 \\ 5 \end{pmatrix} + \begin{pmatrix} 12 \\ -10 \end{pmatrix} + \begin{pmatrix} -3 \\ 9 \end{pmatrix} = \begin{pmatrix} 11 \\ 4 \end{pmatrix}$$

Add the numbers in the top line to get 11, the x component, and the bottom line to get 4, the y component. This is 11i + 4j.

The modulus or magnitude of $x\mathbf{i} + y\mathbf{j}$ is $\sqrt{x^2 + y^2}$.

EXAMPLE 14

The vector \mathbf{a} is equal to $5\mathbf{i} - 12\mathbf{j}$. Find $|\mathbf{a}|$, and find a unit vector in the same direction as \mathbf{a} .

$$|\mathbf{a}| = \sqrt{5^2 + (-12)^2} = \sqrt{169} = 13$$


A unit vector in the same direction of **a** is $\frac{\mathbf{a}}{|\mathbf{a}|}$

$$\frac{\mathbf{a}}{|\mathbf{a}|} = \frac{5\mathbf{i} - 12\mathbf{j}}{13} = \frac{5}{13}\mathbf{i} - \frac{12}{13}\mathbf{j}$$

You can also express this as $\frac{1}{13}\begin{pmatrix} 5\\ -12 \end{pmatrix}$.

HINT

From Pythagoras' theorem, the magnitude of xi + yj, represented by the hypotenuse, is $\sqrt{x^2 + y^2}$



EXAMPLE 15

Given that $\mathbf{a} = 5\mathbf{i} + \mathbf{j}$ and $\mathbf{b} = -2\mathbf{i} - 4\mathbf{j}$, find the exact value of $|2\mathbf{a} + \mathbf{b}|$.

$$2\mathbf{a} + \mathbf{b} = 2 {5 \choose 1} + {-2 \choose -4} = {10 \choose 2} + {-2 \choose -4} = {8 \choose -2}$$

$$|2\mathbf{a} + \mathbf{b}| = \sqrt{8^2 + (-2)^2} = \sqrt{68} = \sqrt{4}\sqrt{17} = 2\sqrt{17}$$

EXERCISE 5

▶ Given that $\mathbf{a} = 3\mathbf{i} + 4\mathbf{j}$ and $\mathbf{b} = -2\mathbf{i} + 2\mathbf{j}$, find

SKILLS

INTERPRETATION

- **b** −3**b** a 2a
- **2** Given that $\mathbf{a} = 4\mathbf{i} + 4\mathbf{j}$, $\mathbf{b} = 2\mathbf{i} + 6\mathbf{j}$ and $\mathbf{c} = 3\mathbf{i} + 4\mathbf{j}$, find
 - a + b + c
- **b** |a + b|

c 2b - c

d |2b - c|

- e 2b + 2c 3a
- 3 \blacktriangleright The points A, B, and C have coordinates (4, -2), (6, 4) and (-2, 6), find
 - a the position vectors of A, B and C

AB

d 4b - 5a

- c AC
- d | OC |

c a + b

- \triangleright P, Q, and R are points with position vectors $2\mathbf{i} 3\mathbf{j}$, $\mathbf{i} + 2\mathbf{j}$ and $4\mathbf{i} 2\mathbf{j}$. Find in terms of \mathbf{i} and \mathbf{j}
 - a \overrightarrow{PQ}

- 5 \blacktriangleright If the coordinates of A are (3, 4) and $AB = 2\mathbf{i} + 2\mathbf{j}$, find the position vector of B.
- 6 \triangleright Given that $\mathbf{a} = 6\mathbf{i} + 3\mathbf{j}$, $\mathbf{b} = 6\mathbf{i} 8\mathbf{j}$ and $\mathbf{c} = -5\mathbf{i} + 14\mathbf{j}$, find the unit vector in the direction of a, b and c.
- **7** Figure 3.2 Given that $2\mathbf{e} 3\mathbf{f} = \begin{pmatrix} 5 \\ 15 \end{pmatrix}$, where $\mathbf{e} = \begin{pmatrix} 4 \\ m \end{pmatrix}$ and $\mathbf{f} = \begin{pmatrix} n \\ -3 \end{pmatrix}$, find the constants m and n.
- 8 If $\mathbf{r} = \begin{pmatrix} 4 \\ -1 \end{pmatrix}$, $\mathbf{s} = \begin{pmatrix} 3 \\ 7 \end{pmatrix}$ and $p\mathbf{r} + q\mathbf{s} = \begin{pmatrix} 7 \\ 37 \end{pmatrix}$, find the constants p and q.
- 9 Given that $\mathbf{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 2 \\ -6 \end{pmatrix}$, find
 - $\mathbf{a} + 3\mathbf{b}$
- **b** |**a** + 3**b**| **c** 4**a** + 2**b**
- **d** |4a + 2b|

EXAM PRACTICE: CHAPTER 7

Relative to a fixed origin O, the point A has position vector $5\mathbf{i} - 6\mathbf{j}$.


The point B is such that $AB = \mathbf{i} + 11\mathbf{j}$.

Show that the triangle *OAB* is isosceles.

[4]

In the diagram below, $BA = \mathbf{a}$ and $BC = \mathbf{c}$. DE is such that DA:DE = 1:5 and AX is such that AX:XC = 3:2.

Prove BX and XE are co-linear



[5]

If the vectors $4\mathbf{i} - 6\mathbf{j}$ and $m\mathbf{i} - 2\mathbf{j}$ are parallel, state the value of m.

[2]

The vectors of A and B are $3\mathbf{i} - 2\mathbf{j}$ and $t\mathbf{i} + \mathbf{j}$. Find the value of t if OAB is a straight line.

[3]

The points *A*, *B*, *C* and *D* form a quadrilateral and:

$$AB = 3i + 5j$$
, $AC = 6i + 6j$ and $BC = 3i + j$

[2]

b Hence, or otherwise, show that *ABCD* is a trapezium.

[3] [2]

f c Find the exact value of |BD|.

d Find a unit vector parallel to *BD*.

[3]

• The point F is on the line BD and BF:FD = 1:2

$$AF = 5\mathbf{i} + 4\frac{1}{3}\mathbf{j}$$

a Find AD.

Point E is on the line AD and ABCE is a parallelogram.

Show that *F* lies on the line *CE*.

[3]

f Find the ratio of *EF*: *FC*.

[4]

Zaynab, Asaad and Ali enter a running competition. They all take different routes, which are described by these vectors, where $\mathbf{s} = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$, $\mathbf{t} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$ and the units are km.

Zaynab: $\mathbf{s} + 2\mathbf{t}$

Asaad: 2s + t

Ali: 5**s** − **t**

a Express each journey as a column vector.

[3]

b They all take 6 hours to complete their routes.i Find the length of each journey in km.

[2]

ii Find the average speed of each runner.

[1]



 \bigcirc OPQR is a parallelogram. M is the midpoint of the diagonal OQ.

$$\overrightarrow{OP} = 2\mathbf{a}$$
 and $\overrightarrow{OR} = 2\mathbf{b}$


a Express \overrightarrow{OM} in terms of **a** and **b**.

[2]

b Use vectors to show that *M* is the midpoint of *PR*.

[2]

R The diagram below shows a regular hexagon.



$$\overrightarrow{OA} = 4\mathbf{a}$$
 $\overrightarrow{OB} = 4\mathbf{b}$

Express in terms of a and/or b

$$\overrightarrow{AB}$$

$$\stackrel{\longrightarrow}{EF}$$

X is the midpoint of BC.

c Express \overrightarrow{EX} in terms of **a** and/or **b**.


Y is the point on AB extended such that AB:BY = 3:2

- **d** Prove that E, X and Y lie on the same straight line.
- Show that the triangle whose vertices have position vectors 2i + 6j, 8i + 2j and 11i + 13j is an isosceles triangle.
- 10 $\mathbf{a} = \begin{pmatrix} 3 \\ -6 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 4 \\ -2 \end{pmatrix}$

Find **b**
$$-2a$$
. [2]

- OAB is a triangle. M is the midpoint of OA. N is the midpoint of OB. $\overrightarrow{OM} = \mathbf{a}$ and $\overrightarrow{ON} = \mathbf{b}$. Show that AB is parallel to MN.
- The diagram shows a parallelogram ABCD with $\overrightarrow{AB} = \mathbf{b}$ and $\overrightarrow{AD} = \mathbf{a}$. E is such that that $\overrightarrow{DE}:\overrightarrow{EC}=3:1$.

$$\overrightarrow{EF} = \lambda \overrightarrow{AE}$$
 and $\overrightarrow{CF} = \mu \overrightarrow{BC}$



a Find \overrightarrow{CF} in terms of μ .

[2]

b Find EF in terms of **a**, **b** and λ .

[2]

[5]

[3]

c Find the values of μ and λ .


[4]

CHAPTER SUMMARY: CHAPTER 7

The velocity of a skier moving down a ski slope has direction and magnitude.




- Vectors that are equal have both the same magnitude and the same direction.
- Two vectors can be added using the 'triangle law'.



- The modulus of the vector is another way of saying its magnitude. The words are interchangeable.
 - The modulus of vector \mathbf{a} is written as $|\mathbf{a}|$.
 - The modulus of vector \overrightarrow{PQ} is written as $|\overrightarrow{PQ}|$.
 - The modulus of $x\mathbf{i} + y\mathbf{j}$ is $\sqrt{x^2 + y^2}$.
- The vector $-\mathbf{a}$ has the same magnitude as the vector \mathbf{a} but in the opposite direction.
- If vector **a** is parallel to vector **b**, $\mathbf{a} = \lambda \mathbf{b}$ where λ is a scalar.
- $\mathbf{a} \mathbf{b}$ is the same as $\mathbf{a} + (-\mathbf{b})$.
- A unit vector is a vector that has a modulus (or magnitude) of 1 unit.
- If $\lambda \mathbf{a} + \mu \mathbf{b} = \alpha \mathbf{a} + \beta \mathbf{b}$ and the non-zero vectors \mathbf{a} and \mathbf{b} are not parallel, then $\lambda = \alpha$ and $\mu = \beta$.
- The position vector of point A is the vector OA, where O is the origin.
- OA is usually written as vector **a**.
- $\overrightarrow{AB} = \mathbf{b} \mathbf{a}$, where \mathbf{a} and \mathbf{b} are the position vectors of A and B respectively.

CHAPTER 8



8 RECTANGULAR CARTESIAN COORDINATES

Straight line graphs are an excellent tool for showing linear growth or a linear relationship between two variables. Scientists and engineers use them on a daily basis.

One example of this is the linear relationship between the flow of water and water level. In May 2006, the Sichuan earthquake in China caused landslides that blocked several rivers, creating natural dams. Lakes formed behind these dams and the largest of these was Lake Tangjiashan. This filled with water at a rate that was about five times the rate at which water could drain from it. As a result, the water level grew to the point where it threatened to flood an area that is home to over one million people.



Scientists used the methods you will cover in this chapter to predict when the lake would overflow. This gave time for towns and villages to be evacuated, saving many lives.

LEARNING OBJECTIVES

- Use the equation of the straight line y = mx + c and ax + by + c = 0
- Work out the gradient of a straight line
- Find the equation of a straight line

- Understand the relationships between the gradients of parallel and perpendicular lines
- Find the distance between two points on a straight line
- Find the coordinates of a point that divides a straight line in a given ratio

STARTER ACTIVITIES

Write out the gradient and *y*-intercept of these lines.

a
$$y = 2x + 5$$

b
$$10x - 5y = 15$$
 c $y = \frac{1}{2}x$

$$y = \frac{1}{3}x$$

$$\mathbf{d} \quad y = \mathbf{6}$$


$$y = \frac{2}{3}x + 9$$

WORK OUT THE GRADIENT OF A STRAIGHT LINE

KEY POINTS

You can work out the gradient (m) of a straight line joining the point with coordinates (x_1, y_1) to the point with coordinates (x_2, y_2) by using the formula

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$



Work out the grad

Work out the gradient of the line joining these pairs of points.

(-2, 7) and (4, 5)

- **b** (2d, -5d) and (6d, 3d)
- **a** $(x_1, y_1) = (-2, 7)$ and $(x_2, y_2) = (4, 5)$

So,
$$m = \frac{5-7}{4-(-2)}$$

= $\frac{-2}{6} = -\frac{1}{3}$

Use $m = \frac{y_2 - y_1}{x_2 - x_1}$

The gradient of the line is $-\frac{1}{3}$

b
$$(x_1, y_1) = (2d, -5d)$$
 and $(x_2, y_2) = (6d, 3d)$
 $3d - (-5d)$

$$m = \frac{3d - (-5d)}{6d - 2d}$$
$$= \frac{8d}{4d} = 2$$

Use
$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

The gradient of the line is 2.

EXAMPLE 2

The line joining (2, -5) to (4, a) has gradient -1. Work out the value of a.

$$m = -1$$
, $(x_1, y_1) = (2, -5)$ and $(x_2, y_2) = (4, a)$

$$m = \frac{a - (-5)}{4 - 2} = -1$$

Use
$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

So
$$\frac{a+5}{2} = -1$$
,

$$a + 5 = -2$$

and a = -

EXERCISE 1

- 1 Work out the gradient of the lines joining these points.
 - a (4, 3) and (6, 4)


b (-3, 2) and (-4, -2)

 $\left(\frac{1}{3}, \frac{2}{3}\right)$ and $\left(\frac{4}{5}, \frac{1}{2}\right)$

- d (2c, 4c) and (5c, 6c)
- **2** The line joining (2, -4) to (5, d) has a gradient of 3. Work out the value of d.
- **3** \blacktriangleright The line joining (3, -6) to (6, α) has a gradient of 3. Find α .
- **4** The line joining (a, 3) to (2, 1) has a gradient of $-\frac{1}{3}$. Find a.
- 5 The line joining (-10a, 6) to (17, -4) has a gradient of $-\frac{1}{5}$. Find a.
- **6** The line joining (3a, a) to (-27, -7) has a gradient of $\frac{23}{25}$. Find a.
- **7** \blacktriangleright The quadrilateral A, B, C, D has coordinates (-6, 1), (-4, 4), (2, 0) and (0, -3).
 - a Find the gradients of the lines AB, BC, CD and DA.
 - b What do these gradients tell us about the quadrilateral?

FIND THE EQUATION OF A STRAIGHT LINE

You can find the equation of a straight line with gradient m that passes through the point with coordinates (x_1, y_1) by using the formula $y - y_1 = m(x - x_1)$



EXAMPLE 3

Find the equation of the line with gradient $-\frac{1}{2}$ that passes through the point (4, -6).

$$m=-\frac{1}{2}, (x_1, y_1)=(4, -6)$$

$$y - (-6) = -\frac{1}{2}(x - 4)$$

$$y + 6 = -\frac{1}{2}(x - 4)$$

$$y + 6 = -\frac{1}{2}x + 2$$

$$y=-\frac{1}{2}x-4$$

EXAMPLE 4

Find the equation of the line with gradient 5 that passes through the point (3, 2).

The gradient is 5, so

$$\frac{y-2}{x-3}=5$$

$$y - 2 = 5(x - 3)$$


$$y - 2 = 5x - 15$$

$$y = 5x - 13$$

(x, y) is any point on the line.

Use
$$y - y_1 = m(x - x_1)$$

Use $y - y_1 = m(x - x_1)$



EXAMPLE 5

A line l is parallel to the line y=2x+3 and passes through the point (1, 2). Find the equation of the line l.

Parallel lines have the same gradient so line l has gradient 2.

$$2 = 2 \times 1 + c$$

To find c, substitute m = 2, x = 1 and y = 2 into y = mx + c

$$c = 0$$

So line l has equation

$$y = 2x$$

KEY POINTS

Parallel lines have the same gradient


EXERCISE 2

SKILLS THINKING

- Find the equation of the line with gradient m that passes through the point (x_1, y_1) when
- **a** m=3 and $(x_1, y_1)=(2, 6)$ **b** m=-6 and $(x_1, y_1)=(3, 6)$ **c** $m=\frac{1}{2}$ and $(x_1, y_1)=(-6, -10)$
- Find the equation of the line that passes through the points (4, 13) and (-15, 6). Write your answer in the form ax + by + c = 0
- Find the equation of the line that passes through the points (22, 9) and (6, 8). Write your answer in the form y = mx + c
- Find the equation of the line that passes through the points (18, 6) and (4, 12). Write your answer in the form y = mx + c
- **5** A line is parallel to the line $y = -\frac{2}{5}x 3$ and intercepts the *y*-axis at (0, -4). Calculate the equation of the line and write your answer in the form ax + by + c = 0.
- 6 A line is parallel to the line 2x 3y + 7 = 0 and passes through the point (0, 3). Write down the equation of the line.
- 7 The line y = 6x 18 meets the x-axis at the point P. Calculate the coordinates of P.
- 8 The line $y = \frac{2}{3}x + 7$ meets the y-axis at point the B. The point C has coordinates (6, 3). Find the equation of the line joining B and C.

UNDERSTAND THE RELATIONSHIP BETWEEN PERPENDICULAR LINES





If a line has a gradient of m, a line perpendicular to it has a gradient of $-\frac{1}{m}$.

You can also say that if two lines are perpendicular, the product of their gradients is -1.

EXAMPLE 6

Work out the gradient of the lines that are perpendicular to the lines with these gradients.

b
$$\frac{1}{2}$$

$$-\frac{2}{5}$$

Gradient m=3 so the gradient of the perpendicular line is $-\frac{1}{3}$. Gradient $=-\frac{1}{m}$ Gradient $m=\frac{1}{2}$ so the gradient of the perpendicular line is -2. $-\frac{1}{2}\times -2=-1$

Gradient =
$$-\frac{1}{m}$$

b Gradient $m = \frac{1}{2}$ so the gradient of the perpendicular line is -2.

$$-\frac{1}{2}\times -2=-1$$

c Gradient $m = -\frac{2}{5}$. so the gradient of the perpendicular line is $\frac{5}{2}$. $-\frac{1}{2} \times -2 = -1$

$$-\frac{1}{2}\times -2 = -\frac{1}{2}$$

129

Show that the line y = 3x + 4 is perpendicular to the line x + 3y - 3 = 0.

The gradient of the line y = 3x + 4 is 3.

Compare
$$y = 3x + 4$$
 with $y = mx + c$ to find m

Rearrange into the form y = mx + c

$$x + 3y - 3 = 0$$

$$3y = -x + 3$$

$$y = -\frac{1}{2}x + 1$$

The gradient of this line is $-\frac{1}{3}$.

The products of the two gradients is $3 \times -\frac{1}{3} = -1$.

The lines are perpendicular because the product of their gradients is -1.

EXAMPLE 8

Work out whether these pairs of lines are parallel, perpendicular, or neither.

a
$$y = -2x + 9$$
 and $y = -2x - 3$

b
$$3x - y - 2 = 0$$
 and $x + 3y - 6 = 0$

$$y = \frac{1}{2}x$$
 and $2x - y + 4 = 0$

a
$$y = -2x + 9$$
 has gradient -2

This is of the form y = mx + c, so m = -2

$$y = -2x - 3$$
 has gradient -2

Since the gradients are equal the two lines are parallel.

b
$$3x - y - 2 = 0$$

so,
$$y = 3x - 2$$

Rearrange the equation

Therefore the gradient of this line is 3.

$$x + 3y - 6 = 0$$

So,
$$y = -\frac{1}{3}x + 2$$

Therefore the gradient of this line is $-\frac{1}{3}$

Since $3 \times \frac{1}{3} = -1$ the lines are perpendicular.

$$y = \frac{1}{2}x$$
 is already in the form $y = mx + c$

So, the gradient is $\frac{1}{2}$.

$$2x - y + 4 = 0$$

So,
$$y = 2x + 4$$

Therefore the gradient of this line is 2.

These two lines are neither parallel nor perpendicular.

EXAMPLE 9

Find an equation of the line that passes through the point (3, -1) and is perpendicular to the line y = 2x - 4

The line y = 2x - 4 has gradient m = 2.

So the gradient of the perpendicular line is $-\frac{1}{2}$.

$$m = -\frac{1}{2}$$
, $(x_1, y_1) = (3, -1)$

$$y - (-1) = -\frac{1}{2}(x - 3)$$

$$y + 1 = -\frac{1}{2}x + \frac{3}{2}$$

$$y=-\frac{1}{2}x+\frac{1}{2}$$

Use the rule $-\frac{1}{m}$ with m=2

Use $y - y_1 = m(x - x_1)$

SKILLS REASONING Work out if these pairs of lines are parallel, perpendicular or neither.

a
$$y = 4x + 2$$

b
$$y = \frac{3}{4}x + 3$$

$$y=-\frac{1}{4}x-7$$

$$y = \frac{3}{4}x - 5$$

$$y = 5x - 3$$

d
$$5x - y - 1 = 0$$

$$5x - y = -4$$

$$y=-\frac{1}{5}x+2$$

e
$$y = -\frac{3}{2}x + 6$$

 $3x + 2y - 3 = 0$

f
$$5x - y + 5 = 0$$

$$2x + 0y - 6 = 0$$



▶ Find the equation of the line that passes through (-3, 8) and is perpendicular to the line y = 5x + 2.

3 Find the equation of the line

- a parallel to the line y = -2x + 3 and passing through (1, 1)
- **b** perpendicular to the line y = 4x 4 and passing through (-2, 6)
- perpendicular to the line 2x + y 9 = 0 and passing through (3, 2)
- **d** perpendicular to the line 4x 6y = -7 and passing through (-3, 4).

 \triangleright Find the equation of a line passing through the point (3, -4) that is perpendicular to the line 8x + 6y = 15. Write your answer in the form ax + by + c = 0.

5 Find the equation of a line passing through the point (6, 3) and perpendicular to the line 4x - 5y = -10. Write your answer in the form ax + by + c = 0.

6 Find the equation of a line that passes through (5, 6) and is perpendicular to the line whose equation is $y = \frac{2}{3}x + 5$


Find the equation of a line that passes through (7, -3) and is perpendicular to 4x + 3y = 12.

The vertices of a quadrilateral PQRS have coordinates P(-1, 5), Q(7, 1), R(5, -3), S(-3, 1). Show that the quadrilateral is a rectangle.

FIND THE DISTANCE BETWEEN TWO POINTS ON A LINE

You can find the distance d between two points (x_1, y_1) and (x_2, y_2) that lie on a straight line using the formula:


$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$



EXAMPLE 10

Find the distance between these pairs of points.

- a (2, 3) and (5, 7)
- **b** (4a, a) and (-3a, 2a)
- c $(2\sqrt{2}, -5\sqrt{2})$ and $(4\sqrt{2}, \sqrt{2})$



Draw a sketch

Let
$$(x_1, y_1) = (2, 3)$$
 and $(x_2, y_2) = (5, 7)$

$$d = \sqrt{(5-2)^2 + (7-3)^2}$$

$$= \sqrt{3^2 + 4^2}$$

$$= \sqrt{25} = 5$$

Use $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Take the positive square root because d is a length

b Let
$$(x_1, y_1) = (4a, a)$$
 and $(x_2, y_2) = (-3a, 2a)$

$$d = \sqrt{(-3a - 4a)^2 + (2a - a)^2}$$

$$= \sqrt{(-7a)^2 + a^2}$$

$$= \sqrt{49a^2 + a^2}$$

$$= \sqrt{50a^2}$$

So
$$d = 5\sqrt{2}a$$

c
$$(x_1, y_1) = (2\sqrt{2}, -5\sqrt{2})$$
 and $(x_2, y_2) = (4\sqrt{2}, \sqrt{2})$

$$d = \sqrt{(4\sqrt{2} - 2\sqrt{2})^2 + (\sqrt{2} - (-5\sqrt{2}))^2}$$

$$= \sqrt{(2\sqrt{2})^2 + (6\sqrt{2})^2}$$

$$= \sqrt{8 + 72}$$

$$= \sqrt{80}$$
So $d = 4\sqrt{5}$

EXERCISE 4


SKILLS REASONING

- Find the distance between these points, leaving your answer in surd form where appropriate.
 - a (1, 2) and (7, 10)
- **b** (3, -7) and (8, 5)
- (-2, 2) and (1, 6)
- d (5, 2) and (1, 6)
- (-1, -5) and (5, 5)
- f (-4a, 0) and (3a, -2a)
- (-7b, 5b) and (2b, -5b)
- h (-2c, 2c) and (-3c, 5c)



FIND THE COORDINATES OF A POINT THAT DIVIDES A LINE IN A GIVEN RATIO

Consider a line between points $A(x_1, y_1)$ and $B(x_2, y_2)$, divided by the point $P(x_p, y_p)$ in a ratio of m:n.



HINT

If m = n then the coordinate P is the midpoint of Aand B

Using similar triangles, you can derive these formulae:

$$x_p = \frac{nx_1 + mx_2}{m+n} \text{ and } y_p = \frac{ny_1 + my_2}{m+n}$$

EXAMPLE 11

Find the coordinates of the point which divides A(0, 6): B(4, 10) in the ratio 3:1.

$$x_p = \frac{(1)(0) + (3)(4)}{3+1} = 3$$

$$(1)(6) + (3)(10)$$

Use
$$x_p = \frac{nx_1 + mx_2}{m + n}$$

$$y_p = \frac{(1)(6) + (3)(10)}{3+1} = 9$$

Use
$$y_p = \frac{ny_1 + my_2}{m+n}$$

Therefore the coordinates are (3, 9)

EXAMPLE 12

A line between points A(-1, 3) and B(5, 7) is divided by point P such that AP:PB = 1:4. Find the coordinates of point *P*.

$$x_p = \frac{4(-1) + (1)(5)}{1 + 4}$$
$$= \frac{1}{5}$$

Use
$$x_p = \frac{nx_1 + mx_2}{m + n}$$

$$y_p = \frac{4(3) + 1(7)}{1 + 4}$$
$$= \frac{19}{5}$$

Use
$$y_p = \frac{ny_1 + my_2}{m+n}$$

Therefore the coordinates of *P* are $\left(\frac{1}{5}, \frac{19}{5}\right)$

EXERCISE 5

- 1 Find the coordinates of the points which divide AB in the given ratios.
- **SKILLS** INTERPRETATION
- **a** A(1, 5):B(-2, 8) in the ratio 1:2.
- **b** A(3, -7): B(-2, 8) in the ratio 3:2.
- A(-2, 5): B(5, 2) in the ratio 4:3.
- **d** A(4, 2):B(6, 8) in the ratio 2:2.
- ≥ Find the coordinates of the point which divides the line segment from (-2, 1) to (2, 3) in the ratio 3:4.
- 3 Find the coordinates of the point that divides the line segment MN with the coordinates of endpoints at M(-4, 0) and M(0, 4) in the ratio 3:1.

[2]

[1]

- A is a point with coordinates (2, 4) and B is a point with coordinates (-1, 0). The line L has the equation 3y = 4 - 2x. Is the line L parallel to AB? Show your working clearly.
 - The straight line L_1 passes through the points (-3, 0) and (9, 10). Find an equation for L_1 in the form ax + by + c = 0, where a, b and c are integers.
 - The equation of a line is given as 4x 5y 8 = 0. Point A(7, 4) and B(2, 0) lie on the line.
 - a Find the length of AB, leaving your answer in surd form. [3]

The point C has coordinates (2, t), where t > 0, and AC = AB.

- **b** Find the value of t.
- Find the area of triangle ABC.
- Find the equation of the line L_1 which is perpendicular to the line y=2x+8 at the point (1, 10). Give your answer in the form ax+by+c=0. [3]
- Find the equation of the line which is perpendicular to the line y = 5x + 8 at the point (5, 33). [2]
- Find the equation of the line which is perpendicular to the line $y=-\frac{1}{3}x+8$ at the point (9, 5). [3]
- A is the point (0, 1) and B is the point (10, 6).
 - a Find the coordinates of the midpoint of AB.

The equation of the straight line through A and B is $y = \frac{1}{2}x + 1$.

- b Write down the equation of another straight line that is parallel to AB and passes through the point (0, 1).
- Prove that the coordinates (3, 2), (1, 2), (-3, -2), (1, 2) form a square. [2]
- Find the area of rectangle ABCD with vertices A(-3, 0), B(3, 2), C(4, -1), and D(-2, -3). [2]
- Quadrilateral ABCD has vertices A(-3, 0), B(2, 4), C(3, 1), and D(-4, -3).

 Calculate the perimeter of the quadrilateral.

b Calculate the area of *DEFG*.


| 11 | The line l_1 passes through the points $A(-1, 2)$ and $B(11, 8)$. | |
|----|---|-----|
| | a Find an equation for l_1 . | [2] |
| | The line l_2 passes through the point $C(10, 0)$ and is perpendicular to l_1 . The lines l_1 and l_2 intersect at the point D . | |
| | b Calculate the coordinates of <i>D</i> . | [2] |
| | c Calculate the length AD . | [2] |
| | f d Hence, or otherwise, calculate the area of the triangle ACD . | [1] |
| 12 | Given the points $A(-1, 2)$ and $B(7, 8)$, determine the coordinates of point P that divides AB in the ratio 1:3. | [3] |
| 13 | Determine the midpoint between $(-7, -2)$ and $(4, 5)$. | [2] |
| 14 | The points $A(-5, 5)$ and $B(9, -2)$ lie on the line L . | |
| | a Find an equation for L in the form $ax + by + c = 0$. | [2] |
| | b Show that the distance between A and B is $7\sqrt{5}$. | [2] |
| 15 | Quadrilateral $DEFG$ has vertices $D(5, 1)$, $E(2, 4)$, $F(-4, 4)$ and $G(-1, 1)$. a Find the equation of the line of DE . | [4] |
| | | |

CHAPTER SUMMARY: CHAPTER 8


The general form of the equation of the straight line is y = mx + c where m is the gradient and c is the y-intercept or constant.

Another form of this equation is ax + by + c = 0 where a, b, c are integers.

- The formula for calculating the gradient is $m = \frac{y_2 y_1}{x_2 x_1}$.
- You can find the equation of the line with gradient m that passes through the point with coordinates (x_1, y_1) by using $y y_1 = m(x x_1)$.
- Two lines with the same gradient are parallel.




- If a line has a gradient of m, a line perpendicular to it has a gradient of $-\frac{1}{m}$.
- You can also say that if two lines are perpendicular, the product of their gradients is -1.



- You can find the distance d between (x_1, y_1) and (x_2, y_2) using the formula: $d = \sqrt{(x_2 x_1)^2 + (y_2 y_1)^2}$
- The coordinates of the point (x_p, y_p) dividing the line joining the points (x_1, y_1) and (x_2, y_2) in the ratio m:n are given by $x_p = \frac{nx_1 + mx_2}{m+n}$ and $y_p = \frac{ny_1 + my_2}{m+n}$.

CHAPTER 9



9 DIFFERENTIATION

In the time of Sir Isaac Newton and Gottfried Leibniz, the founders of **calculus**, one of the biggest problems was poor navigation at sea. Shipwrecks occurred because the captain did not know the actual position of his ship. This was mainly due to a poor understanding of how the earth, planets and stars moved in relation to each other. Calculus (differentiation and integration) was developed to improve this understanding and help improve navigation at sea.

Differentiation is used to solve many real life problems. For example, it is used to determine maximum and minimum values, which you will learn about in this chapter. This technique is very useful for companies to determine costs, and for engineers and architects to calculate the amount of building material required.



LEARNING OBJECTIVES

- Find the gradient function of a curve and differentiate a function that has multiple powers of \boldsymbol{x}
- Differentiate e^{ax} , $\sin ax$ and $\cos ax$
- Use the chain rule to differentiate more complicated functions
- Use the product rule to differentiate more complicated functions

- Use the quotient rule to differentiate more complicated functions
- Find the equation of the **tangent** and **normal** to the curve y = f(x)
- Find the stationary points of a curve and calculate if they are minimum or maximum points
- Apply what you have learned about turning points to solve problems

STARTER ACTIVITIES

Simplify

a
$$a^6 \times a^7$$

$$(a^2)^{-1}$$

$$(a^3)^{-\frac{1}{2}}$$

$$\mathbf{m} \left(\frac{a}{b}\right)^2$$

$$\left(\frac{a^2}{b^3}\right)^4$$

$$\left(\frac{a^8b^2}{c^6}\right)^{-\frac{1}{2}}$$

b
$$\frac{a^7}{a^{-3}}$$

$$(a^2)^{-3}$$

$$(a^6)^{\frac{1}{3}}$$

$$(a^2 \times b^{-4})^3$$

$$(m^{-1}n^3)^{-2}$$

$$\sqrt{\frac{m^2}{x}}$$

$$\frac{a^{-5}}{a^9}$$

g
$$(a^{-2})^{-4}$$

$$(\alpha^9)^{-\frac{1}{3}}$$

$$(a^3b^{\frac{1}{2}})^4$$

$$\left(\frac{a^6}{h_{10}}\right)^{\frac{1}{2}}$$

$$\mathbf{w} \left(\frac{x^2y}{z^3}\right)^{-1}$$

d
$$a^{-4} \times a^{-2}$$

$$(a^{-12})^{-\frac{1}{4}}$$

$$(a^2b^{-2})^{-2}$$

$$\left(\frac{a^2}{m^4}\right)^{-1}$$


$$\mathbf{x} \quad \left[(a^3b^{-8})^{-\frac{1}{3}} \right]^2$$


FIND THE GRADIENT FUNCTION OF A CURVE AND DIFFERENTIATE A FUNCTION THAT HAS MULTIPLE POWERS OF x

On a straight line graph, the gradient is constant, the same everywhere along the line: it has a fixed value.

However, on a curved graph the gradient is always changing. The gradient depends on where you are along the x-axis. The different gradients can be shown by tangents, lines that touch the curve in one place only.

Differentiation is the process by which you produce an equation to calculate the gradient at a specific point.





As a general rule, if;

$$f(x) = ax^n$$

then...

$$f'(x) = n\alpha x^{n-1}.$$

This is the formula for the curve. i.e. the function

This is the gradient function for f(x), the formula that gives the gradient at any point.

It is also known as the derivative, or 'derived functon'

KEY POINTS

A function can be written as f(x) or as an equation in the form $y = ax^2 + bx + c$

The notation for the gradient function is: f'(x) or $\frac{dy}{dx}$

EXAMPLE 1

Find the **derived function** when f(x) equals

b
$$x^{\frac{1}{2}}$$

d
$$\frac{x}{x^5}$$

d
$$\frac{x}{r^5}$$
 e $x^2 \times x^3$

 $a 6x^5$

The power 6 is reduced to the power 5 and the 6 multiplies the answer

b
$$f(x) = x^{\frac{1}{2}}$$

The power
$$\frac{1}{2}$$
 is reduced to $\frac{1}{2} - 1 = -\frac{1}{2}$, and the $\frac{1}{2}$ multiplies the answer

$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}}$$

This is then written in an alternative form

c
$$f(x) = x^{-2}$$

$$f'(x) = -2x^{-3}$$

The power -2 is reduced to -3 and the -2 multplies the answer

$$=-\frac{2}{x^3}$$

This is also rewritten in an alternative form using knowledge of negative powers

d Let
$$f(x) = \frac{x}{x^5}$$

= x^{-4}

Simplify using rules of powers to give one simple power, i.e. subtract 1 - 5 = -4

So
$$f'(x) = -4x^{-5}$$

= $-\frac{4}{x^5}$

Reduce the power -4 to give -5, then multiply your answer by -4

e Let
$$f(x) = x^2 \times x^3$$

= x^5

Add the powers this time to give 2 + 3 = 5

So
$$f'(x) = 5x^4$$

Reduce the power 5 to 4 and multiply your answer by 5

EXAMPLE 2

Use standard results to differentiate

$$x^3 + x^2 - x^{\frac{1}{2}}$$

b
$$2x^{-3}$$

$$\frac{1}{3}x^{\frac{1}{2}} + 4x^2$$

$$y = x^3 + x^2 - x^{\frac{1}{2}}$$

So
$$\frac{dy}{dx} = 3x^2 + 2x - \frac{1}{2}x^{-\frac{1}{2}}$$

= $3x^2 + 2x - \frac{1}{2\sqrt{x}}$

Differentiate each term as you come to it. First x^3 , then x^2 , then $x^{\frac{1}{2}}$

b
$$y = 2x^{-3}$$

So
$$\frac{dy}{dx} = -6x^{-4}$$
$$= -\frac{6}{x^4}$$

Differentiate x^{-3} , then multiply the answer by 2

$$y = \frac{1}{3}x^{\frac{1}{2}} + 4x^2$$

So
$$\frac{dy}{dx} = \frac{1}{3} \times \frac{1}{2}x^{-\frac{1}{2}} + 8x$$

= $\frac{1}{6} \times x^{-\frac{1}{2}} + 8x$
= $\frac{1}{6\sqrt{x}} + 8x$

Take each term as you come to it, and treat each term as a multiple

EXAMPLE 3

Let
$$f(x) = 4x^2 - 8x + 3$$

- a Find the gradient of y = f(x) at the point $(\frac{1}{2}, 0)$.
- **b** Find the coordinates of the point on the graph of y = f(x) where the gradient is 8.
- **c** Find the gradient of y = f(x) at the points where the curve meets the line y = 4x 5.

a As
$$f(x) = 4x^2 - 8x + 3$$

$$f'(x) = 8x - 8 + 0$$

First find f'(x), the derived function

So
$$f'\left(\frac{1}{2}\right) = -4$$

Then substitute the x-coordinate value to obtain the gradient

b
$$f'(x) = 8x - 8 = 8$$

Put the gradient function equal to 8

so
$$x = 2$$

granical survey of an area

so
$$x = 2$$

Then solve the equation you have obtained to give the value of xSubstitute this value for x into f(x) to give the value of y

The point where the gradient is 8 is (2, 3).

and y = f(2) = 3

8 8 is (2, 3). Interpret your answer in words

$$4x^2 - 8x + 3 = 4x - 5$$

$$4x^2 - 12x + 8 = 0$$

$$x^2 - 3x + 2 = 0$$

$$(x-2)(x-1)=0$$

So
$$x = 1$$
 or $x = 2$

At x = 1 the gradient is 0.

At x = 2 the gradient is 8, as in part b.

Put f(x) = 4x - 5, then rearrange and collect terms to give a quadratic equation

Divide by the common factor 4

Solve the quadratic equation by factorising

Substitute the values of x into f'(x) = 8x - 8to give the gradients at the specified points

EXAMPLE 4

Use standard results to differentiate.

a
$$\frac{1}{4\sqrt{x}}$$

b
$$x^3(3x+1)$$

$$\frac{x-2}{r^2}$$

a Let
$$y = \frac{1}{4\sqrt{x}}$$

= $\frac{1}{4}x^{-\frac{1}{2}}$

Therefore
$$\frac{dy}{dx} = -\frac{1}{8}x^{-\frac{3}{2}}$$

Express the 4 in the denominator as a multiplier of
$$\frac{1}{4}$$
 and express the x term as power $-\frac{1}{2}$

efore
$$\frac{dy}{dx} = -\frac{1}{8}x^{-\frac{3}{2}}$$
 Differentiate by reducing the power of x and multiplying $\frac{1}{4}$ by $-\frac{1}{2}$

b Let
$$y = x^3(3x + 1)$$

= $-\frac{1}{8}x^{\frac{3}{2}}$
= $3x^4 + x^3$

Therefore
$$\frac{dy}{dx} = 12x^3 + 3x^2$$
$$= 3x^2(4x + 1)$$

c Let
$$y = \frac{x-2}{x^2}$$

 $= \frac{1}{x} - \frac{2}{x^2}$
 $= x^{-1} - 2x^{-2}$
Therefore $\frac{dy}{dx} = -x^{-2} + 4x^{-3}$

Express the single fraction as two separate fractions, and simplify
$$\frac{x}{x^2}$$
 to $\frac{1}{x}$

Then express the rational expressions as negative powers of
$$x$$
, and differentiate

Find f'(x), given that f(x) equals

SKILLS

- **c** $x^{\frac{1}{2}}$ **d** $x^{\frac{3}{2}}$

- REASONING
- \sqrt{x} g $\sqrt[3]{x}$ h x^{-4} i $\frac{1}{x^2}$ l $x^3 \times x^6$ m $x^{-4} \times x^{\frac{1}{2}}$

 $=\frac{-(x-4)}{r^3}$

2 Differentiate

$$a 3x^2$$

b
$$7x^4$$

$$x^{-2}$$

$$d = \frac{3}{4}$$

$$e^{5x^{5}}$$

f
$$3\sqrt{x}$$

g
$$\frac{3}{\sqrt{x}}$$

h
$$\frac{5}{\sqrt[3]{x}}$$

3 Differentiate

$$y = x^5 + x^{-2}$$

b
$$y = \frac{1}{2}x^{-2} + 4x^{-2}$$

b
$$y = \frac{1}{2}x^{-2} + 4x^2$$
 c $y = x^{\frac{1}{8}} + 3x^{-2} + 9x^{\frac{1}{8}}$

4 Find
$$\frac{dy}{dx}$$
 when

$$y = x^6 + x^2 + 2x$$

b
$$y = x^3 + 3$$

$$y = 6x^3 - 5x^2 + 3x^{-2} + 6x + 3$$

5 Find f'(x) when

a
$$f(x) = 3x + \frac{2}{3}x^{\frac{1}{3}}$$

$$f(x) = 3x^{\frac{1}{6}} + 2x^{\frac{3}{4}}$$

b
$$f(x) = 3x^{\frac{1}{6}} + 2x^{\frac{3}{4}}$$
 c $f(x) = 3 - 8x^{-1} + x^{-\frac{2}{3}}$

6 Find the gradient of the curve at point *A* for

a
$$f(x) = x^4 + 3x^2 + 6x + 3$$
 and $A = (0, 0)$

b
$$f(x) = 3x^2 + 2x^{-1}$$
 and $A = (2, 13)$

c
$$f(x) = 4x^2 + 3x^{-1} + x^{\frac{1}{2}} + 157$$
 and $A = (4, 13)$

7 Differentiate

$$\frac{1}{3}x^3(x-2)$$

b
$$3\sqrt{x} + \frac{1}{2x}$$

$$\frac{2x+3}{x}$$

$$\frac{3x^2-6}{x}$$

$$x(x^2 - x + 2)$$

b
$$3\sqrt{x} + \frac{1}{2x}$$
 c $\frac{2x+3}{x}$ **e** $x(x^2 - x + 2)$ **f** $(3x - 2)(4x + \frac{1}{x})$

USE THE CHAIN RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS

Consider the function $y = (1 + x)^{10}$.

One way of calculating $\frac{dy}{dx}$ is to use the binomial expansion, but this is a very long and inefficient way of calculating it.

Another way is to use the chain rule.

EXAMPLE 5

Differentiate $y = (3x^4 + x)^5$

$$Let u = 3x^4 + x$$

$$\frac{\mathrm{d}u}{\mathrm{d}x} = 12x^3 + 1$$

$$y = u^{5}$$

$$\frac{\mathrm{d}y}{\mathrm{d}u} = 5u^4$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$
$$= 5u^4 \times 12x^3 + 1$$

 $= 5(3x^4 + x)^4(12x^3 + 1)$

Using the chain rule $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

Substitute $u = 3x^4 + x$ back into the equation

Differentiate: $y = \sqrt{5x^2 + 1}$

Let
$$u = 5x^2 + 1$$

Using the chain rule $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

$$\frac{\mathrm{d}u}{\mathrm{d}x} = 10x$$

$$y = \sqrt{u}$$

$$y = u^{\frac{1}{2}}$$

Using the rules of indices

$$\frac{\mathrm{d}y}{\mathrm{d}u} = \frac{1}{2}u^{-\frac{1}{2}}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

$$=\frac{1}{2}u^{-\frac{1}{2}}\times 10x$$

$$=5xu^{-\frac{1}{2}}$$

$$=5x(5x^2+1)^{-\frac{1}{2}}$$

$$=\frac{5x}{\sqrt{5x^2+1}}$$

 $u = 5x^2 + 1$ back into the equation.

EXAMPLE 7

Given that $y = (x^2 - 7x)^4$

Calculate $\frac{dy}{dx}$ using the chain rule.

$$y=(x^2-7x)^4$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

$$let u = x^2 - 7x \qquad \frac{du}{dx} = 2x - 7$$

$$y = u^4 \qquad \qquad \frac{\mathrm{d}y}{\mathrm{d}u} = 4 \, u^3$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4u^3(2x - 7)$$

$$\frac{dy}{dx} = 4(x^2 - 7x)^3(2x - 7)$$

Differentiate \boldsymbol{u} in terms of \boldsymbol{x}

Differentiate \boldsymbol{y} in terms of \boldsymbol{u}

Substitute expressions for $\frac{dy}{du}$ and $\frac{du}{dx}$ into $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

Substitute in $u = x^2 - 7x$

EXAMPLE 8

Differentiate

$$y = \sin(2x^2 + 3)$$

$$y = \sin(2x^2 + 3)$$

Let
$$u = 2x^2 + 3$$
 so $\frac{du}{dx} = 4x$

$$y = \sin u$$
 so $\frac{dy}{du} = \cos u$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \cos u \times 4x$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4x\cos(2x^2 + 3)$$

Substitute $u = 2x^2 + 3$ back into the equation

EXERCISE 2

SKILLS

CRITICAL **PROBLEM** SOLVING

Differentiate

$$(3x + 1)^4$$

b
$$(2x-4)^6$$

$$c (1 + r^2)^3$$

$$(1-5x)^7$$

c
$$(1 + x^2)^3$$
 d $(1 - 5x)^7$ **e** $(2 + 3x)^{-1}$

$$(6x + 2)^{\frac{1}{2}}$$

f
$$(6x+2)^{\frac{1}{2}}$$
 g $\left(\frac{2x^4+6}{2}\right)^7$ **h** $\sqrt{7-x}$ **i** $4(2-7x)^4$ **j** $6(7-3x)^{-3}$

h
$$\sqrt{7-x}$$

$$4(2-7x)^4$$

$$\mathbf{j}$$
 6(7 - 3 \mathbf{x})

Differentiate

a
$$6e^{2x}$$

d
$$e^{x^2-3}$$

b
$$10e^{4-x}$$
 c $3e^{-5x}$ **d** e^{x^2-3x} **e** $5e^{x^2+4x}$

3 Differentiate

$$a \sin(2x + 1)$$

b
$$\cos(2x^2 + 2)$$
 c $\cos^3 x$

$$c \cos^3 x$$

d
$$2\sin^2 2x$$

5 Find an equation for the tangent to each curve at the point on the curve with the given x-coordinate.

$$u = (4x - 6)^4$$
 at $x = 2$

b
$$y = \sin(2x + 6)$$
 at $x = -3$

DIFFERENTIATE e^{ax} , $\sin ax$ AND $\cos ax$

In Chapter 10 you will learn about radians. Radians are just an alternative way of measuring angles.

The conversion between radians and degrees is:

π radians = 180 degrees

These rules apply when radians are used.

KEY POINTS

If
$$y = \sin ax$$
 then $\frac{dy}{dx} = a \cos ax$

If
$$y = a \cos x$$
 then $\frac{dy}{dx} = -a \sin ax$


You will also need to differentiate the exponential function you met in Chapter 1.

This function, e^x , is very special, as the gradient of the curve is the same as the value of the function itself.

KEY POINTS

In other words, if
$$y = e^x$$
 then $\frac{dy}{dx} = e^x$

It follows that if
$$y = e^{ax}$$
 then $\frac{dy}{dx} = e^{ax}$



Differentiate

$$y = \sin 4x$$

$$y = 2\sin 5x$$

$$y = 4\cos x$$

$$\mathbf{d} \quad y = \mathrm{e}^{\frac{1}{2}x}$$

$$y = 3e^{2x}$$

$$y = \sin 4x$$

Using the chain rule =
$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{dy}{dx}$$

Let
$$u = 4x$$

$$\frac{\mathrm{d}u}{\mathrm{d}x} = 4$$

$$y = \sin u$$

$$\frac{\mathrm{d}y}{\mathrm{d}u} = \cos u$$

Therefore

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \cos u \times 4$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4\cos u$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4\cos 4x$$

as
$$u = 4x$$

$$\mathbf{b} \quad y = 2\sin 5x$$

Using the chain rule =
$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{dy}{dx}$$

Let
$$u = 5x$$

$$\frac{\mathrm{d}u}{\mathrm{d}x} = 5$$

$$y = 2 \sin u$$

$$\frac{\mathrm{d}y}{\mathrm{d}u} = 2\cos u$$

Therefore

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 2\cos u \times 5$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 10\cos u$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 10\cos 5x$$

as
$$u = 5x$$

$$y = 4\cos x$$

Using the chain rule =
$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

Let
$$u = x$$

$$\frac{\mathrm{d}u}{\mathrm{d}x} = 1$$

$$y = 4\cos u$$

$$\frac{\mathrm{d}y}{\mathrm{d}u} = -4\sin u$$

Therefore

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -4\sin u \times 1$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -4\sin u$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -4\sin x$$

as
$$u = x$$

$$\mathbf{d} \quad y = \mathrm{e}^{\frac{1}{2}x}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2}\mathrm{e}^{\frac{1}{2}x}$$

If
$$y = e^{ax}$$
 then $\frac{dy}{dx} = ae^{ax}$ with $a = \frac{1}{2}$

$$y = 3e^{2x}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 6\mathrm{e}^{2x}$$

If
$$y = ke^{ax}$$
 then $\frac{dy}{dx} = kae^{ax}$ with $a = 2$ and $k = 3$

EXERCISE 3

SKILLS ANALYSIS REASONING

Differentiate

a $\sin 3x$

- $-\cos x$
- c $-3\sin 2x$

 $d e^{2x}$

- e $3\sin x 2\cos 3x$ f $2e^{2x} + 4x$

2 Find $\frac{dy}{dx}$ when

 $y = \sin 6x$

- **b** $y = 3\cos\frac{1}{2}x$
- $y = \frac{1}{2}\sin 8x + 3\cos 8x + 54$
- **d** $y = 3e^{4x}$

$$y = -\frac{1}{2}e^{3x} + 3e^{5x} - 48x$$

- Find the gradient of the curve with the equation $y = 3e^{-x}$ at the point (0, 3).
- Find the gradient of the curve with the equation $y = 3 \sin x$ at the point $x = \frac{\pi}{3}$.
- Find the gradient of the curve with the equation $y = 5\cos 2x$ at the point $x = \frac{\pi}{4}$.

USE THE PRODUCT RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS

You need to be able to differentiate functions that are multiplied together, using the product rule.

If
$$y = uv$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = u\frac{\mathrm{d}v}{\mathrm{d}x} + v\frac{\mathrm{d}u}{\mathrm{d}x}$$

EXAMPLE 10

Given that $f(x) = x(2x + 1)^3$ find f'(x)

$$f(x) = x(2x + 1)^3$$

$$let u = x so \frac{du}{dx} = 1$$

let
$$v = (2x + 1)^3$$
 so $\frac{dv}{dx} = 3(2x + 1)^2(2)$
= $6(2x + 1)^2$

$$\frac{dy}{dx} = (x)(6(2x+1)^2) + (2x+1)^3(1)$$

$$= 6x(2x+1)^2 + (2x+1)^3$$

$$= (2x+1)^2(6x+(2x+1))$$

$$= (2x+1)^2(8x+1)$$

Apply the product rule

Use the chain rule

$$Use \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

 $(2x + 1)^2$ is a common factor to both terms, so factorise

You can also now simplify the part in brackets

EXAMPLE 11

Given that $f(x) = x^2 \sqrt{3x - 1}$, find f'(x)

$$f(x) = x^2 \sqrt{3x - 1}$$

$$let u = x^2 \qquad so \quad \frac{du}{dx} = 2x$$

let
$$v = (3x - 1)^{\frac{1}{2}}$$
 so $\frac{dv}{dx} = \frac{1}{2}(3x - 1)^{-\frac{1}{2}}(3)$
= $\frac{3}{2}(3x - 1)^{-\frac{1}{2}}$

Apply the product rule

$$\frac{dy}{dx} = (x^2)(\frac{3}{2}(3x-1)^{-\frac{1}{2}}) + (3x-1)^{\frac{1}{2}}(2x)$$

$$= \frac{3x^2}{2}(3x-1)^{-\frac{1}{2}} + 2x(3x-1)^{\frac{1}{2}}$$
Use $\frac{dy}{dx} = u\frac{dv}{dx} + v\frac{du}{dx}$

EXAMPLE 12

Differentiate

$$y = e^x \sin x$$

$$\mathbf{b} \quad y = x \mathbf{e}^{x^2}$$

 $y = e^x \sin x$

Let
$$u = e^x$$
 so $\frac{du}{dx} = e^x$

 $=\frac{3x^2}{2\sqrt{3x-1}}+2x\sqrt{3x-1}$

Let
$$v = \sin x$$
 so $\frac{dv}{dx} = \cos x$

Apply the product rule

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{e}^x(\cos x) + \sin x(\mathrm{e}^x)$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{e}^x(\cos x + \sin x)$$

 $\mathsf{Use}\,\frac{\mathsf{d} y}{\mathsf{d} x} = u\frac{\mathsf{d} v}{\mathsf{d} x} + v\frac{\mathsf{d} u}{\mathsf{d} x}$

HINT

You will also need to use the chain rule to differentiate e^{x²} $y = xe^{x^2}$

Let
$$u = x$$
 so $\frac{du}{dx} = 1$

Let
$$v = e^{x^2}$$
 so $\frac{dv}{dx} = 2xe^{x^2}$

$$\frac{dy}{dx} = x(2xe^{x^2}) + e^{x^2}(1)$$

$$= 2x^2e^{x^2} + e^{x^2}$$

$$= e^{x^2}(2x^2 + 1)$$

Apply the product rule

Use
$$\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

EXERCISE 4

SKILLS

CRITICAL THINKING PROBLEM SOLVING 1 Use the product rule to differentiate

$$x(1 + 4x)^4$$

$$2x(1+3x^2)^2$$

$$x\sqrt{x-2}$$

d
$$2x^4(5+x)^3$$

$$4x^2(5x-2)^{-1}$$

$$x^3\sqrt{3x+1}$$

2 b Use the product rule to differentiate

$$a xe^{3x}$$

b
$$(x^2 - 4)e^{-2x}$$

c
$$2xe^{-3x}$$

d
$$(4x)e^{x^3}$$

$$e^{-(6x+1)e^{x^2}}$$

$$f (5x + 2)e^{-x^2}$$

3 > Use the product rule to differentiate

 $\mathbf{a} \quad x \cos x$

- b $\cos^2 x \sin x$
- c $4x \sin 6x$


 $\mathbf{d} = \mathbf{e}^{x}(\cos x)$

 $e^{x}(\sin x)$

4 Find the value of $\frac{dy}{dx}$ at the on the curve

- a $x^2(4x-3)^3$ where x=2
- **b** $4x(2x-2)^{\frac{1}{2}}$ where x=3
- $(2x + 1)(x + 3)^2$ where x = 4.

Find the equation of the tangent to the curve at the point given by the *x*-coordinate $f(x) = x^2(4x - 1)^{\frac{1}{2}}$ where x = 1.



KEY POINTS

You need to be able to differentiate functions using the quotient rule:

If $y = \frac{u}{v}$ where u and v are functions of x then $\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$

EXAMPLE 13

Given that
$$y = \frac{x^2}{2x+5}$$
 find $\frac{dy}{dx}$

$$y=\frac{x^2}{2x+5}$$

Let
$$u = x^2$$
 so $\frac{du}{dx} = 2x$

Let
$$v = 2x + 5$$
 so $\frac{dv}{dx} = 2$

$$\frac{dy}{dx} = \frac{(2x + 5)(2x) - (x^2)(2)}{(2x + 5)^2}$$
$$= \frac{4x^2 + 10x - 2x^2}{(2x + 5)^2}$$

$$=\frac{2x(x+5)}{(2x+5)^2}$$

Apply the quotient rule

Differentiate

Differentiate

Substitute each component into $\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$

Simplify/work out parts of the numerator

Group together like terms

EXAMPLE 14

Differentiate

a
$$y = \frac{e^{2x+3}}{x}$$

$$\mathbf{b} \quad y = \frac{\mathbf{e}^x}{\sin x}$$

$$y = \frac{e^{2x+3}}{x}$$

Let
$$u = e^{2x+3}$$
 so $\frac{du}{dx} = 2e^{2x+3}$

Let
$$v = x$$
 so $\frac{dv}{dx} = 1$

Therefore

$$\frac{dy}{dx} = \frac{x(2e^{2x+3}) - e^{2x+3}(1)}{x^2}$$
$$= \frac{2xe^{2x+3} - e^{2x+3}}{x^2}$$

Apply the quotient rule

Substitute in each component into $\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$

Simplify

$$=\frac{e^{2x+3}(2x-1)}{x^2}$$

$$y = \frac{e^x}{\sin x}$$

Apply the quotient rule

Let
$$u = e^x$$
 so $\frac{du}{dx} = e^x$

Let
$$v = \sin x$$
 so $\frac{dv}{dx} = \cos x$

$$\frac{dy}{dx} = \frac{\sin x (e^x) - e^x (\cos x)}{(\sin x)^2}$$
$$= \frac{e^x (\sin x - \cos x)}{\sin^2 x}$$

Substitute in each component into
$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{v\frac{\mathrm{d}u}{\mathrm{d}x} - u\frac{\mathrm{d}v}{\mathrm{d}x}}{v^2}$$

Simplify

EXERCISE 5

SKILLS CRITICAL

THINKING **PROBLEM** SOLVING

Use the quotient rule to differentiate

$$\frac{6x}{x-1}$$

b
$$\frac{3x}{6x+3}$$

$$\frac{3x^2}{(2x-3)^3}$$

$$\frac{6x^2}{(x+2)^2}$$

$$\frac{\sqrt{x+3}}{4x^2}$$

$$\frac{x+6}{\sqrt{4x+3}}$$

Use the quotient rule to differentiate

$$\frac{x}{e^{3x}}$$

$$b \frac{e^{x^2}}{x+1}$$

$$\frac{e^{x^2}}{x}$$

$$\frac{e^{3x}}{1-e^{3x}}$$

3 > Use the quotient rule to differentiate

$$\frac{\cos x}{x}$$

$$\frac{e^x}{\sin x}$$

c
$$\frac{3x^2+1}{\cos x}$$
 d $\frac{2x-1}{\sin x}$

d
$$\frac{2x-1}{\sin x}$$

4 Find the equation of the tangent to each curve at the point given by the x-coordinate.


a
$$y = \frac{3x^2 + 1}{(x + 1)^2}$$
 at $x = 3$

b
$$y = \frac{\sin x}{x}$$
 at $x = \pi$

FIND THE EQUATION OF THE TANGENT AND NORMAL TO THE CURVE y = f(x)

A tangent to a curve is a straight line that touches the curve at one point only. As a result the gradient of the tangent is the same as the gradient of the curve at the point where it touches the curve.


The 'normal' is a straight line perpendicular to the tangent at the point where it touches the curve.



You can use differentiation to find the tangent to a curve at a particular point.

EXAMPLE 15

Find the equation of the tangent to the curve $y = x^3 - 3x^2 + 2x - 1$ at the point (3, 5).



$$y = x^3 - 3x^2 + 2x - 1$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 3x^2 - 6x + 2$$

Differentiate to get the gradient function

$$f(3) = 3(3)^2 - 6(3) + 2$$

Substitute the x value in to find the gradient

$$f(3) = 11$$

The gradient at (3, 5) is 11

$$y - y_1 = m(x - x_1)$$

$$y - 5 = 11(x - 3)$$

$$y - 5 = 11x - 33$$

$$y = 11x - 28$$

Use the equation for a straight line

Substitute in the values of the coordinates and the gradient

EXAMPLE 16

Find the equation of the normal to the curve with the equation $y = 3\cos 2x$ at the point $\left(\frac{\pi}{4}, 0\right)$

$$y = 3\cos 2x$$

$$f'(x) = 3(-\sin 2x) \times 2$$

$$= -6 \sin 2x$$

$$f'\left(\frac{\pi}{4}\right) = -6\sin\frac{2\pi}{4}$$
$$= -6$$

Substitute the *x* value in to find the gradient

The gradient of the tangent at $(\frac{\pi}{4}, 0)$ is -6.

So the gradient of the normal is $\frac{1}{6}$.

Product of the gradients of perpendicular lines is -1

 $y - y_1 = m(x - x_1)$

Use the formula for a straight line

 $y-0=\frac{1}{6}\left(x-\frac{\pi}{4}\right)$

Substitute values of the coordinates and the gradient in

 $y = \frac{1}{6}x - \frac{\pi}{24}$

This is the equation of the normal at $(\frac{\pi}{4}, 0)$

Find the equations of the tangent and normal of these equations at the given coordinates.

SKILLS

INTERPRETATION

a
$$y = x^2 - 8x + 11$$
 at (3, -4)

b
$$y = \frac{1}{2}x^2 - x^{-3} + 4x + 2$$
 at $\left(1, \frac{11}{2}\right)$

$$y = \sqrt{x} - \frac{3}{x^2}$$
 at (1, -2)

d
$$y = e^{2x}$$
 at (1, e^2)

$$e \quad y = e^{-3x} \text{ at } \left(1, \frac{1}{e^2}\right)$$

FIND THE STATIONARY POINTS OF A CURVE AND CALCULATE WHETHER THEY ARE MINIMUM OR MAXIMUM STATIONARY POINTS


A point where the gradient of f(x) stops increasing and starts decreasing is called a **maximum point**.

A point where the gradient of f(x) stops decreasing and starts increasing is called a **minimum point**.

These are known as **turning points**, and occur where f'(x) = 0.

To find the coordinates of these points, you need to

- Differentiate f(x) to find the gradient function.
- Solve f'(x) = 0 (as this represents the gradient being 0)
- Substitute the value(s) of x into the original equation to find the corresponding y-coordinate



You can also find out whether stationary points are maximum or minimum

points by finding the value of f''(x) (i.e. $\frac{d^2y}{dx^2}$) at the stationary point.

This is because f''(x) measures the change in gradient.

You obtain f''(x) by differentiating f(x) twice.

If
$$\frac{dy}{dx} = 0$$
 and $\frac{d^2y}{dx^2} < 0$, the point is a maximum.

If
$$\frac{dy}{dx} = 0$$
 and $\frac{d^2y}{dx^2} > 0$, the point is a minimum.

EXAMPLE 17

Find the coordinates of the turning point on the curve $y = x^4 - 32x$, and state whether it is a maximum or minimum.

$$y = x^{4} - 32x$$

$$\frac{dy}{dx} = 4x^{3} - 32$$

$$4x^{3} - 32 = 0$$

$$4x^{3} = 32$$

$$x = 2$$

$$y = x^{4} - 32x$$

$$y = (2)^{4} - 32(2)$$

$$y = -48$$

Differentiate to find the derivative function

Set the derivative function equal to 0

Simplify

To find the value of y at x = 2 substitute x = 2 into the original equation

The stationary point is at (2, -48)

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4x^3 - 32$$

$$\frac{d^2y}{dx^2} = 12x^2 = 12(2)^2 = 48$$

Differentiate again

Substitute in x = 2

So the stationary point is a MINIMUM.

Positive number = minimum point and negative number = maximum point

EXAMPLE 18

Find the stationary points on the curve

$$f(x) = 2x^3 - 15x^2 + 24x + 6$$

and state whether they are minimum or maximum points.

$$f(x) = 2x^3 - 15x^2 + 24x + 6$$

$$f'(x) = 6x^2 - 30x + 24$$

Differentiate to find the derivative function

$$6x^2 - 30x + 24 = 0$$

Set the derivative function equal to 0

$$6(x^2 - 5x + 4) = 0$$

Factorise

$$6(x^2-4)(x-1)=0$$

Factorise again

$$x = 1 \text{ or } x = 4$$

Write the solutions

Substituting into the original formula will give these coordinates as stationary points.

$$(1, 17)$$
 and $(4, -10)$

$$f'(x) = 6x^2 - 30x + 24$$

$$f''(x) = 12x - 30$$

Differentiate again

$$f''(1) = 12(1) - 30$$

$$f''(4) = 12(4) - 30$$

Substitute in
$$x = 1$$

Substitute in
$$x = 4$$

$$= -18$$

$$= 18$$


So (1, 17) is a maximum and (4, -10) is a minimum

HINT

'Minima' and 'maxima' are the plural forms of 'minimum' and 'maximum'.

Note: This is an extension activity and will not be in the exam.

Consider the curve $f(x) = 6x - x^2$ below. The gradient of the tangent at any point is given by f'(x) = 6 - 2x.



At the point f'(1) the gradient is 4. When the tangent has a positive gradient, the graph is going up and the function is increasing.

At the point f'(4) the gradient is -2. When the tangent has a negative gradient the graph is going down the curve and the function is decreasing.

At the point f'(3) the gradient is 0 and it is a stationary point.

KEY POINT

$$f'(x_0) > 0 \Rightarrow f$$
 is increasing at $x = x_0$

$$f'(x_0) < 0 \Rightarrow f$$
 is decreasing at $x = x_0$

$$f'(x_0) = 0 \Rightarrow f$$
 is stationary at $x = x_0$

EXERCISE 7

SKILLS DECISION MAKING 1 Find the values of x for which $\frac{dy}{dx} = 0$

$$y = x^2 + 12x$$

b
$$y = 5x^2 + 3x - 2$$

$$y = 4x^2 - 3x - 2$$

d
$$y = 2x + \frac{18}{x}$$

2 Find the *x*-coordinates of the points where the gradient is zero. Establish whether these points are maximum or minimum points.

$$u = 6x^2 + 2x$$

b
$$y = 10 + 7x - 6x^2$$

$$y = 10 - 2x + 4x^2$$

d
$$y = x^3 - x^2 - x + 1$$

e
$$y = x + \frac{1}{x}$$

$$y = x^2 + \frac{54}{x}$$

APPLY WHAT YOU HAVE LEARNT ABOUT TURNING POINTS TO SOLVE PROBLEMS

Whenever you see a question asking about the maximum value or minimum value of a quantity, you are likely to use differentiation at some point. Most questions will involve creating a formula, for example for volume or area, and then calculating the maximum value of it.

EXAMPLE 19

a A large tank (shown) is to be made from $54\,\mathrm{m}^2$ of sheet metal. It has no top. Show that the volume of the tank is given by $V=18x-\frac{2}{3}x^3$



- **b** Calculate the values of *x* that will give the largest volume possible, and calculate what this volume is.
- Justify that the value of V you have found is a maximum.

a Step 1

Try to make formulae using the information you have.

$$V = x^2y$$

Formula for volume

$$SA = 2x^2 + 3xy$$

Formula for the surface area (no top)

Step 2

Find a way to remove a constant, in this case y. You can rewrite the surface area formula in terms of y.

$$2x^2 + 3xy = 54$$
 so $54 - 2x^2 = 3xy$ and $y = \frac{54 - 2x^2}{3x}$

Step 3

Substitute the surface area formula into the volume formula, to replace y.

$$V = x^2y$$
 so $V = x^2\left(\frac{54 - 2x^2}{3x}\right)$ and $V = \frac{54x^2 - 2x^4}{3x}$

$$V=18x-\frac{2}{3}x^3$$

Simplify

b
$$\frac{dV}{dx} = 18 - 2x^2$$

Differentiate to find the derivative function

$$18 - 2x^2 = 0$$

Set the derivative function equal to 0 (since you are looking for the maximum)

$$2x^{2} = 18$$

$$x = \pm 3$$

$$V = 18(3) - \frac{2}{3}(3)^{3}$$

$$= 36 \,\mathrm{m}^{2}$$

There is a maximum at x = +3 (since a length cannot be negative)

Substitute x = 3 into the formula for volume


$$\frac{d^2V}{dx^2} = -4x$$

When
$$x = 3$$
, $\frac{d^2V}{dx^2} = -4 \times 3 = -12$

This is negative, so V = 36 is the maximum value of V.

EXAMPLE 20

The diagram shows a minor sector OMN of a circle with centre O and radius r cm. The perimeter of the sector is $100 \, \text{cm}$ and the area of the sector is $A \, \text{cm}^2$.



a Show that $A = 50r - r^2$

Given that r varies, find

- **b** The value of r for which A is a maximum and show that A is a maximum.
- The value of $\angle MON$ for this maximum area.
- The maximum area of the sector OMN.
- a Let the perimeter of the sector be P, so

$$P = 2r + r\theta$$

This is the sum of two radii (2r) and an arc length $MN(r\theta)$

$$\theta = \frac{100 - 2r}{r}$$
 equation (1)

Rearrange and substitute P = 100

Let the area of the sector be A, so

$$A = \frac{1}{2}r^2\theta \qquad \text{(equation 2)}$$

Substitute equation 1 into equation 2

$$A = \frac{1}{2}r^2\left(\frac{100 - 2r}{r}\right)$$
$$= 50r - r^2$$

The area formula is in terms of two variables r and θ , so you need to substitute for θ so that the formula is in terms of one variable, r

$$\frac{dA}{dr} = 50 - 2r$$

When
$$\frac{dA}{dr} = 0$$
, $r = 25$

Also $\frac{d^2A}{dr^2} = -2$, which is negative

Use the method which you learned to find stationary values: put the first derivative equal to zero, then check the sign of the second derivative

So the area is a maximum when r = 25.

$$\theta = \frac{100 - 50}{25}$$

Substitute
$$r = 25$$
 into $\theta = \frac{100 - 2r}{r}$

So angle MON = 2 radians

d The maximum value of the area is

$$50 \times 25 - 25^2 = 625 \, \text{cm}^2$$

Use
$$A = 50r - r^2$$

EXERCISE 8

SKILLS

CRITICAL THINKING PROBLEM SOLVING A rectangular field is **fenced** on three sides. A house forms the fourth side of the rectangle. Given that the total length of the fence is 80 m, show that the area, A, of the field is given by the formula A = y(80 − 2y), where y is the perpendicular distance from the house to the end of the field.

Given that the area is a maximum for this length of fence, find the dimensions of the **enclosed** field and the area which is enclosed.



A closed cylinder has a total surface area equal to 600π . Show that the volume, $V \text{cm}^3$, of this cylinder is given by the formula $V = 300\pi r - \pi r^3$, where r cm is the radius of the cylinder.

Find the maximum volume of such a cylinder.


3 ► A shape consists of a rectangular base with a semi-circular top, as shown. Given that the perimeter of the shape is 40 cm, show that its area is given by the formula

$$A=40r-2r^3-rac{\pi r^2}{2}$$
 where r cm is the radius of the semicircle.

Find and justify the maximum value for this area.



- 4 A wire of length 2 m is bent into the shape shown, made up of a rectangle and a semi-circle.
 - a Find an expression for y in terms of x.
 - **b** Show that the area is given by $A = \frac{x}{8}(8 4x \pi x)$




5 The shape shown is a wire frame in the form of a large rectangle split by parallel lengths of wire into 12 smaller equal-sized rectangles.

The total length of wire used to complete the whole frame is $1512 \,\mathrm{mm}$ and the width of one of the smaller rectangles is $x \,\mathrm{mm}$. Show that the area of the whole shape is $A \,\mathrm{mm}^2$,

where
$$A = 1296x - \frac{108x^2}{7}$$

Find and justify the maximum area which can be enclosed in this way.



EXAM PRACTICE: CHAPTER 9

1
$$f(x) = x^3 + 3x$$
 $x > 0$

- a Differentiate to find f'(x). [2]
- **b** Given that f'(x) = 15, find the value of x.
- Given that $y = 5x 7 + \frac{2\sqrt{x} + 3}{x}$ x > 0 find $\frac{dy}{dx}$
- f(x) = $\frac{(3-4\sqrt{x})(3-4\sqrt{x})}{x^{\frac{1}{2}}}$ x > 0
 - a Show that $f(x) = 9x^{-\frac{1}{2}} + Px^{\frac{1}{2}} + Q$, where P and Q are constants to be found. [2]
 - **b** Find f'(x). [4]
 - **c** Evaluate f'(9). [1]
- A curve C has equation $y = 2x^3 + kx^2 + 7$
 - a Find $\frac{dy}{dx}$ [2]
 - b The point P, where x = -2, lies on C. The tangent to C at the point P is parallel to the line with equation $y = \frac{17}{2}x + 5$ Find the value of k.
 - f c Find the value of the y-coordinate of P.
- The curve C has equation $y = x^3 2x^2 x + 9$ x > 0
 - **a** The point *P* has *x*-coordinate 2. If the *x*-coordinate lies on the curve, find the value of the *y*-coordinate. [1]
 - Find the equation of the tangent to C at P.The point Q also lies on C.[2]
 - Given that the tangent to C at Q is perpendicular to the tangent to C at P, find the x-coordinate of Q, giving your answer in surd form.
 [3]
- A curve has equation $y = \frac{x^2 5x 24}{x}$
 - a Find $\frac{dy}{dx}$ in its simplest form. [4]
 - **b** Find an equation of the tangent at the point x = 2. [3]
- Find $\frac{dy}{dx}$ if $y = 3\sin 5x + 4\cos 3x$
- Find $\frac{dy}{dx}$ if $y = \cos\left(\frac{x}{2}\right)$

$$f(x) = e^{2x} + 3x^2 + 4\sin 4x$$
. Find $f'(x)$

[3]

Find the gradient of the curve $y = 2e^{-2x}$ at point (0, 2).

[3]

[3]

- Find the gradient of the curve with equation $y = 4\cos 2x$ at the point where $x = \frac{\pi}{4}$. [3]
- $12 \quad f(x) = \frac{x+k}{x-2k}$

Find f'(x). [3]

13 $f(x) = x^2 + 3 + \frac{4}{(x-2)}$

Find the equation of the normal to the curve y = f(x) at the point where x = 3.

- Differentiate $y = 6e^{2x^3}$ [3]
- Differentiate $y = 3\sin^2 2x$ [4]

Q16 HINT

 $x \in \mathbb{R}$ means xis a real number. It can be positive, negative or zero, including integers, rational numbers

- The function $g(x) = \frac{e^x}{(e^x 2)^2} x \in \mathbb{R}, x \neq \ln 2$
 - a Find g'(x). [4]
 - **b** Find the value of x for which g'(x) = 1[2]
- Differentiate with respect to x, $\frac{\sin 4x}{x^3}$, giving your answer in its simplest form. [3]
- Differentiate $x^2 \cos 3x$ with respect to x. [3]
- Given $f(x) = (3x^2 1)(x^2 + 5x + 2)$, find the derivative of f(x). [3]
- Find the equation of the tangent to the curve $f(x) = x \sin x$ where $x = \frac{3}{2}\pi$. [5]
- 21 A curve has the equation $y + 10 = 12x^{\frac{1}{2}} x^{\frac{3}{2}}$ [4]
 - a Find the coordinates of the turning point.
 - **b** Determine the nature of the turning point.
- $y = x^2 px^{\frac{1}{2}}$ [4]
 - a Show that $\frac{dy}{dx} = 2x \frac{1}{2}kx^{-\frac{1}{2}}$
 - **b** Given that y is decreasing at x = 4, find the possible set of values for k

- A sector *OMN* of a circle has area 100 cm².
 - a Show that the perimeter of this sector is given by the formula $P=2r+\frac{200}{r},\,r>\sqrt{\frac{100}{\pi}}$

[5]

b Find the minimum value for the perimeter of such a sector.

CHAPTER SUMMARY: CHAPTER 9

You should know, and be able to use, these standard formulae:

| Function | Derivative |
|-----------|------------|
| x^n | nx^{n-1} |
| $\sin ax$ | $a\cos ax$ |
| $\cos ax$ | $-a\sin x$ |
| e^{ax} | ae^{ax} |

- You also need to know these rules.
 - Chain rule:

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u} \times \frac{\mathrm{d}u}{\mathrm{d}x}$$

■ Product rule:

If
$$y = uv$$
 then $\frac{dy}{dx} = u\frac{dv}{dx} + v\frac{du}{dx}$

■ Quotient rule:


If
$$y = \frac{u}{v}$$
 then $\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$

A turning point is a point where $\frac{dy}{dx} = 0$

If
$$\frac{dy}{dx} = 0$$
 and $\frac{d^2y}{dx^2} > 0$ then the point is a minimum.

If
$$\frac{dy}{dx} = 0$$
 and $\frac{d^2y}{dx^2} < 0$ then the point is a maximum.

CHAPTER 10



10 INTEGRATION

Integration has many uses but it is especially important in *kinematics*. Integration helps engineers understand the relationship between displacement, velocity and *acceleration*.

You already know how to find the area of common shapes, but how do you find the area under a curve? Integration helps you solve this problem and you'll learn about it in this chapter. You will also take this a step further to calculate the *volume* of solids which have curved surfaces.



LEARNING OBJECTIVES

- Use integration to reverse the process of differentiation
- Understand how calculus is related to problems involving kinematics
- Use integration to find areas
- Use Integration to find the *volume of revolution*
- Relate rates of change to each other

STARTER ACTIVITIES

Copy and complete the table below.

| f(x) | GRADIENT FUNCTION f'(x) | f'(2) | f'(-1) | f'(0) |
|---------------------------------|-------------------------|-------|--------|-------|
| y = (x+3)(x-6) | | | | |
| $y = \frac{x^2 + x^3}{x}$ | | | | |
| $y=x(x^5-7x)$ | | | | |
| $y = \frac{x^2 \times x^3}{x}$ | | | | |
| $y = \frac{1}{4}x + 2$ | | | | |
| $y = \frac{(x^2 + 3x)(1-x)}{x}$ | | | | |

INTEGRATION AS THE REVERSE OF DIFFERENTIATION

The indefinite integral of x^n is denoted by $\int x^n dx$

This is the constant of integration and it is important in indefinite integrals because the derivative only tells you about the change that occurred. If you integrate the derivative, you will calculate the entire change that occurred over the range of integration. However, it doesn't tell you where you started, hence c is a generic 'starting point'

The expression to be integrated

 $\int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad n \neq 1$ The elongated S means integrate

The constant of integration has to be included in all indefinite integrals

The dx tells you which letter is the variable to integrate with respect to

The rules for integration you need for this course are

KEY POINTS

$$\int x^n dx = \frac{x^{n+1}}{n+1} + c, \quad (n \neq 1)$$

This is the one you will most commonly meet

KEY POINTS

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

This arises from the equation showing the unusual character of e: $\int e^x dx = e^x + c$

KEY POINTS

$$\int \sin x \, dx = -\cos x + c$$

Notice the minus sign

KEY POINTS

$$\int \sin ax \, \mathrm{d}x = -\frac{1}{a} \cos ax + c$$

KEY POINTS

$$\int \cos x \, \mathrm{d}x = \sin x + c$$

No minus sign

KEY POINTS

$$\int \cos ax \, \mathrm{d}x = \frac{1}{a} \sin ax + c$$

EXAMPLE 1

Integrate

a
$$\int 2x^3 dx$$

$$\frac{dy}{dx} = 6x + 2x^{-3} - 3x^{\frac{1}{2}}$$

d
$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}} - \frac{1}{2}x^{-\frac{3}{2}}$$

$$\int (x^{\frac{1}{2}} + 2x^3) dx$$

$$\int (3x^2 + px^{-2} + q^2) dx$$

a $\int 2x^3$ $=\frac{2x^4}{4}+c$ $=\frac{1}{2}x^4+c$

Increase the power by one, and divide by the new power

$$\int 3x^{\frac{1}{2}} \\
= \frac{3x^{\frac{3}{2}}}{\frac{3}{2}} + c \\
= 2x^{\frac{3}{2}} + c$$

Increase the power by one, and divide by the new power

c
$$\frac{dy}{dx} = 6x + 2x^{-3} - 3x^{\frac{1}{2}}$$

$$y = \frac{6}{2}x^2 + \frac{2}{-2}x^{-2} - \frac{3}{\frac{3}{2}}x^{\frac{3}{2}} + c$$

Integrate each part separately

$$= 3x^{2} - x^{-2} - 2x^{\frac{3}{2}} + c$$
$$= 3x^{2} - \frac{1}{x^{2}} - 2\sqrt{x^{3}} + c$$

Simplify

d
$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}} - \frac{1}{2}x^{-\frac{3}{2}}$$

$$f(x) = \frac{\frac{1}{2}x^{\frac{1}{2}}}{\frac{1}{2}} - \frac{\frac{1}{2}x^{-\frac{1}{2}}}{-\frac{1}{2}} + c$$

Integrate each part separately

$$= x^{\frac{1}{2}} + x^{-\frac{1}{2}} + c$$
$$= \sqrt{x} + \frac{1}{\sqrt{x}} + c$$

Simplify if possible

e
$$\int (x^{\frac{1}{2}} + 2x^3) dx$$

= $\frac{2}{3}x^{\frac{3}{2}} + \frac{2}{4}x^4 + c$

Integrate each part separately

$$3^{4} + \frac{2}{3}\sqrt{x^{3}} + \frac{1}{2}x^{4} + c$$

Simplify

HINT

p and q^2 should be treated as if they were just numbers.

EXAMPLE 2

$$\int (3x^2 + px^{-2} + q^2) dx$$

Integrate each part separately

$$= \frac{3}{3}x^3 + \frac{p}{-1}x^{-1} + q^2x + c$$

Simplify

$$=x^3-\frac{p}{x}+q^2x+c$$

Find $\int (2\cos x + 3e^{-x} - \sin 2x) dx$

$$\int (2\cos x + 3e^{-x} - \sin 2x) dx = \int 2\cos x dx + \int 3e^{-x} dx - \int \sin 2x dx$$
$$= 2\sin x - 3e^{-x} + \frac{1}{2}\cos 2x + c$$

Integrate each term separately using standard integration rules

SKILLS

REASONING

Integrate with respect to x

$$a \int (x^3 + 3x) dx$$

d
$$\int (5x^{\frac{1}{2}} - 3x^2) dx$$

$$\int (4x^2 - 3x^{-4} + r) dx$$

b
$$\int (4x^3 - 2x) dx$$
 c $\int (3x^{-2} + 4) dx$

$$= \int (30x^2 - x^{-3}) dx$$

d
$$\int (5x^{\frac{1}{2}} - 3x^2) dx$$
 e $\int (30x^2 - x^{-3}) dx$ **f** $\int (x^2 + x^{-\frac{1}{2}} - x^{\frac{3}{2}} + 4) dx$

h
$$\int (px^4 + 2t + 3x^{-2}) dx$$

g
$$\int (4x^2 - 3x^{-4} + r) dx$$
 h $\int (px^4 + 2t + 3x^{-2}) dx$ i $\int \left(\frac{3 - x^3}{2^{x^3}}\right) dx$

2 Integrate with respect to x

a
$$\int (2x + 3)x^2 dx$$

d
$$\int \left(\frac{2x+3}{\sqrt{x}}\right) dx$$

$$\int \left(\frac{2x^2+3}{x^2}\right) dx$$

e
$$\int \left(\frac{1}{4x^3} - \frac{3}{x^2}\right) dx$$

c
$$\int (2x + 3)^2 dx$$

f $\int (x^{\frac{5}{3}} + x^{-\frac{7}{4}}) dx$

3 Integrate with respect to x

$$a 2\cos x$$

d
$$3e^{4x}$$

g
$$\sin \frac{1}{2}x$$

 \mathbf{b} 2 sin 3x

$$h 5 \cos 3x$$

c 4ex

f
$$5e^{-3x}$$

$$i$$
 4 sin 3 x

4 Integrate with respect to x

a
$$2(\cos x - \sin x + 3x)$$
 b $7e^x + 3\sin x - 2x^{-2}$ **c** $e^x + \sin x + 2\cos x$

$$7e^x + 3\sin x - 2x^{-2}$$

$$e^x + \sin x + 2\cos x$$

d
$$3x^2 + 2\sin 2x + 3e^{4x}$$
 e $4(\cos x + e^x + x^2)$

UNDERSTAND HOW CALCULUS IS RELATED TO PROBLEMS INVOLVING **KINEMATICS**

KEY POINTS

You need to understand how displacement (s), velocity (v) and acceleration (a) can relate to calculus. The relationship between these variables in calculus is:

$$v = \frac{ds}{dt}$$

and
$$a = \frac{di}{dt}$$

and
$$a = \frac{dv}{dt}$$
 (or $a = \frac{d^2s}{dt^2}$. This is known as a second derivative.)

These can also be written as

$$s = \int v dt$$

and
$$v = \int a dt$$

EXAMPLE 3

The velocity, $v \, \mathrm{m} \, \mathrm{s}^{-1}$, of a ball after t seconds, is given by $v = 8 + 10t - t^2$. ($t \ge 0$)

- a Find the acceleration after t seconds.
- Calculate t when the acceleration is zero.
- Hence, or otherwise, find the maximum velocity.

$$a \quad a = \frac{\mathrm{d}v}{\mathrm{d}t} = 10 - 2t$$

Integrate the function for v to find a

b
$$a = 10 - 2t$$

$$0 = 10 - 2t$$

Substitute a = 0

$$2t = 10$$

$$t = 5$$
 seconds

Solve

c The maximum velocity occurs when $\frac{dv}{dt} = 0$ and t = 5.

Therefore, $v_{\text{max}} = 8 + 50 - 25 = 33 \,\text{m}\,\text{s}^{-1}$.

EXAMPLE 4

The velocity, $v \text{ m s}^{-1}$, of a particle after t seconds is given by

$$v = 12t - 8t^3 \quad (t \ge 0)$$

Given that the initial displacement is 10 m, find

an expression for s in terms of t

b the displacement when t = 2.

a
$$s = \int v dt$$

= $\int (12t - 8t^3) dt$
= $6t^2 - 2t^4 + c$

$$10 = 6(0)^2 - 2(0)^4 + c$$

$$c = 10$$

Thus,
$$s = 6t^2 - 2t^4 + 10$$

b When
$$t=2$$

$$s = 6(2)^2 - 2(2)^4 + 10$$

$$s = 2 \,\mathrm{m}$$

To find *c* substitute initial conditions t = 0 and s = 10 into $v = 12t - 8t^3$

Substitute t = 2 into $s = 6t^2 - 2t^4 + 10$

EXERCISE 2

SKILLS

CRITICAL

- ▶ The displacement, s metres, of a particle after t seconds is given by $s = 250 + 10t^2$. Find an expression for the velocity.
- **2** The displacement, s metres, of a particle after t seconds is given by $s = 60 + 48t 16t^2$. Find an expression for the velocity.
- 3 \triangleright The displacement, s metres, of a particle after t seconds is given by $s = 25 + 30t 5t^2$
 - a Find an expression for the velocity.
 - **b** Work out the velocity after 3 seconds. Clearly state your units.
- 4 \triangleright The velocity, $v \text{ m s}^{-1}$, of a particle after t seconds is given by v = 160 32t.
 - a Find an expression for the acceleration.
 - **b** At t = 0 the particle's displacement is 384 m from its starting point. Find
 - an expression for this displacement
 - ii the time when the particle passes through the origin.
- 5 The velocity, $v \, \text{m} \, \text{s}^{-1}$, of a particle after t seconds is given by $v = 24 + 6t t^2$
 - a Find an expression for its acceleration.
 - Calculate its acceleration after 2 seconds.
 - When s = 0, t = 0
 - **d** Find the displacement when t = 2.

USE INTEGRATION TO FIND AREAS

The definite integral is defined as

$$\int_a^b f'(x) dx = [f(x)]_a^b = f(a) - f(b)$$


a and b are called the 'limits' of the integral

provided f' is the derived function of f throughout the **interval** (a, b).

In particular if you wish to find the area between a curve, the x-axis and the lines x = a and x = b you use

Area =
$$\int_a^b y \, dx$$

where y = f(x) is the equation of the curve.




EXAMPLE 5

Find the area of the region R bounded by the curve with equation y = (4 - x)(x + 2) and the positive x- and y-axes.

When
$$x = 0$$
, $y = 8$

When
$$y = 0$$
, $x = 4$ or -2



The area of R is given by

$$A = \int_{0}^{4} (4 - x)(x + 2) dx$$
So
$$A = \int_{0}^{4} (8 + 2x - x^{2}) dx$$

$$A = \left[8x + x^{2} - \frac{1}{3}x^{3} \right]_{0}^{4}$$

$$= \left(8(4) + 4^{2} - \frac{4^{3}}{3} \right) - \left(8(0) + 0^{2} - \frac{0^{3}}{3} \right)$$

$$= \left(32 + 16 - \frac{64}{3} \right) - (0)$$

So the area is $26\frac{2}{3}$ square units

Expand the brackets to leave the function in a form that you can integrate

Integrate

Substitute in limits of 4 and 0

EXAMPLE 6

The region R is enclosed by the curve with equation $y = x^2 + \frac{4}{x^2}$, x > 0, the x-axis and the lines x = 1 and x = 3. Find the area of R.

Area =
$$\int_{1}^{3} \left(x^2 + \frac{4}{x^2}\right) dx$$

This curve is not one you would be expected to sketch but the limits of the integral are simply x = 1 and x = 3, so you can write down an expression for the area without referring to a sketch

$$=\int_{1}^{3}(x^{2}+4x^{-2})\mathrm{d}x$$
 Write the expression in a form suitable for integrating
$$=\left[\frac{1}{3}x^{3}-4x^{-1}\right]_{1}^{3}$$
 Integrate
$$=\left[\frac{1}{3}x^{3}-\frac{4}{x}\right]_{1}^{3}$$
 Simplify
$$=\left(\frac{1}{3}(3)^{3}-\frac{4}{3}\right)-\left(\frac{1}{3}(1)^{3}-\frac{4}{1}\right)$$
 Substitute in the limits 3 and 1
$$=\left(9-\frac{4}{3}\right)-\left(\frac{1}{3}-4\right)$$
 Simplify
$$=13-\frac{5}{3}$$


$$=11\frac{1}{3}$$

You may be able to use a **graphical calculator** to check your answer, but you must show your working.

In the examples thus far, the area that was calculated was above the x-axis. If the area between the curve and the x-axis lies below the x-axis, then $\int y dx$ will give a negative answer.

EXAMPLE 7

Find the area of the finite region bounded by the curve y = x(x - 3) and the x-axis.



First draw a sketch. The solutions to y = x(x - 3) are x = 0 and x = 3. So the curve crosses the x-axis - (0, 0) and (3, 0)

In this case you can see that the region is *below* the *x*-axis

Area =
$$\int_0^3 x(x-3) dx$$

= $\int_0^3 (x^2-3x) dx$ Multiply out the brackets
= $\left[\frac{1}{3}x^3-\frac{3}{2}x^2\right]_0^3$ Integrate and use square brackets
= $\left(\frac{1}{3}(3)^3-\frac{3}{2}(3)^2\right)-\left(\frac{1}{3}(0)^3-\frac{3}{2}(0)^2\right)$ Substitute in the limits

$$= \left(9 - \frac{27}{2}\right) - 0$$
$$= -\frac{9}{2}$$

Thus, the area is $\frac{9}{2}$ square units.

Work out each part

The minus sign indicates that the area is below the x-axis


EXAMPLE 8

Find the area between the curve y = x(x + 1)(x - 1) and the *x*-axis.

First let's try this in the same way as in the previous examples.

Start with a sketch.

You can see this time that part of the curve is above the x-axis and part is below



Area =
$$\int_{-1}^{1} x(x + 1)(x - 1) dx$$

= $\int_{-1}^{1} (x^3 - x) dx$
= $\left[\frac{1}{4}x^4 - \frac{1}{2}x^2\right]_{-1}^{1}$
= $\left(\frac{1}{4}(1)^4 - \frac{1}{2}(1)^2\right) - \left(\frac{1}{4}(-1)^4 - \frac{1}{2}(-1)^2\right)$
= $\left(-\frac{1}{4}\right) - \left(-\frac{1}{4}\right)$
= 0

Expand the brackets

Integrate and use square brackets

Use the limits

From the sketch you can see that this is clearly not the correct answer and this method will not work. As a result, you must approach this type of question differently. You need to integrate each section separately and then combine them as positive values.

Area
$$=\int_{-1}^{1} x(x+1)(x-1) dx = \int_{-1}^{0} x(x+1)(x-1) dx + \int_{0}^{1} x(x+1)(x-1) dx$$

Area above the x-axis Area below the x-axis

Area above the *x*-axis =
$$\int_{-1}^{0} x(x+1)(x-1) dx$$

= $\int_{-1}^{0} (x^3 - x) dx$
= $\left[\frac{1}{4}x^4 - \frac{1}{2}x^2\right]_{-1}^{0}$
= $\left(\frac{1}{4}(0)^4 - \frac{1}{2}(0)^2\right) - \left(\frac{1}{4}(-1)^4 - \frac{1}{2}(-1)^2\right)$
= $0 - \left(-\frac{1}{4}\right)$
= $\frac{1}{4}$

Use the limits -1 and 0 as the curve lies above the x-axis in this region

Expand the brackets

Integrate

Use the limits

So the area above the *x*-axis is $\frac{1}{4}$ square unit.

Area below the x-axis =
$$\int_0^1 x(x+1)(x-1) dx$$

$$= \int_0^1 (x^3 - x) dx$$

$$= \left[\frac{1}{4} x^4 - \frac{1}{2} x^2 \right]_0^1$$

$$= \left(\frac{1}{4} (1)^4 - \frac{1}{2} (1)^2 \right) - \left(\frac{1}{4} (0)^4 - \frac{1}{2} (0)^2 \right)$$

$$= -\frac{1}{4} - 0$$

$$= -\frac{1}{4}$$

Expand the brackets

Integrate


Use the limits

The minus sign indicates that the area is beneath the *x*-axis

So the area below the *x*-axis is $\frac{1}{4}$ square unit.

The total area of the curve between x = -1 and x = 1 is $\frac{1}{4} + \frac{1}{4} = \frac{1}{2}$ square unit.

Sometimes you may need to find the area between a curve and a line, or between two curves.




KEN BUINTS

The area between a line (y_1) and a curve (y_2) is given by:

Area =
$$\int_a^b y_1 dx - \int_a^b y_2 dx$$

EXAMPLE 9

Below is a diagram showing the line y = x and the curve y = x(4 - x). Find the area bounded by the line and the curve.



$$x = x(4-x)$$

$$x = 4x - x^2$$

$$x^2 - 3x = 0$$

$$x(x-3)=0$$

$$x = 0 \text{ or } x = 3$$

Equate the equation of the line and the equation of the curve to find the *x*-values where the line crosses the curve

Expand the bracket

Area =
$$\int_0^3 (x(4-x) dx - \int_0^3 x dx$$

Use $\int_a^b y_1 dx - \int_a^b y_2 dx$ to find the area

$$=\int_0^3 (3x - x^2) dx$$

$$= \left[\frac{3}{2}x^2 - \frac{1}{3}x^3 \right]_0^3$$

$$= \left(\frac{3}{2}(3)^2 - \frac{1}{3}(3)^3\right) - \left(\frac{3}{2}(0)^2 - \frac{1}{3}(0)^3\right)$$

$$=\left(\frac{27}{2}-9\right)-0$$

$$=\frac{9}{2}$$
 square units

SKILLS

DECISION MAKING CRITICAL THINKING Sketch these curves and find the area of the finite region bounded by the curves and the

a
$$y = x(x - 2)$$

b
$$y = (x + 2)(x - 3)$$

$$y = -(x + 4)(x - 1)$$

d
$$y = x(x-2)(x-5)$$

2 Find the area between the curve with equation y = f(x), the x-axis and the lines x = aand x = b in each case.

a
$$f(x) = 4x^2 - 2x + 2$$

b
$$f(x) = 3x - x^2$$
 $a = -1, b = 1$

c
$$f(x) = x^3 - 4x$$

$$a = 1, b = 2$$

a
$$f(x) = 4x^2 - 2x + 2$$
 $a = 0, b = 3$
b $f(x) = 3x - x^2$ $a = -1, b = 2$
c $f(x) = x^3 - 4x$ $a = 1, b = 2$
d $f(x) = \frac{4x^4 - x}{2x}$ $a = -1, b = 2$

e
$$f(x) = 8x^{-3} + \sqrt{x}$$
 $a = 1, b = 4$

$$a = 1, b = 4$$

3 Find the area of the region between the curves or lines represented by these equations.

a
$$y = x^2$$
 and $y = 2x$

b
$$y = x^2$$
 and $y = 4$

$$y = x(2 - x)$$
 and $y = x$

d
$$y = x^2 - 2$$
 and $y = \frac{1}{2}x^2$

$$y = x^2 - 1$$
 and $y = x + 1$

Evaluate

a
$$\int_{1}^{2} 3x^2 dx$$

b
$$\int_{1}^{2} \frac{1}{x^{2}} dx$$

c
$$\int_{4}^{9} \frac{1}{x^{\frac{1}{2}}} dx$$


d
$$\int_{0}^{4} (x^3 - 2x - 3x^{\frac{1}{2}}) dx$$

Find the area enclosed by the curve $y = x^2 - x - 1$ and the line y = 5.


Find the area between $y = 1 + 4x^{-2}$ and the x-axis from x = 1 to x = 2.

7 The diagram shows the curve $y = x^2$ and the line y = 9, with the enclosed region shaded.

- a Find the points of intersection A and B.
- **b** Find the area of the shaded region.




- 8 The diagram shows the curve $y = x^2 2x + 1$ and the line y = x + 1, with the enclosed region shaded.
 - a Find the points of intersection of the curve and the line.
 - **b** Find the area of the shaded region.




9 The diagram shows the curve $y = x^2 + 3x$.

Find the area of the shaded region between x = -1 and x = 2.




- The diagram shows the finite region R, bounded by the curve with equation y = x(4 + x), the line y = 12 and the y-axis.
 - a Find the coordinates of point A.
 - **b** Show that the area of the shaded region is $13\frac{1}{3}$ units.




11 Show that the area between the curve $y = 4x - x^2$ and the x-axis from x = 0 to x = 5 is $8\frac{1}{3}$

USE INTEGRATION TO FIND A VOLUME OF REVOLUTION

You can also use integration to find volumes.






KEY POINTS

The volume of revolution of a solid rotated 2π radians around the x-axis between x=a and x=b is given by: Volume $=\pi\int_a^b y^2 dx$

EXAMPLE 10

Find the volume of revolution formed by rotating the area enclosed by the curve $y = x + x^2$, the x-axis and the coordinates x = 1, x = 3.

First sketch the curve.



$$V = \pi \int_1^3 (x + x^2)^2 \, \mathrm{d}x$$

$$= \pi \int_{3}^{3} (x^2 + 2x^3 + x^4) dx$$

$$= \pi \left[\frac{1}{3}x^3 + \frac{1}{2}x^4 + \frac{1}{5}x^5 \right]_1^3$$

$$= \pi \left[\left(\frac{1}{3} (3)^3 + \frac{1}{2} (3)^4 + \frac{1}{5} (3)^5 \right) - \left(\frac{1}{3} (1)^3 + \frac{1}{2} (1)^4 + \frac{1}{5} (1)^5 \right) \right]$$

$$= \pi \left[\frac{1557}{20} - \frac{47}{60} \right]$$

$$=77\frac{1}{15}\pi$$
 cubic units

Use $V = \pi \int_{a}^{b} y^2 dx$

Expand

Integrate

Substitute in the limits

Simplify

EXAMPLE 11

The region R is bounded by the curve with the equation $y = x^2 - 1$, the x-axis and the lines x = 1and x = -1.

- Find the area of R
- Find the volume of the solid formed when the region R is rotated through 2π radians about the

a Area =
$$\int_{-1}^{1} (x^2 - 1) dx$$

= $[2x - x]_{-1}^{1}$
= $(2(1) - (1)) - (2(-1) - (-1))$
= 2

b
$$V = \pi \int_{\pi}^{b} y^{2} dx$$

$$= \pi \int_{\pi}^{1} (x^{2} - 1)^{2} dx$$

$$= \pi \int_{-1}^{1} (x^{4} - 2x^{2} + 1) dx$$

$$= \pi \left[\frac{x^{5}}{5} - \frac{2x^{3}}{3} + x \right]^{1}$$

$$= \pi \left\{ \left(\frac{1}{5} - \frac{2}{3} + 1 \right) - \left(-\frac{1}{5} + \frac{2}{3} - 1 \right) \right\}$$
$$= \frac{16\pi}{15}$$

EXAMPLE 12

The region R is bounded by the curve with the equation $y = \sin 2x$, the x-axis and the lines x = 0 and $x = \frac{\pi}{2}$.

- a Find the area of R.
- **b** Find the volume of the solid formed when the region R is rotated through 2π radians about the x-axis.
- a Area = $\int_{\frac{\pi}{2}}^{0} \sin 2x \, dx$ = $\left[-\frac{1}{2} \cos 2x \right]_{0}^{\frac{\pi}{2}}$ = $\left(\left(-\frac{1}{2} \right) (-1) \right) \left(\left(-\frac{1}{2} \right) (1) \right)$ = 1
- b Volume = $\pi \int_{\frac{\pi}{2}}^{0} \sin^{2} 2x \, dx$ = $\left(\frac{1}{2}\right) \pi \int_{\frac{\pi}{2}}^{0} \frac{1}{2} (1 - \cos 4x) \, dx$ = $\left(\frac{1}{2}\right) \pi \left[x - \frac{1}{4} \sin 4x\right]_{0}^{\frac{\pi}{2}}$ = $\left(\frac{1}{2}\right) \pi \left(\frac{\pi}{2} - 0\right) - (0)$ = $\frac{\pi^{2}}{4}$

Use $\cos 2A = 1 - 2\sin^2 A$ Rearrange to give $\sin^2 A = \dots$

Note that $2 \times 2x$ gives 4x in the cos term

Multiply out and integrate

EXERCISE 4

SKILLS

PROBLEM SOLVING INTERPRETATION

- 1 Find the volume of the solid formed by rotating the area enclosed by the curve $y = x^2$, the x-axis and the line x = 2.
- Find the volume of the solid formed by rotating the area enclosed by the curve $y = x^3$, the x-axis and the line x = 2.
- **3** Find the volume of the solid formed by rotating the area enclosed by the curve $y = x^3 + 1$, the x-axis and the line x = 2.
- 4 The region R is bounded by the curve with equation y = f(x), the x-axis and the lines x = a and x = b. In each part find the exact value of:
 - i the area of R,
 - ii the volume of the solid of revolution formed by rotating R through 2π radians about the x-axis.

a
$$f(x) = \frac{2}{1+x}$$
; $a = 0$, $b = 1$

b
$$f(x) = \ln x$$
; $a = 1$, $b = 2$

c
$$f(x) = x\sqrt{4-x^2}$$
; $a = 0$, $b = 2$

RELATE RATES OF CHANGE TO EACH OTHER

Differential equations arise from many problems in mechanics, physics, chemistry, biology and economics. As their name suggests, these equations involve differential coefficients and so equations of the form

$$\frac{dy}{dx} = 3y$$
, $\frac{ds}{dt} = 2 + 6t$, $\frac{d^2y}{dt^2} = -25y$ $\frac{dP}{dt} = 10 - 4P$

are differential equations (x, y, t, s) and P are variables).

KEY POINTS


You can set up simple differential equations from information given in context. This may involve using connected rates of change, or ideas of proportion.

EXAMPLE 13

The image shows a right circular cylindrical metal rod which is expanding as it is heated. After t seconds the radius of the rod is x cm and the length of the rod is 5x cm.

$$rac{\mathrm{d}A}{\mathrm{d}t}=$$
 0.032 cm 2 s $^{-1}$ and $rac{\mathrm{d}A}{\mathrm{d}x}=$ 2 πx

Find $\frac{dx}{dt}$ when the radius of the rod is 2 cm, giving your answer to 3 significant figures.




$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{0.016}{\pi x}$$

When $x = 2 \, \text{cm}$

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{0.016}{2\pi}$$

$$\frac{\mathrm{d}x}{\mathrm{d}t} = 0.00255\,\mathrm{cm}\,\mathrm{s}^{-1}$$



EXAMPLE 14

A vase with a circular cross-section is shown. Fluid is flowing into the vase.


When the depth of the water is h cm, the volume of water Vcm3 is given by

$$V=4\pi h(h+4), \qquad 0 \leqslant h \leqslant 25$$

Water flows into the vase at a constant rate of $80\pi \text{cm}^3\text{s}^{-1}$ and

$$\frac{\mathrm{d}V}{\mathrm{d}t} = 80\pi \text{ and } \frac{\mathrm{d}V}{\mathrm{d}h} = 8\pi h + 16\pi$$

Find the rate of change of the depth of the water, in cm s⁻¹, when h=6



Find dh

$$\frac{dh}{dt} = \frac{dV}{dt} \div \frac{dV}{dh}$$
$$\frac{dh}{dt} = 80\pi \times \frac{1}{8\pi h + 16\pi}$$

When
$$h = 6$$

$$\frac{\mathrm{d}h}{\mathrm{d}t} = 80\pi \times \frac{1}{8\pi h + 16\pi}$$

$$\frac{dh}{dt} = \frac{80\pi}{64}$$

$$\frac{\mathrm{d}h}{\mathrm{d}t} = 1.25\,\mathrm{cm}\,\mathrm{s}^{-1}$$

You can relate one rate of change to another. This is useful when a situation involves more than two variables.

EXAMPLE 15

Given that the area of a circle, A, is related to its radius r by the formula $A=\pi r^2$, and the rate of change of its radius in cm is given by $\frac{\mathrm{d}r}{\mathrm{d}t}=5$, find $\frac{\mathrm{d}A}{\mathrm{d}t}$ when r=3

$$A = \pi r^2$$

$$\frac{dA}{dr} = 2\pi r$$

As A is a function of r integrate to find $\frac{dA}{dr}$

Using
$$\frac{\mathrm{d}A}{\mathrm{d}t} = \frac{\mathrm{d}A}{\mathrm{d}r} \times \frac{\mathrm{d}r}{\mathrm{d}t}$$

$$\frac{\mathrm{d}A}{\mathrm{d}t} = 2\pi r \times 5$$

 $=30\pi$

Substitute in values of $\frac{dA}{dr}$ and $\frac{dr}{dt}$

when r = 3

EXAMPLE 16

The volume of a **hemisphere** V cm³ is related to its radius r cm by the formula $V=\frac{2}{3}\pi r^3$ and the total surface area S cm² is given by the formula $S=\pi r^2+2\pi r^2=3\pi r^2$. Given that the rate of increase in volume, in cm³ s⁻¹, $\frac{\mathrm{d}V}{\mathrm{d}t}=6$, find the rate of increase of surface area $\frac{\mathrm{d}S}{\mathrm{d}t}$.

$$V=rac{2}{3}\pi r^3$$
 and $S=3\pi r^2$

$$\frac{dV}{dr} = 2\pi r^2$$
 and $\frac{dS}{dr} = 6\pi r$

Now
$$\frac{\mathrm{d}S}{\mathrm{d}t} = \frac{\mathrm{d}S}{\mathrm{d}r} \times \frac{\mathrm{d}r}{\mathrm{d}V} \times \frac{\mathrm{d}V}{\mathrm{d}t}$$

$$= 6\pi r \times \frac{1}{2\pi r^2} \times 6$$

$$= \frac{18}{r}$$

This is area of circular base plus area of curved surface.

As V and S are functions of r, find $\frac{dV}{dr}$ and $\frac{dS}{dr}$

Use an extended chain rule together with the property that $\frac{\mathrm{d}r}{\mathrm{d}V}$ = 1 \div $\frac{\mathrm{d}V}{\mathrm{d}r}$

You can use this formula for finding approximations.

$$\delta y \approx \frac{\mathrm{d}y}{\mathrm{d}x} \times \delta x$$

This formula shows an approximate relationship between a small change in the variable, x, δx and the equivalent small change in the variable y, δy .

EXAMPLE 17

Given that $y = 4x^3$, find the approximate percentage change in y for a 1% change in x.

$$\delta x$$
 is 1% of x
so $\delta x = \frac{x}{100}$
 $\delta y \approx \frac{12x^3}{100}$

$$y = 4x^3$$

so
$$\frac{\mathrm{d}y}{\mathrm{d}x} = 12x^2$$

$$\delta y \approx 12 \, x^2 \times \frac{x}{100}$$

Differentiate

Substitute into
$$\delta y \approx \frac{\mathrm{d}y}{\mathrm{d}x} \times \delta x$$

The percentage change in y will be $\frac{\delta y}{u} \times 100$

Percentage change =
$$\frac{12x^3}{4x^3 \times 100} \times 100$$

= 3%

Substitute
$$\delta y \approx \frac{12 x^3}{100}$$

EXERCISE 5

SKILLS INTERPRETATION REASONING

- ▶ Given that $V = \frac{1}{3}\pi r^3$ and that $\frac{dV}{dt} = 8$ find $\frac{dr}{dt}$ when r = 3.
- ► Given that $y = xe^x$ and that $\frac{dx}{dt} = 5$ find $\frac{dy}{dt}$ when x = 2.
- Given that $r = 1 + 3\cos\theta$ and that $\frac{d\theta}{dt} = 3$, find $\frac{dr}{dt}$ when $\theta = \frac{\pi}{6}$.
- 5 If $y = 3x^2$, find the approximate percentage change in y due to a change of 1% in the value of x.
- 6 If $y = 9x^3$, find the approximate percentage change in y due to a change of 1% in the value of x.
- 7 In a study of the water loss of picked leaves the mass M grams of a single leaf was measured at times t days after the leaf was picked. It was found that the rate of loss of mass was proportional to the mass M of the leaf. Write down a differential equation for the rate of change of mass of the leaf.
- **8** A curve C has equation y = f(x), y > 0. At any point P on the curve, the gradient of C is proportional to the product of the x and the y coordinates of P. The point A with coordinates (4, 2) is on C and the gradient of C at A is $\frac{1}{2}$. Show that $\frac{dy}{dx} = \frac{xy}{16}$

Note: These are extension activities and will not be in the exam.

9 At time t seconds the surface area of a cube is $A \text{ cm}^2$ and the volume is $V \text{ cm}^3$.

The surface area of the cube is expanding at a constant rate of $2 \text{ cm}^2 \text{ s}^{-1}$.

Show that
$$\frac{dV}{dt} = \frac{1}{2}V^{\frac{1}{3}}$$
.

10 An inverted conical funnel is full of salt. The salt is allowed to leave by a small hole in the vertex. It leaves at a constant rate of 6 cm³ s⁻¹.

Given that the angle of the cone between the slanting edge and the vertical is 30°, show that the volume of the salt is $\frac{1}{9}\pi h^3$, where h is the height of salt at time t seconds.

Show that the rate of change of the height of the salt in the funnel is inversely proportional to h^2 . Write down the differential equation relating h and t.

EXAM PRACTICE: CHAPTER 10

 $\frac{dy}{dx} = (x^2 + 3)^2 \text{ and the point } C \text{ with coordinates (3, 20) lies on the curve } C.$

Find the equation of the curve in the form y = f(x).

[3]

The gradient function of a curve is $f'(x) = -x^3 + 2x^{-2} - \frac{5}{2}x^{-3}$ find the equation of the curve given that it passes through the point (1, 7).

[3]

Given that $y = 5x^3 - 5x^{-2}$, $x \ne 0$, find $\int y dx$

[2]

 $\frac{dy}{dx} = \frac{x+9}{\sqrt{x}}. \text{ Find } y.$


[3]

Integrate $5e^x - 4\cos x + 2x^3$

[4]

The diagram shows a sketch of the curve $y = 10 + 8x + x^2 - x^3$. The x-coordinate of point A is 2. The region R, shown shaded, is bounded by the curve, the y-axis and the line from O to A, where O is the origin.

Find the exact value of the shaded region R.



[5]


- The line y = 10 x intersects the curve with equation $y = 10x x^2 8$ at A and B.
 - Find the coordinates of A and B.

[2]

b Calculate the exact value of the region bounded by the curve and line.

[4]

The diagram shows the curve $y = 4 + 3x - x^2$



a Find the coordinates where the curve crosses the x-axis.


[2]

b Find the exact area of the shaded region.

[3]

[3]


The diagram shows the curve C with the equation y = x(x + 4)(x - 2). The graph intersects the x-axis at x = 0, x = -4 and x = 2. The finite region, shown shaded in the diagram, is bounded by the curve C and the x-axis. Find the total area of the finite region shown shaded in the diagram.




[4]

[4]

The diagram shows a sketch of the curve $y = x^2 - 3x + 2$. Find the sum of the areas. [4]



Find the area bounded by the line y = -x - 11 and the curve $y = 9 - x^2$ shown in the diagram.




- Find the volume of the solid formed when the area enclosed by the curve $y=x^3$, the x-axis and the line x=2 performs one revolution about the x-axis. [5]
- Find the volume of the solid formed when the area enclosed by the curve $y = x^3 + 1$, the x-axis and the line x = 1 performs one revolution about the x-axis. [5]
- Find the volume of the solid formed when the area enclosed by the curve $y = x^2 + 3$ and the lines x = 1 and x = 2 performs one revolution about the x-axis. [5]
- A metal cube expands uniformly as it is heated. At time t seconds, the length of each edge of the cube is x cm, and the volume of the cube is V cm³ and $\frac{dV}{dx} = 3x^2$. The volume, V cm³, increases at a constant rate of 0.048 cm³ s⁻¹

Find
$$\frac{dx}{dt}$$
 when $x = 8$.

[5]

If the radius of a spherical bubble increases from 1 cm to 1.02 cm, find (to 2 significant figures) the approximate increase in the volume of the bubble.





CHAPTER SUMMARY: CHAPTER 10

You should know, and be able to use, these standard integral formulae.

| FUNCTION | INTEGRAL | | |
|-----------------|-----------------------------------|--|--|
| x^n | $\frac{1}{n+1}x^{n-1}, n \neq -1$ | | |
| $\sin ax$ | $-\frac{1}{a}\cos ax + c$ | | |
| $\cos ax$ | $\frac{1}{a}\sin ax + c$ | | |
| e ^{ax} | $\frac{1}{a}e^{ax} + c$ | | |

- You should also know how to calculate areas and volumes:
 - Area between a curve and x-axis = $\int_a^b y \, dx$, $y \ge 0$
 - Area between a curve and y-axis = $\int_{c}^{d} x \, dy$, $x \ge 0$
- Volume of revolution about the x-axis = $\pi \int_a^b y^2 \mathrm{d}x$
- Volume of revolution about the y-axis = $\pi \int_{\pi}^{d} x^2 dy$

CHAPTER 11



11 TRIGONOMETRY



Hummingbirds are tiny, colourful birds with beautiful feathers. They are admired for the sound of their wings, which flap up to 80 times per second, and for their song.

Sound is vibrations. These can by analysed as a combination of sine waves (sine is a trigonometric function – you can see a sine wave on page 191): any sound can be created with the right combination. When physicists or engineers investigate sound, they are actually investigating sine waves. Without their mathematical understanding of these waves, their analysis of sound would be very limited.

LEARNING OBJECTIVES

- Measure angles in radians
- Calculate arc length in a circle using radians
- Calculate the area of a sector using radians
- Understand the basic trigonometrical ratios and sine, cosine and tangent graphs
- Use sine and cosine rules

- Use sine and cosine rules to solve problems in 3D
- Use trigonometry identities and formulae to solve problems
- Solve trigonometric equations
- Use trigonometric formulae to solve equations

Q1 HINT

This is a way of remembering how to compute the sine, cosine, and tangent of an angle. In other words, what are the standard formulae for sine, cosine and tangent?

STARTER ACTIVITIES

- Write down the meaning of SOHCAHTOA.
- An aeroplane climbs at an angle of 15 degrees with the ground. What is the distance it has travelled (to the nearest hundred metres) when it has attained an altitude of 400 metres?
- The angle of elevation to the top of a tree is 20°. How far away are you from the tree, if the tree's height is 15 m? Round your answer to 3 significant figures.

MEASURE ANGLES IN RADIANS


Radians are an alternative to degrees. Some calculations involving circles are easier when radians are used, as opposed to degrees.

KEY POINTS

- A radian is the angle subtended at the centre of a circle by an arc whose length is equal to that of the radius of the circle.
- 1 radian = $\frac{180^{\circ}}{\pi}$.

HINT

The symbol for radians is $^{\circ}$, so θ° means that θ is in radians. If there is no symbol with an angle you should assume that it is in radians, unless the context makes it clear that it is in degrees.



Since $180^{\circ} = \pi \text{ rad}$, $1^{\circ} = \frac{\pi}{180} \text{ rad}$

EXAMPLE 1

Convert these angles into degrees.

- **b** $\frac{4\pi}{15}^{c}$
- a $\frac{7\pi^{c}}{8}$
 - $=\frac{7}{8}\times 180$

Remember that π rad = 180°

 $= 157.5^{\circ}$

Check using your calculator

 $\frac{4\pi^{c}}{15}$ $=\frac{4}{15}\times180$ $= 48^{\circ}$

EXAMPLE 2

Convert these angles into radians.

- a 150°
- b 110°

HINT

Your calculator will give the decimal answer 1.91986 These answers, in terms of π , are exact.

a
$$150^{\circ} = 150 \times \frac{\pi}{180}^{\circ}$$

$$=\frac{5\pi}{6}^{c}$$

b
$$110^{\circ} = 110 \times \frac{\pi}{180}^{\circ}$$

$$=\frac{11}{18}\pi^{c}$$

EXERCISE 1

SKILLS REASONING Convert these angles (in radians) to degrees.

$$\frac{1}{10}\pi$$

b
$$\frac{1}{20}\pi$$
 c $\frac{5}{12}\pi$ **d** $\frac{1}{2}\pi$

h
$$\frac{7}{9}\pi$$

Use your calculator to convert these to degrees.

b 10

d 1.37°

$$\sqrt{2}^{c}$$

f 3.14°

a 4.49°

▶ Use your calculator to write down the value of these trigonometric functions.

a
$$\sin\left(\frac{1}{4}\right)^{c}$$

b $\tan \sqrt{2}^c$

c cos 0.56c

d
$$\cos(3.2)^c$$

e sin 3.2°

tan 3.2°

 \triangleright Convert these angles to radians, giving your answer as multiples of π .

b 10°

d 30°

f 120°

135°

h 240°

315°

k 330°

5 Use your calculator to convert these angles to radians, giving your answer to 3 significant figures.

b 75°

c 100°

e 235°


325°

CALCULATE ARC LENGTH AND THE AREA OF A CIRCLE USING RADIANS

The formula for the length of an arc is simpler in radians than in degrees.

KEY POINTS

• To find the arc length l of a circle use the formula $l=r\theta$, where r is the radius of the circle and θ is the angle, in radians, contained by the sector.



EXAMPLE 3


Show that the length of an arc is $l = r\theta$.

The circle has centre O and radius r. The arc AB has length l. The length of the arc is proportional to the angle around O.

So
$$\frac{l}{2\pi r}=rac{ heta}{2\pi}$$
 which means $l=r heta$

HINT

Length of arc Angle AOB $\frac{\text{Circumference}}{\text{Circumference}} = \frac{\text{Figure AOB}}{\text{Total angle around } 0}, \text{ with both angles in radians}$




EXAMPLE 4

Find the length of the arc of a circle of radius 5.2 cm, given that the arc subtends an angle of 0.8 c at the centre of the circle.

Arc length =
$$5.2 \times 0.8$$
 Use $l = r\theta$ with $r = 5.2$ and $\theta = 0.8$ = 4.16 cm

EXAMPLE 5

An arc AB of a circle, with centre O and radius r cm, subtends an angle of θ radians at O. The **perimeter** of the sector AOB is P cm. Express r in terms of P and θ .



Draw a diagram to display the data


HINT

The perimeter is given by the arc AB, by OA and by OB. The length of the arc AB is $r\theta$.


$$P = r\theta + 2r$$
$$= r(\theta + 2)$$
$$r = \frac{P}{r}$$

EXAMPLE 6

The border of a garden pond consists of a straight edge AB of length 2.4 m, and a curved part C, as shown in the diagram below. The curved part is an arc of a circle, centre θ and radius 2 m. Find the length of C.



C subtends the reflex angle θ at O, so the length of Cis 2θ measured in metres, because the radius is 2m



You can use the isosceles triangle *AOB* to find the angle AOB inside the triangle

(Use your calculator in radian mode)

$$\sin x = \frac{1.2}{2}$$
, so $x = 0.6435^{\circ}$

The acute angle AOB is 2x rad = 2(0.6435)

$$= 1.287^{\circ}$$


So
$$\theta = (2\pi - 1.287) \text{ rad} = 4.996^{\circ}$$

So the length of C is $2\theta = 9.99 \,\mathrm{m}$ (3 s.f.)

Use θ + the acute angle AOB is $2\pi^{c}$


The formula for the area of a sector is also simpler with radians.

• To find the area A of a sector of a circle use the formula $A = \frac{1}{2}r^2\theta$, where r is the radius of the circle and θ is the angle, in radians, contained by the sector.



EXAMPLE 7

Show that the area of the sector of a circle with radius r is $A = \frac{1}{2}r^2\theta$



The circle has centre O and radius r.

The sector POQ has area A.

The area of the sector is proportional to the angle around O, so

$$\frac{A}{\pi r^2} = \frac{\theta}{2\pi}$$

$$A = \frac{1}{r^2}$$


$$\frac{\text{Area of sector}}{\text{Area of circle}} = \frac{\text{Angle } POQ}{\text{Total angle around } O}$$

$$A = \frac{1}{2}r^2\theta$$

Rearrange

EXAMPLE 8

A plot of land is in the shape of a sector of a circle of radius 55 m. The length of fencing that is erected along the edge of the plot to enclose the land is 176 m. Calculate the area of the plot of land.



Draw a diagram to include all the data and let the angle of the sector be θ .

$$Arc AB = 176 - (55 + 55) = 66 m$$

As the perimeter is given, first find length of arc AB

$$66 = 55\theta$$

Use formula for arc length, $l = r\theta$, where θ is in radians

So
$$\theta = 1.2 \text{ rad.}$$

Area of plot =
$$\frac{1}{2}$$
(55)²(1.2)
= 1815 m²

Use the formula for area of a sector, $A = \frac{1}{2}r^2\theta$



SKILLS DECISION MAKING

- An arc AB of a circle, centre O and radius r cm, subtends an angle θ radians at O. The length of AB is L cm.
 - a Find L when

$$r = 8, \theta = 0.45$$

i
$$r = 8, \ \theta = 0.45$$
 ii $r = 5.5, \ \theta = \frac{3}{2}\pi$ iii $r = 22, \ \theta = 0.7$

iii
$$r = 22, \theta = 0.7$$

b Find *r* when

i
$$L = 10, \theta = 0.5$$

ii
$$L = 2.34, \theta = 0.7$$

i
$$L = 10, \ \theta = 0.5$$
 ii $L = 2.34, \ \theta = 0.7$ iii $L = 2.5, \ \theta = \frac{5}{12}\pi$

 \bullet Find θ when

$$L = 10, r = 8.4$$

$$L = 5.5, r = 5.625$$

ii
$$L = 5.5, r = 5.625$$
 iii $L = \sqrt{12}, r = \sqrt{2}$

d Find the area of the sector contained by angle θ when these statements are true.


i
$$r = 8, \theta = 0.43$$


ii
$$r = 3.5, \ \theta = \frac{3\pi}{2}$$

iii
$$r = 22, \theta = 0.7$$


Find the area of the sector and the arc length of these (angles given in radians).













Find the area the shaded area (angles given in radians).




b




CHAPTER 11


4 Find the area of the shaded region in this diagram.




- 5 The area of sector *OAB* is 250 cm². The radius is 11 cm.
 - a Find θ .
 - b Find the length of the arc.




- 6 A sector of a circle has radius 6 cm and angle 2.2°. Calculate the area and perimeter.
- 7 Referring to the diagram, find
 - a the perimeter of the shaded region
 - b the area of the shaded region.




- 8 In the diagram, AB is the diameter of a circle with centre O and radius 4 cm. The point C is on the circle such that $\angle COB$ is $\frac{2}{3}\pi$. Find
 - a the perimeter of the shaded region
 - b the area of the shaded region.



- The arc AB of a circle with centre O and radius r is such that, $\angle AOB = 0.25^{\circ}$. Given that the perimeter of the minor sector AOB is 45 cm,
 - a calculate r
 - b show that the area of the minor segment AOB is 50 cm²
 - f c find the area of the segment enclosed by the chord AB and the minor arc AB.
- In the diagram, BC is the arc of a circle with centre O and radius 7 cm. The angle subtended by the arc is 1.9°.
 The points A and D are such that OA = OD = 3cm.
 Calculate the area of the shaded region.





- In the diagram, *OAB* is a sector of a circle with centre *O* and radius R cm. $\angle AOB = 2\theta$ radians. A circle with centre C and radius r cm touches the arc AB at T, and touches OA and OB at D and E respectively, as shown.
 - a Write down, in terms of R and r, the length of OC.
 - **b** Using $\angle OCE$, show that $R \sin \theta = r(1 + \sin \theta)$.
 - c Given that $\sin \theta = \frac{3}{4}$ and that the perimeter of the sector OAB is 21 cm, find R, giving your answer to 3 significant figures.






b Calculate the area of the **flowerbed**.






13 There is a straight path of length 70 m from the point A to the point B. The points are joined by a railway track in the form of an arc of the circle whose centre is C and whose radius is 44 m. as shown in the diagram.





- i the length of the railway track
- ii the shortest distance from C to the path
- iii the area of the region bounded by the railway track and the path.



UNDERSTAND THE BASIC TRIGONOMETRICAL RATIOS AND SINE, COSINE AND TANGENT GRAPHS

You need to memorise the exact values of certain trigonometrical ratios and you need to be able to find the exact values of some trigonometrical ratios.


You can find the trigonometrical ratios of angles 30°, 45° and 60° exactly. Consider an equilateral triangle ABC of side 2 units. If you drop a perpendicular from A to meet BC at D, then BD = DC = 1unit, the angle BAD is 30°, and the angle ABD is 60°.

Using Pythagoras' theorem in the triangle ABD

$$AD^2 = 2^2 - 1^2 = 3$$

So $AD = \sqrt{3}$ units.

Using the triangle
$$ABD$$
, $\sin 30^\circ = \frac{1}{2}$, $\cos 30^\circ = \frac{\sqrt{3}}{2}$, $\tan 30^\circ = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$ and $\sin 60^\circ = \frac{\sqrt{3}}{2}$, $\cos 60^\circ = \frac{1}{2}$, $\tan 60^\circ = \sqrt{3}$



If you now consider an isosceles right-angled triangle PQR, in which PQ=QR=1 unit, then the ratios for 45° can be found.

Using Pythagoras' theorem

$$PR^2 = 1^2 + 1^2 = 2$$

So
$$PR = \sqrt{2}$$
 units.


Then
$$\sin 45^\circ = \cos 45^\circ = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$
 and $\tan 45^\circ = 1$.

 $\sqrt{2}$ R 1

HINT

The graph of $\sin \theta$, where θ is in radians, has period 2π

You need to be able to recognise the sine graph.



Functions that repeat themselves after a certain interval are called periodic functions, and the interval is called the **period** of the function. You can see that $\sin \theta$ is periodic with a period of 360°.

HINT

Because it is periodic There are many symmetry properties of $\sin \theta$ but you can see from the graph that


$$\sin(\theta + 360^{\circ}) = \sin\theta$$
 and $\sin(\theta - 360^{\circ}) = \sin\theta$

Two further symmetry properties of $\cos \theta$ are

 $cos(\theta + 360^{\circ}) = cos \theta$ and $cos(\theta - 360^{\circ}) = cos \theta$

$$\sin(90^{\circ} - \theta) = \sin(90^{\circ} + \theta)$$

You need to be able to recognise the cosine graph.



Like $\sin \theta$, $\cos \theta$ is periodic with a period of 360°. In fact, the graph of $\cos \theta$ is the same as that of $\sin \theta$ when it has been translated by 90°.

HINT

Because it is periodic

HINT

Symmetry about $\theta = 0^{\circ}$

 $\cos(-\theta) = \cos(\theta)$

You need to be able to recognise the tangent graph



The tan function behaves very differently from the sine and cosine functions but it is still periodic, it repeats itself in cycles of 180° so its period is 180°.

The period symmetry properties of $\tan \theta$ are

$$tan(\theta + 180^{\circ}) = tan \theta$$

$$\tan(\theta - 180^\circ) = \tan\theta$$

EXAMPLE 9

Find the exact value of

cos 405°

b tan 120°

c sin 300°

 $\cos 405 = \cos(360 + 45)$

You can see from the cosine graph that $\cos x$ has a period of 360°

$$= \cos 45$$

$$=\frac{1}{\sqrt{2}}$$

Use triangle *PRQ* above

b tan 120 = tan (180 - 60)

You can see from the tangent graph that tan x has a period of 180°

$$= \tan(-60)$$

$$= -tan 60$$

$$= -\sqrt{3}$$

Use triangle ABD above

sin 300 = sin (360 - 60)

You can see from the sine graph that $\sin x$ has a period of 360°

$$= \sin(-60)$$

$$= -\sin 60$$

$$=-\frac{\sqrt{3}}{2}$$

Use triangle ABD above

EXERCISE 3

Express these trigonometric ratios using either 30°, 45° or 60°. (Do not use a calculator.)

- $c \sin(-60^\circ)$ d sin 330°
- e tan 210°

- **SKILLS ANALYSIS**
- a sin 135° b cos 120°
- $h \cos(-150^{\circ})$


tan 135°

- g sin 405°
- cos 330°
- tan 300°

- k sin 420°
- sin(-240°)
- m tan (−120°)
- n cos 225°
- tan 300°


2 The curve shows the graph $y = \cos x$ in the interval $0 \le x \le 4\pi$ ($0 \le x \le 720^\circ$).

Write down in radians any points where the curve intersects the x-axis.



USE SINE AND COSINE RULES

In your IGCSE Mathematics course you should have met the **sine rule**, the **cosine rule** and the formula for the **area of a triangle**.



The formulae for these are

Sine Rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ or $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$

= 6.22

Cosine Rule: $a^2 = b^2 + c^2 - 2bc \cos A$


Area of a triangle: $\frac{1}{2}ab\sin C$

HINT

Note that side a is opposite angle A

EXAMPLE 10

In the triangle ABC, AB = 8 cm, the angle BAC is 30°, and the angle BCA is 40°. Find BC.




Always draw a diagram and carefully then add the data

Using the sine rule
$$\frac{\alpha}{\sin A} = \frac{c}{\sin C}$$
 gives $\frac{x}{\sin 30} = \frac{8}{\sin 40}$ so $x = \frac{8 \sin 30}{\sin 40}$

Substitute $c=8\,\mathrm{cm},\,C=40^\circ,\,A=30^\circ,\,a=x$

EXAMPLE 11

In the triangle ABC, AB = 4 cm, AC = 12 cm, and the angle ABC is 64° . Find the angle ACB.



Using the sine rule
$$\frac{\sin C}{c} = \frac{\sin B}{b}$$

gives
$$\frac{\sin C}{4} = \frac{\sin 64}{12}$$


$$\sin C = \frac{4\sin 64}{12}$$
$$= \frac{\sin 64}{3}$$

and
$$C = \sin^{-1} \left(\frac{\sin 64}{3} \right)$$
$$= 17.4^{\circ}$$

EXAMPLE 12

In the triangle ABC, AB = 4 cm, AC = 3 cm, and the angle ABC is 44° . Work out two possible values for the angle ACB.

Substitute $b = 12 \,\mathrm{cm}, c = 4 \,\mathrm{cm}, B = 64^{\circ}$



Here $b = 3 \, \text{cm}, c = 4 \, \text{cm}, B = 44^{\circ}.$

Here C > B, as c > b, and so there will be two possible results. The diagram shows why.

With $B=44^{\circ}$ and c=4 cm drawn, imagine putting a pair of compasses at A, then drawing an arc with centre A and radius 3 cm. This will intersect BC at C_1 and C_2 showing that there are two triangles ABC_1 and ABC_2 where b = 3 cm, c = 4 cm and $B = 44^\circ$.

This would not happen if AC > 4 cm

 $\frac{3}{\sin 44} = \frac{4}{\sin C}$ Using the sine rule $\sin C = \frac{4\sin 44}{3}$ $C = \sin^{-1}\left(\frac{4\sin 44}{3}\right)$ gives One solution is


This corresponds to the triangle ABC_2 .

The other solution is $C = 180^{\circ} - 67.9^{\circ}$ As $\sin(180^{\circ} - x) = \sin x$ $= 112.1^{\circ}$

This corresponds to the triangle ABC_1 .

EXAMPLE 13

In the triangle PQR, PQ = 5.9 cm, QR = 8.2 cm and PR = 10.6 cm. Find the size of the angle PQR.



Using the cosine rule
$$\cos Q = \frac{p^2 + r^2 - q^2}{2pr}$$

gives
$$\cos Q = \frac{8.2^2 + 5.9^2 - 10.6^2}{2 \times 8.2 \times 5.9}$$
$$= -0.1065...$$


so
$$Q = \cos^{-1}(-0.1065...)$$

= 96.1°

Substitute $p=8.2\,\mathrm{cm},\,r=5.9\,\mathrm{cm}$ and $q=10.6\,\mathrm{cm}$

EXAMPLE 14

In the triangle ABC, AB = 5 cm, BC = 6 cm, and the angle ABC is x° . The area of the triangle ABC is 12 cm². Given that AC is the longest side, find the value of x.



Using Area =
$$\frac{1}{2}ac\sin B$$

gives
$$12 = \frac{1}{2} \times 6 \times 5 \times \sin x$$

Substitute $a=6\,\mathrm{cm},\,c=5\,\mathrm{cm},B=x^\circ$ and the area = $12\,\mathrm{cm}^2$

so
$$\sin x = 0.8$$

There are two possible solutions for x.

One is $x = \sin^{-1}(0.8) = 53.1^{\circ}$. However this cannot be the solution because B must be the largest angle, and if both A and C are smaller than 53.1° , the sum of the angles of the triangle cannot be 180° .


The other solution for x is $180^{\circ} - 53.1^{\circ} = 126.9^{\circ}$. This is the correct solution.


It can be written as 127° to 3 significant figures.


SKILLS


DECISION MAKING PROBLEM SOLVING


In each triangle below find the values of x, y and z.


























Find the area of each triangle.







- In triangle ABC, AB = 11 cm, BC = 9 cm and CA = 10 cm. Find the area of the triangle.
- A bird leaves a point A and flies 3.2 km, on a bearing of 128°, to a point B. It then flies 4.7 km, on a bearing of 066° , to point C. Show that angle ACB is 24.5° .
- **5** \triangleright ABCD is a quadrilateral. AB = 7 cm, AD = 6 cm and BC = 9 cm. Angle ABC = 75° and angle $ADC = 90^{\circ}$. Calculate the perimeter of ABCD.
- Prove the sine rule for a general triangle ABC.
- Prove the cosine rule for a general triangle ABC.
- The longest side of a triangle has length (2x-1) cm. The other sides have lengths (x-1) cm and (x + 1) cm. Given that the largest angle is 120°, find
 - the value of x.

the area of the triangle.

6cm

USE SINE AND COSINE RULES TO SOLVE PROBLEMS IN 3D


You need to be able to solve problems in 3D.

EXAMPLE 15

HINT

AC was calculated in part α

The diagram below shows a cuboid with dimensions 8 cm, 6 cm and 5 cm.



- Find length AC.
- **b** Find length AG.
- Find angle GAC.

8 cm

a AC is the diagonal across the base ABCD of the cuboid. Start by sketching this base and draw AC.

Identify a relevant right-angled triangle (triangle ABC in this case). Use Pythagoras' theorem to find AC.

tify a relevant right-angled triangle (triangle
$$ABC$$
 in this case). Pythagoras' theorem to find AC .
$$AC^2 = AB^2 + BC^2$$

$$AC^2 = 6^2 + 8^2$$


$$AC^2 = AB^2 + BC^2$$

$$AC^2 = 6^2 + 8^2$$

$$AC^2 = 100$$

So
$$AC = 10 \text{ cm}$$

b Identify a relevant right-angled triangle. Use triangle AGC (you use this triangle because C is vertically below G).



Use Pythagoras' theorem to find AG.


$$AG^2 = AC^2 + CG^2$$

$$AG^2 = 10^2 + 5^2$$

$$AG^2 = 125$$

So
$$AG = 11.2 \,\text{cm} (3 \,\text{s.f.})$$

• Find angle GAC. This is the triangle drawn in part b. Use trigonometry to find angle GAC.




$$\tan \theta = \frac{5}{10}$$
. therefore, theta = 26.6 degrees.

EXERCISE 5

SKILLS


DECISION MAKING PROBLEM SOLVING

The diagram below shows a square-based pyramid. The length AB is 8 cm. Point E is vertically above point X (not marked), the centre point of square ABCD. The height of the pyramid, EX, is 7 cm.




- Calculate length AC.
- Calculate angle EAC.


- Calculate length AE.
- Calculate the area of face AEB.
- A cube has sides of 6 cm. Find
 - the length AB
 - **b** the angle between AB and the horizontal plane.




- The diagram represents a cuboid ABCDEFGH. $CD = 10 \, \text{cm}$, $BC = 8 \, \text{cm}$ and $BF = 4 \, \text{cm}$.
 - a Calculate the length of AG. Give your answer to 3 significant figures.
 - **b** Calculate the size of the angle between AG and the face ABCD. Give your answer to 3 significant figures.



The diagram represents a ski slope. AEFD is a rectangle. ABCD is a square. EF and FC are perpendicular to plane ABCD. AB = AD = 60 m. Angle ABE = 90°. Angle $BAE = 30^{\circ}$. Calculate the size of the angle that the line DE makes with the plane ABCD.



- **5** \triangleright *ABCDEFGH* is a cuboid. *M* is the midpoint of *HG*. *N* is the midpoint of *DC*.
 - Calculate BN.
 - **b** Work out the size of the **obtuse** angle between the planes *MNB* and *CDHG*.



- 6 ► A mobile cell tower 150 m tall has two support cables running 300 metres due east and x metres due south, anchored at A and B. The angle of inclination to the horizontal of the latter cable is 50°
 - **a** Find the distance between the cable attached to *B* and the horizontal plane.
 - **b** Find the distance between A and B.

USE TRIGONOMETRY IDENTITIES TO SOLVE PROBLEMS

You should already know the identity

$$\tan\theta \equiv \frac{\sin\theta}{\cos\theta}$$

You also need to know the following. The first of these is not given in the exam.

$$\sin^2\theta + \cos^2\theta = 1$$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

EXAMPLE 16

Simplify these expressions.

a
$$\sin^2 3\theta + \cos^2 3\theta$$

b
$$5 - 5\sin^2\theta$$

$$\frac{\sin 2\theta}{\sqrt{1-\sin^2 2\theta}}$$

a
$$\sin^2 3\theta + \cos^2 3\theta = 1$$

This is the same as $\sin^2 \theta + \cos \theta = 1$ with θ replaced by 3θ

b
$$5 - 5\sin^2\theta = 5(1 - \sin^2\theta)$$

$$=5\cos^2\theta$$

Use
$$\sin^2 \theta + \cos^2 \theta = 1$$
 which gives $1 - \sin^2 \theta = \cos^2 \theta$

$$\frac{\sin 2\theta}{\sqrt{1 - \sin^2 2\theta}} = \frac{\sin 2\theta}{\sqrt{\cos^2 2\theta}}$$

Use
$$1 - \sin^2 2\theta = \cos^2 2\theta$$

$$\sqrt{1 - \sin^2 2\theta} \qquad \sqrt{\cos^2 2\theta}$$
$$= \frac{\sin 2\theta}{|\cos 2\theta|}$$

Since
$$\sqrt{\cos^2 2\theta} = |\cos 2\theta|$$

$$= \tan 2\theta$$
 or $-\tan 2\theta$

as
$$|\cos 2\theta| = \pm \cos 2\theta$$

EXAMPLE 17

Show that
$$\frac{\cos^4 \theta - \sin^4 \theta}{\cos^2 \theta} = 1 - \tan^2 \theta$$

When you have to prove an identity like this, you often need to use basic identities you know like $\sin^2 \theta + \cos^2 \theta = 1$

Usually the best strategy is to start with the more complicated side (here the left-hand side, LHS) and try to produce the expression on the other side.

LHS =
$$\frac{\cos^{4}\theta - \sin^{4}\theta}{\cos^{2}\theta}$$
=
$$\frac{(\cos^{2}\theta - \sin^{2}\theta)(\cos^{2}\theta + \sin^{2}\theta)}{\cos^{2}\theta}$$
=
$$\frac{\cos^{2}\theta - \sin^{2}\theta}{\cos^{2}\theta}$$
=
$$\frac{\cos^{2}\theta}{\cos^{2}\theta} - \frac{\sin^{2}\theta}{\cos^{2}\theta}$$
=
$$1 - \tan^{2}\theta$$

Factorise the numerator as the difference of two squares

Use that
$$\cos^2 \theta + \sin^2 \theta = 1$$

EXAMPLE 18

Use the formula for $\sin(A + B)$ and $\cos(A + B)$ to prove that

$$a \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

= RHS

a
$$tan(A+B) = \frac{tanA + tanB}{1 - tanA tanB}$$
 b $tan(A-B) = \frac{tanA - tanB}{1 - tanA tanB}$

a
$$\tan(A+B) = \frac{\sin(A+B)}{\cos(A+B)} = \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B}$$

Dividing the top and bottom by $\cos A \cos B$ gives

$$\tan(A + B) = \frac{\frac{\sin A \cos B}{\cos A \cos B} + \frac{\cos A \sin B}{\cos A \cos B}}{\frac{\cos A \cos B}{\cos A \cos B} - \frac{\sin A \sin B}{\cos A \cos B}}$$
$$= \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

Cancel terms as shown and use the result $\tan \theta = \frac{\sin \theta}{}$

b Replace B by -B in the result above.

$$\tan(A - B) = \frac{\tan A + \tan(-B)}{1 - \tan A \tan(-B)}$$
$$= \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

Use the result $tan(-\theta) = -tan \theta$

EXAMPLE 19

Show, using the formula for $\sin(A - B)$, that $\sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$

You know the exact form of sin and cos for many angles, for example 30°, 45°, 60°, 90°, 180. You can use two of these to write 15°.

$$sin 15^{\circ} = sin (45^{\circ} - 30^{\circ}) = sin 45^{\circ} cos 30^{\circ} - cos 45^{\circ} sin 30^{\circ}
= \frac{\sqrt{2}}{2} \times \frac{\sqrt{3}}{2} - \frac{\sqrt{2}}{2} \times \frac{1}{2}
= \frac{1}{4} (\sqrt{2} \times \sqrt{3} - \sqrt{2})
= \frac{1}{4} (\sqrt{6} - \sqrt{2})$$

EXAMPLE 20

Given that $2\sin(x+y) = 3\cos(x-y)$, express $\tan x$ in terms of $\tan y$.

 $2\sin x \cos y + 2\cos x \sin y = 3\cos x \cos y + 3\sin x \sin y$

Expand $\sin(x + y)$ and $\cos(x - y)$

 $\frac{2\sin x\cos y}{\cos x\cos y} + \frac{2\cos x\sin y}{\cos x\cos y} = \frac{3\cos x\cos y}{\cos x\cos y} + \frac{3\sin x\sin y}{\cos x\cos y}$

Divide by $\cos x \cos y$

 $2\tan x + 2\tan y = 3 + 3\tan x \tan y$

Use
$$\frac{\cos x}{\cos x} = \frac{\cos y}{\cos y} = 1$$
, $\frac{\sin x}{\cos x} = \tan x$ and $\frac{\sin y}{\cos y} = \tan y$

so $\tan x(2-3\tan y)=3-2\tan y$

and
$$\tan x = \frac{3 - 2 \tan y}{2 - 3 \tan y}$$

EXERCISE 6

SKILLS

DECISION

MAKING

ANALYSIS

1 > Simplify each of these expressions.

a $1 - \sin^2 \theta$

b $1 - \cos^2 \frac{1}{2} \theta$

c $6\sin^2 4\theta + 6\cos^2 4\theta$

d $\cos^2 A - 1$

 $\frac{\sin 4\theta}{\tan 4\theta}$

f $\frac{\sqrt{1-\cos^2\theta}}{\cos\theta}$

$$\frac{\sqrt{1 - \cos^2 3A}}{\sqrt{1 - \sin^2 3A}}$$

h
$$(1 + \cos x)^2 + (1 - \cos x)^2 + 2\sin^2 x$$

$$\sin^4 x + \sin^2 x \cos^2 x$$

$$\sin^4 x + 2\sin^2 x \cos^2 x + \cos^4 x$$

2 Prove that

$$a (\cos x + \sin x)^2 \equiv 1 + 2\sin x \cos x$$

b
$$\frac{1}{\cos \theta} - \cos \theta \equiv \sin \theta \tan \theta$$

$$\frac{\cos^2\theta}{1-\sin\theta}\equiv 1+\sin\theta$$

$$\frac{1 + \sin x}{\cos x} = \frac{\cos x}{1 - \sin x}$$

$$e \tan x + \frac{1}{\tan x} \equiv \frac{1}{\sin x \cos x}$$

$$\cos^2 A - \sin^2 A \equiv 2\cos^2 A - 1 \equiv 1 - 2\sin^2 A$$

$$g 2 - (\sin x - \cos x)^2 \equiv (\sin x + \cos x)^2$$

h
$$\sin^2 x \cos^2 y - \cos^2 x \sin^2 y \equiv \sin^2 x - \sin^2 y$$

3 Express these as a single sine, cosine or tangent.

$$a \sin 15\cos 20 + \cos 15\sin 20$$

$$\frac{\tan 76 - \tan 45}{1 + \tan 76 \tan 45}$$

d
$$\cos 2x \cos x - \sin 2x \sin x$$

e
$$\cos 4\theta \cos 3\theta - \sin 4\theta \sin 3\theta$$

$$\frac{\tan 2x + \tan 3x}{1 - \tan 2x \tan 3x}$$

$$g \sin(A + B)\cos B - \cos(A + B)\sin B$$

h
$$\cos\left(\frac{3x+2y}{2}\right)\cos\left(\frac{3x-2y}{2}\right)-\sin\left(\frac{3x+2y}{2}\right)\sin\left(\frac{3x-2y}{2}\right)$$

4 Calculate, without using your calculator, the exact value of

$$a \sin 30\cos 60 + \cos 30\sin 60$$

$$\frac{\tan 45 + \tan 15}{1 - \tan 45 \tan 15}$$

e
$$\frac{1 - \tan 15}{1 + \tan 15}$$

5 Prove these identities.

$$\sin(A + 60) + \sin(A - 60) \equiv \sin A$$

$$\frac{\sin(x+y)}{\cos x \cos y} \equiv \tan x + \tan y$$

$$\mathbf{c} \quad \cos\left(\theta + \frac{\pi}{3}\right) + \sqrt{3}\sin\theta \equiv \sin\left(\theta + \frac{\pi}{6}\right)$$

SOLVE TRIGONOMETRIC EQUATIONS

You can use your calculator to find a solution to an equation such as $\sin x = 0.4$. Here, the solution is x = 23.6

However, by analysing the sine, cosine and tangent graphs, you can see other possible solutions.

Here, the solutions are

$$x = 180 - 23.6 = 156.4$$

$$x = 23.6 + 360 = 383.6$$

and so on.

The below table summarises this situation for sine, cosine and tangent.

| EQUATION | 1st SOLUTION (CALC) | 2nd SOLUTION | 3rd SOLUTION | 4th SOLUTION |
|--------------|---------------------|--------------|---------------|---------------|
| $\sin x = k$ | а | 180 - a | All cases are | All cases are |
| $\cos x = k$ | а | 360 - a | ±360 | 2nd solution |
| tan x = k | а | 180 + a | | ±360 |

EXAMPLE 21

Solve, in the interval $0 \le x < 360$, giving your answers to the nearest degree.

a
$$\sin x = 0.5$$

b
$$4\sin x = -3$$

c
$$2\tan(x+1)=0$$

a
$$\sin x = 0.5$$

$$x = 30^{\circ}$$

Using a formula table or a calculator

or

$$x = 180 - 30$$

= 150°

b
$$4\sin x = -3$$

$$\sin x = -\frac{3}{4}$$

Rearrange into the form $\sin x = k$.

$$x = -48.6^{\circ}$$

This is outside the range so you need to find a second and third solution.

$$x = 180 - (-48.6)$$

Use the formula table to calculate the correct range of answers

$$= 228.6^{\circ}$$

or

$$x = -48.6 + 360 = 311.4^{\circ}$$
 (to 3 s.f.)

Therefore, $x = 229^{\circ}$ or 311°

c
$$2 \tan x + 1 = 0$$

$$\tan x = -\frac{1}{2}$$

Rearrange into the form $\tan x = k$

$$x = -26.6$$

Use the formula table to calculate the correct range of answers

or

$$x = -26.6 + 180$$

$$= 153.4$$

$$x = -26.6 + 360$$

$$= 333.4$$

so
$$x = 155^{\circ}$$
 or $x = 333^{\circ}$ (to 3 s.f.)

EXAMPLE 22

Solve, in the interval $0 \le x < 360$, giving your answers to the nearest degree.

$$\cos(x-25)=0.4$$

b
$$tan 2x = 3$$

$$\cos(x-25)=0.4$$

Let
$$y = x - 25$$

Then
$$\cos y = 0.4$$

Substitute y = x - 25 into the equation

$$y = 66.4$$

Solve for $\cos y = 0.4$

or

$$y = 360 - 66.4$$

$$= 293.6$$

Therefore,

$$x - 25 = 66.4$$
 or 293.6

$$x = 91.4^{\circ} \text{ or } 319^{\circ}$$

Then use x = y + 25 to find the x values to the nearest degree

b tan 2x = 3

$$y = 2x$$

Substitute y = 2x into the equation

$$tan y = 3$$

$$y = 71.6$$
 or 251.6 or 431.6 or 611.6

Solve for tan y = 3

Therefore

$$x = 35.8^{\circ} \text{ or } 126^{\circ} \text{ or } 216^{\circ} \text{ or } 306^{\circ}$$

Then use $x = \frac{y}{2}$ to find the x values to the nearest degree

You may also need to solve equations in radians.

Solve $2\cos 2x + 1 = 0$ for $-\pi \le x \le \pi$. Give your answers in multiples of π .

Solve $2\cos 2x + 1 = 0$ for $-\pi \le x \le \pi$. Give your answers in multiples of π .

$$2\cos 2x + 1 = 0$$

First rearrange into the form $\cos 2x = k$

Using a calculator gives $\frac{2\pi}{3}$

Let
$$y = 2x$$

$$\cos y = -\frac{1}{2}$$

$$y=\frac{2\pi}{3}$$

$$x = \frac{\pi}{3}$$
 Since $y = 2x$ divide by 2 to find x . This is the first solution

or

$$y = 2\pi - \frac{2\pi}{3}$$

$$=\frac{4\pi}{3}$$
$$x=\frac{2\pi}{3}$$

$$y = \frac{3\pi}{3} - 2\pi$$

$$x = -\frac{3}{2\pi}$$

Use the formula table for the second solution

Since y = 2x divide by 2 to find x. This is the second solution

Use the formula table for the third solution

Since y = 2x divide by 2 to find x. This is the third solution

or

$$y = \left(2\pi - \frac{2\pi}{3}\right) - 2\pi$$
$$= -\frac{2\pi}{3}$$

Use the formula table for the fourth solution

 $x = -\frac{\pi}{3}$

Since y = 2x divide by 2 to find x. This is the third solution

(Use the table to find all the values in the acceptable range)

So the solutions in the range $-\pi \le x \le \pi$ are

$$x = \frac{\pi}{3}, \frac{2\pi}{3}, -\frac{2\pi}{3}, -\frac{\pi}{3}$$

EXAMPLE 24

Solve, in the interval given, giving your answer to 3 significant figures.

a
$$\sin\left(x-\frac{\pi}{2}\right)=\frac{3}{4}$$

$$0 \le x \le 2\pi$$

b
$$\sin^2\left(x + \frac{\pi}{6}\right) = \frac{1}{2}$$

$$-\pi \leq x \leq \pi$$

Solve for $y = \frac{3}{4}$

 $a \sin\left(x - \frac{\pi}{2}\right) = \frac{3}{4}$

Let
$$y = x - \frac{\pi}{2}$$

Then
$$\sin y = \frac{3}{4}$$

$$y = 0.848 \text{ rad}$$

$$y = 2.29 \text{ rad}$$

Therefore,

$$x - \frac{\pi}{2} = 0.848$$
 or 2.29

$$x = 2.42 \text{ or } 3.86$$

205

Let
$$y = x + \frac{\pi}{6}$$

Then
$$\sin^2 y = \frac{1}{2}$$

$$\sin y = \pm \frac{\sqrt{2}}{2}$$

$$\left(\text{Solve for } \sin y = \frac{\sqrt{2}}{2}\right)$$

$$y = \frac{1}{4}\pi \text{ or } \frac{3}{4}\pi \text{ or } -\frac{3}{4}\pi \text{ or } -\frac{1}{4}\pi$$

 $y = \frac{1}{4}\pi \text{ or } \frac{3}{4}\pi \text{ or } -\frac{3}{4}\pi \text{ or } -\frac{1}{4}\pi$ (Use the table to find all the values in the acceptable range)

Therefore.

$$x = \frac{1}{12}\pi, \frac{7}{12}\pi, -\frac{11}{12}\pi, -\frac{5}{12}\pi$$

EXERCISE 7

SKILLS

Solve these equations for θ , in the interval $0 < \theta \le 360^{\circ}$.

a
$$\sin \theta = -\frac{1}{2}$$
 b $\tan \theta = \sqrt{3}$ **c** $\cos \theta = -1$ **d** $\sin \theta = \sin 15$ **e** $\cos \theta = -\cos 40$ **f** $2 \tan \theta = 2$ **g** $\cos \theta = 0$ **h** $\sin \theta = -0.766$

$$e \cos \theta = -\cos 40$$

$$2 \tan \theta = 2$$

$$\cos \theta = 0$$

h
$$\sin \theta = -0.766$$

$$7 \sin \theta = 5$$

$$j \quad 2\cos\theta = -\sqrt{2}$$

Solve these equations for x or θ . Give your answers to 2 decimal places or in terms of π where appropriate, in the intervals indicated.

a
$$\sin x = -\frac{\sqrt{3}}{2}, -180 \le x \le 180$$

b
$$2 \sin x = -0.3, -180 \le x \le 180$$

$$\cos x = -0.809, -180 \le x \le 180$$

d
$$\cos x = 0.84, -360 \le x \le 0$$

c
$$\cos x = -0.809, -180 \le x \le 180$$

e $\tan x = \frac{\sqrt{3}}{3}, 0 \le x \le 720$
f $\tan x = 2.90, 80 \le x \le 440$
g $\sin \theta = 0, -2\pi \le \theta \le 2\pi$
i $\cos \theta = -\frac{1}{2}, -2\pi \le \theta \le \pi$
j $2(1 + \tan \theta) = 1 - 5\tan \theta, -2\pi$

f
$$tan x = 2.90, 80 \le x \le 440$$

g
$$\sin \theta = 0, -2\pi \le \theta \le 2\pi$$

h
$$\sin \theta = \frac{1}{\sqrt{2}}, -2\pi \le \theta \le \pi$$

i
$$\cos \theta = -\frac{1}{2}, -2\pi \le \theta \le \pi$$

$$\int_{0}^{\sqrt{2}} 2(1 + \tan \theta) = 1 - 5\tan \theta, -\pi \le \theta \le 2\pi$$

3 Find the values of θ , in the interval $0 \le \theta \le 360$, for which

a
$$\sin 4\theta = 0$$

b
$$\cos 3\theta = -1$$

c
$$tan 2\theta = 1$$

d
$$\tan \frac{1}{2}\theta = -\frac{1}{\sqrt{3}}$$
 e $\sin(-\theta) = \frac{1}{\sqrt{2}}$ **f** $2\sin(\theta - 20) = 1$

$$e \sin(-\theta) = \frac{1}{\sqrt{2}}$$

f
$$2\sin(\theta - 20) = \frac{1}{2}$$

g
$$\tan(\theta + 75) = \sqrt{3}$$

 \triangleright Solve, in the intervals indicated, these equations for θ , where θ is measured in radians. Give your answer in terms of π or to 2 decimal places.

a
$$\sin \theta = 0, -2\pi \le \theta \le 2\pi$$

b
$$\sin \theta = \frac{\sin \theta}{\cos \theta}, \ 0 \le \theta \le 2\pi$$

5 Solve these equations in the interval given, giving your answers to 3 significant figures where appropriate.

a
$$\cos(3x+60) = \frac{1}{2}, -90 \le x \le 90$$
 b $\cos(x+\frac{\pi}{4}) = \frac{3}{4}, 0 \le x \le 2\pi$

b
$$\cos(x + \frac{\pi}{4}) = \frac{3}{4}, 0 \le x \le 2\pi$$

c
$$tan(4x - 30) = 2, 90 \le x \le 270$$

d
$$\sin(2x - \pi) = -\frac{1}{2}, -\pi \le x \le \pi$$

e
$$\cos^2(x - \frac{\pi}{16}) = \frac{1}{4}, -\pi \le x \le \pi$$

USE TRIGONOMETRIC FORMULAE TO SOLVE EQUATIONS

EXAMPLE 24

Solve these equations for $0 \le x \le 360^\circ$.

a
$$2\sin x = \tan x$$

a $2\sin x = \tan x$

b
$$\sin^2(x-30^\circ)=\frac{1}{2}$$

When factorising
$$\sin x(2\cos x - 1) = 0$$
, do not cancel. This would cause the solutions $x = 0^\circ$, 180° to be lost.

so
$$2\cos x \sin x = \sin x$$

and
$$\sin x(2\cos x - 1) = 0$$

Rearrange

Use that $\tan x = \frac{\sin x}{\cos x}$

$$\sin x = 0$$
, which gives $x = 0^{\circ}$ or 180° or 360°

or
$$2\cos x - 1 = 0$$

 $\cos x = \frac{1}{2}$, which gives $x = 60^{\circ}$ or 300°

The solutions are $x = 0^{\circ}$ or 60° or 180° or 300° in the range $0 \le x \le 360^{\circ}$

HINT

If one of the solutions was $x = 90^{\circ}$, it would have been rejected. The reason is that the equation was multiplied by a factor $\cos x$ because there was a $\cos x$ in the denominator on the rhs of $2\sin x = \frac{\sin x}{\cos x}$. However, $\cos 90^\circ = 0$, so this would have been incorrect. Take care when multiplying by trigonometric functions.

b
$$\sin^2(x - 30) = \frac{1}{2}$$

 $\sin(x - 30) = \pm \frac{1}{\sqrt{2}}$

Take the square root of both sides

$$\mathsf{Let}\ y = x - \mathsf{30}$$

So the solutions of $\sin y = \frac{1}{\sqrt{2}}$ in the interval $0 \le x \le 360^\circ$ are

$$y=45^{\circ}$$
 or 135°

Recall that
$$\sin 45 = \frac{1}{\sqrt{2}}$$

$$x - 30^{\circ} = 45^{\circ} \text{ or } 135^{\circ}$$

As
$$y = x - 30$$

$$x = 75^{\circ} \text{ or } 165^{\circ}$$

The solutions of $\sin y = -\frac{1}{\sqrt{2}}$ in the interval $0 \le x \le 360^\circ$ are

$$y = -45^{\circ} \text{ or } 180^{\circ} - (-45^{\circ}) \text{ or } -45^{\circ} + 360^{\circ}$$

$$x - 30^{\circ} = -45^{\circ} \text{ or } 225^{\circ} \text{ or } 315^{\circ}$$
 As $y = x - 30^{\circ}$

As
$$y = x - 30^{\circ}$$

$$x = -15^{\circ} \text{ or } 255^{\circ} \text{ or } 345^{\circ}$$

The solution $x = -15^{\circ}$ is not in the range $0 \le x \le 360^{\circ}$ so it is rejected.

The acceptable solutions are $x = 75^{\circ}$ or 165° or 255° or 345°

You may need to solve trigonometric equations in radians.

EXAMPLE 25

Find the values of x, in the interval $-\pi < x \le \pi$, that satisfy the equation

$$2\cos^2 x + 9\sin x = 3\sin^2 x$$

Give your answers in radians correct to 3 significant figures.

$$2\cos^2 x + 9\sin x = 3\sin^2 x$$

Use $\cos^2 x + \sin^2 x = 1$ to replace $\cos^2 x = 1 - \sin^2 x$

$$2(1 - \sin^2 x) + 9\sin x = 3\sin^2 x$$

Let $\sin x = s$

$$2 - 2s^2 + 9s = 3s^2$$

Substitute $s = \sin x$

$$5s^2 - 9s - 2 = 0$$

Rearrange

$$(5s + 1)(s - 2) = 0$$

Factorise

$$s = 2 \text{ or } s = -\frac{1}{5}$$

 $\sin x = 2$ has no solutions

$$\sin x = -\frac{1}{5}$$

gives
$$x = -0.201$$
 (3 s.f.)

Use a calculator in 'rad' mode to get this solution. You can find another solution noting that $\sin(\pi - x) = \sin x$ when x is in radians. This gives: $x = \pi - (-0.201) = 3.34$ (3 s.f.).

You can find further solutions considering $-0.201 \pm 2\pi$ and $3.34 \pm 2\pi$. You could also consider solutions like $-0.201 \pm 4\pi$, but it is clear that these would all be outside the interval $-\pi < x \le \pi$.

SKILLS

CRITICAL THINKING PROBLEM SOLVING ANALYSIS


- **1** Solve these equations for θ , in the interval $0 < \theta \le 360$.
 - a $\cos \theta = \sqrt{3} \sin \theta$
 - **b** $\sin \theta = -\cos \theta$
- **2** Solve $2\cos\theta 3\sin\theta = 0$, $0 < \theta \le 2\pi$ where θ is measured in radians.
- **3** Solve these equations for θ in the interval $0 < \theta \le 360$.
 - a $4\cos^2\theta = 1$
 - **b** $2\cos^2\theta = 5\cos\theta 2$
 - $\sin\theta + 2\cos^2\theta + 1 = 0$
 - d $4\cos^2\theta 5\sin\theta = 5$
- Solve these equations for x, in the interval $0 \le x \le 2\pi$. Give your answers to 3 significant figures or in the form $\frac{a}{b}\pi$, where a and b are integers.
 - a $2\sin^2(x + \frac{\pi}{3}) = 1$
 - **b** $6\sin^2 x + \cos x = 4$
 - $\cos^2 x 6\sin x = 5$

EXAM PRACTICE: CHAPTER 11

- The diagram below shows a sector of a circle of radius 6 cm with centre A. Given that the size of angle BAC is 0.95 radians, find
 - a the area of the sector ABC [2]
 - **b** the length of the arc BC.

The point D lies on the line AC and is such that AD = BD. The length AD is 5.16 cm.

- f c Find the perimeter of R.
- d Find the area of R. [2]



2.2rad
A
4cm

The shape BCD is shown above.


The straight lines DB and DC are equal in length. The curve BC is an arc of a circle with centre A and radius 6 cm. The size of $\angle BAC$ is 2.2 radians and AD = 4 cm.

- Show that the area of the sector BAC is $39.6 \,\mathrm{cm}^2$.
- **b** Calculate the size of $\angle DAC$, in radians. [3]
- c Calculate the total area of the shape. [4]
- In the triangle ABC, AB = 11 cm, BC = 7 cm and CA = 8 cm.
 - **a** Find the size of angle *C*, giving your answer to 3 significant figures. [2]
 - **b** Find the area of triangle ABC, giving your answer in cm² to 3 significant figures. [3]

- Show that the area of a triangle is $\frac{1}{2}ab\sin C$ [3]
- a Given that $\sin \theta = 5 \cos \theta$, find $\tan \theta$. [3]
 - **b** Hence, or otherwise, find the values of θ in the interval $0 \le \theta < 360^{\circ}$ for which $\sin \theta = 5 \cos \theta$. [2]
- Solve $2\cos 2x + 1 = 5\sin x$, for the interval $0 \le x < 2\pi$.
- Simplify
 - $\mathbf{a} \quad \sin^2 3\theta \sin^2 3\theta \cos^2 3\theta \tag{4}$
- a Given that $2(\sin x + 2\cos x) = \sin x + 5\cos x$, find the exact value of $\tan x$.
 - **b** Given that $\sin x \cos y + 3\cos x \sin y = 2\sin x \sin y 4\cos x \cos y$, express $\tan y$ in terms of $\tan x$.
- Find, giving your answers in terms of π , all values of θ in the interval $0 < \theta < 2\pi$ for which
 - **a** $\tan(\theta + \frac{\pi}{3}) = 1$ [3]
 - $\mathbf{b} \quad \sin 2\theta = -\frac{\sqrt{3}}{2} \tag{4}$
- Find the values of x in the interval $0 \le x \le 360^\circ$ for $2\sin x = \cos(x 60)$.
- Prove that $(\cos \theta \tan \theta)^2 + (\sin \theta + 1)^2 \equiv 2 + \tan^2 \theta$ [4]
- 12 a Sketch, for $0 \le x \le 2\pi$, the graph of $y = \sin\left(x + \frac{\pi}{6}\right)$.
 - b Write down the exact coordinates of the points where the graph meets the axes. [1]
- Solve $3\sin^2\theta 2\cos^2\theta = 1$ for the interval $0^\circ \le \theta < 360^\circ$. [4]
- Prove that $\frac{\cos A}{\sin B} \frac{\sin B}{\cos A} \equiv \frac{\cos (A+B)}{\sin B \cos B}$ [5]
- The region R is bounded by the curve $y = \sin 2x$, the x-axis and the vertical lines x = 0 and $x = \frac{\pi}{2}$. Find the volume of the solid formed when the region is rotated 2π radians about the x-axis.

CHAPTER SUMMARY: CHAPTER 11

- 1 radian = $\frac{180}{\pi}$ degrees.
- The length of an arc of a circle is $l = r\theta$
- The area of a sector is $A = \frac{1}{2}r^2\theta$




The sine rule is

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
or

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

The cosine rule is

$$a^2=b^2+c^2-2bc\cos A$$
 or $b^2=a^2+c^2-2ac\cos B$ or $c^2=a^2+b^2-2ab\cos C$



You can find an unknown angle using rearranged form of the cosine rule.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$
 or $\cos B = \frac{a^2 + c^2 - b^2}{2ac}$ or $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$

You can find the area of a triangle using the formula

Area =
$$\frac{1}{2}ab \sin C$$

if you know the length of two sides (a and b) and the value of the angle C between them.

You need to know these identities.

$$\blacksquare$$
 $\sin^2\theta + \cos^2\theta = 1$

$$\blacksquare$$
 $\sin(A + B) = \sin A \cos B + \cos A \sin B$

The table below will help you solve trigonometrical equations.

| EQUATION | 1st SOLUTION (CALC) | 2nd SOLUTION | 3rd SOLUTION | 4th SOLUTION |
|--------------|---------------------|----------------|---------------|---------------|
| $\sin x = k$ | а | 180 - a | All cases are | All cases are |
| $\cos x = k$ | а | 360 - a | ±360 | 2nd solution |
| $\tan x = k$ | а | 180 + <i>a</i> | | ±360 |

GLOSSARY

acceleration the rate at which the velocity of an object increases

adjacent next to one another

approximation 'approximate' means 'almost exact'

area of a triangle $\frac{1}{2}ab\sin C$

arithmetic sequence where the terms have a common difference

asymptote a line that a curve approaches, but never reaches

attribute if people attribute something to a person, they believe that person did it, said it, etc

base of a logarithm $y = \lg x$ means the same as $10^y = x$. Here the *base* of the logarithm is 10. Another common base is the constant e. In this case can write $\log_e x$ as $\ln x$

bearing an angle measured clockwise from north. Always expressed with three digits, e.g. 045°

beehive a container where people keep bees

binomial series means the same as 'Binomial Expansion'

calculus the part of mathematics that deals with changing quantities, such as the speed of a falling object or the slope of a curve

Cartesian describes the standard x-y graph and its axes

chime (of a clock) to sound a small bell

coefficient in the expression $3x^2 + 4x + 2$, 3 is the coefficient of x^2 and 4 is the coefficient of x

column matrix e.g. $\begin{pmatrix} x \\ y \end{pmatrix}$

common difference the difference between any two consecutive terms in an arithmetic sequence (or series), shared by (*common* to) the whole sequence. In the arithmetic sequence 2, 4, 6, 8, 10 ... the *common difference* is 2

common ratio the same as a common difference (above), except that multiplication rather than addition is used. In the geometric sequence 2, 4, 8, 16, 32 ... the *common ratio* is 2

cosine rule $a^2 = b^2 + c^2 - 2bc\cos A$

constant a term that does not include a variable. In the expression $3x^2 + 4x + 2$, 2 is a *constant*

cube a six-sided solid whose sides are all squares

cuboid a six-sided solid whose sides are all rectangles, placed at right angles

dam a wall built across a river or stream to stop the water from flowing, especially in order to make a lake or produce electricity

deduce to reach a logical conclusion. If x + 2 = 3, you deduce that x = 1

derivative, derived function the result of differentiating

determine to find out. By analysing the equation of a curve, you can *determine* whether it is a circle or not

displacement change of position

divisible able to be divided without remainder. $x^2 + 3x + 2 = (x + 1)(x + 2)$, hence $x^2 + 3x + 2$ is divisible by (x + 1) and (x + 2), but no other expression

due north/south/east/west directly to the north/south/
east/west

earnings the money that you receive for the work that you do

elongated longer than usual

equivalent having the same value or meaning as something else

expansion multiplying out terms in brackets. The opposite of **factorisation**

exponential function a function where the variable occurs as an exponent, e.g. e^{2x+1}

factor theorem if f(x) is a polynomial and f(p) = 0 then (x - p) is a factor of f(x)

factorial the result when you multiply an integer by all the integers below it, down to 1. It is written as an exclamation mark. $5! = 5 \times 4 \times 3 \times 2 \times 1$

factorise to rewrite an expression using brackets. You *factorise* $x^2 + 3x + 2$ to get (x + 1)(x + 2)

finite having an end

fluctuate if an amount *fluctuates*, it keeps changing both up and down

function x + 2, 2x, x^2 , $\ln x$, $\sin x$, e^x and so on are *functions* of x. An equation such as $y = 2x^2 + x$ can be written as $f(x) = 2x^2 + x$

geometric sequence (or **series**) each term is multiplied by a multiplier to produce the next term. 2, 4, 8, 16, 32 ... is a *geometric sequence* where the multiplier is 2

global warming the gradual heating up of the Earth gradient slope

graph (verb) to draw a mathematical function as a graphgraphical calculator a calculator that can display graphshemisphere half of a sphere

hence means the same as 'therefore': 2x = 6, hence x = 3

hexagon a shape with six sides

identity (e.g. trigonometrical identity) a statement which is always true, wherever the value of the variables involved. $\tan \theta \equiv \frac{\sin \theta}{\cos \theta}$ is an identity. Notice that you show this by using \equiv instead of =

inequality the opposite of an equation. The equals sign (=) is replaced by an *inequality* sign: >, \ge , <, \le , <> or \ne . For example ' $x \ge 1$ ' means 'x is greater than or equal to 1'

infinite series doesn't end

integer a whole number

intercept where curves or lines cross the x-axis or y-axis

interpret to explain the meaning of something

interval the *interval* of $\int_a^b x \, dx$ is [a, b]

invest to give money to a company such as a bank in order to obtain a profit

kinematics the study of moving objects

landslide a sudden fall of a lot of earth or rocks down a hill, cliff etc

legend a historical story that may not be true

LHS left-hand side

linear where the variables have the power 1.

Hence y = 2x + 3 is *linear* but $y = x^2$ and $y = \frac{1}{x}$

(i.e. $y = x^{-1}$) are not. A *linear function* can be represented by a *straight line*

linear growth where the value over time can be expressed as a straight line

linear programming solving a problem by drawing graphs of inequalities

linear relationship where the power of the variables is 1. y = 4x + 1 (i.e. $y^1 = 4x^1 + 1$) shows a *linear relationship* between x and y, which can be plotted as a *line*. On the other hand, $y = x^2 + x + 1$, $y = \frac{1}{x}$, $y = \sin x$, $y = \lg(x)$ and $y = e^x$ show *non-linear* relationships

local a *local* maximum is greater than *nearby* values on the curve, but not necessarily greater than values *further* away

long division in algebra, a method of dividing one polynomial by another

between x and y, which produce curves

magnitude if a **vector** is drawn on a graph, its magnitude (or *size* or *modulus*) is the distance from the origin to its tip. The magnitude is not a vector itself, but a scalar.

By Pythagoras, the magnitude of $inom{x}{y}$ is $\sqrt{x^2+y^2}$

manipulate (a mathematical expression) to work on it so as to change it into a simpler or more useful form

mathematical modelling using mathematics to understand and/or solve a real-world problem

midpoint a point that is half of the way through or along something

modulus of a vector the vector's magnitude

modulus the positive value of an expression, shown as |x|. |1| and |-1| are both equal to +1. Useful in statements such as |x| < 1, meaning that x is a fraction between -1 and +1. Note that the modulus of a *vector* has a different meaning

multiply out expand. If you multiply out (x + 1)(x + 2), you get $x^2 + 3x + 2$

non-perfect square an expression that can't be factorised into the form $(ax \pm b)^2$

normal is perpendicular to a tangent

obtuse between 90° and 180° ($\frac{\pi}{2}$ and π)

outcome the final result of something – used especially when no one knows what it will be until it actually happens

overlap if two or more things *overlap*, part of one thing covers part of another

parallel two lines that are the same distance apart along their whole length

parallelogram a quadrilateral in which opposite sides are parallel and equal

Pascal's triangle a triangle of numbers where each number is the two numbers above it added together except for the edges, which are all 1.

per annum for each year

perfect square an expression that can be factorised into the form $(ax \pm b)^2$

perimeter the whole length of the border around an area or shape

period how often a curve repeats itself. $y = \sin \theta$ has a period of 360° (2 π)

perpendicular gradient the gradient of a line at right angles (*perpendicular*) to another line

plane a flat surface, for example the side of a cuboid

plot a small piece of land for building or growing things on

polynomial usually has three or more terms, e.g. $x^2 + 5x + 2$

product rule if
$$y = uv$$
 then $\frac{dy}{dx} = u\frac{dv}{dx} + v\frac{du}{dx}$

quadratic an expression such as $x^2 + 2x + 3$, containing a variable to the power of 2 (but no higher)

quadrilateral a shape that has four sides

quotient rule
$$\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$$

radians π radians = 180 degrees

rational any real number, including integers, that can be written as a fraction, even if it produces a repeating decimal. Many functions are irrational: most roots, logarithms and trigonometrical functions, and the constants π and e

rational exponent a rational power e.g. $(a + b)^3$, $(a + b)^{-3}$, $(a + b)^{\frac{2}{7}}$, $(a + b)^{-\frac{2}{7}}$. An expression with a rational *power* may not produce a rational *number*

real any number representing a quantity. 3, -3, $\sqrt{3}$, $\frac{1}{3}$, $\lg 3$, $\sin 3$, π and e are all real. $\sqrt{-3}$ and $\ln (-3)$ are both unreal. Not all real numbers are rational

rebound bounce

reciprocal $\frac{31}{4}$ is the *reciprocal* of $\frac{4}{31}$, and vice versa

reciprocal function $y = \frac{k}{x}$ where k is a constant. Called 'reciprocal' because the variable x has been moved to the denominator; the reciprocal of x is $\frac{1}{x}$

rectangular Cartesian coordinates (x, y)

recurrence relationship where the next term of a sequence is derived from the preceding term(s). The pattern repeats itself (*recurs*) throughout the sequence

remainder the expression left over after dividing

remainder theorem if a polynomial f(x) is divided by (ax - b) then the remainder is $f(\frac{b}{a})$

RHS right-hand side

rotate to turn around an axis or centre

rules of indices how to combine powers, e.g. $x^2 \times x^3 = x^5$

satisfy if a statement is true when another is true, the first *satisfies* the second. The statement x = 2 *satisfies* the equation (x - 1)(x - 2) = 0

savings account a bank account in which you keep money that you want to save, and which pays you interest on that money

scalar a single quantity (unlike a vector).

A vector multiplied by a scalar gives another vector:

$$\begin{pmatrix} x \\ y \end{pmatrix} \times 3 = \begin{pmatrix} 3x \\ 3y \end{pmatrix}$$

sector a part of a circle bounded by an arc and two radii

segment part of a line between two points

sequence a set of terms in which each term is defined by the previous one(s). A sequence may be arithmetic (3, 5, 7, 9, 11 ...), geometric (3, 6, 12, 24, 48 ...), Fibonacci (0, 1, 1, 2, 3...) and so on

series a **sequence** in which the terms are added together: 3 + 5 + 7 + 9 + 11..

shift a change in position

Sigma notation/∑ **notation** uses the Greek 'S' to indicate a Sum of elements, for example in a series

similar triangles have equal angles but not equal sides

simultaneous equations two or more equations, including two or more variables, that must be solved together. If the equations are linear, you will need as many equations as variables. y = x + 1 and 3y = 7x - 5 are two simultaneous equations in two variables.

sine rule
$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
 or $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$

size of a vector is the vector's magnitude

sketch (noun or verb) a sketch is a drawing that explains something without necessarily being accurate

spot to detect something that is hard to see

straight line graph where the variables have a linear relationship

stretch normally means 'enlarge in length, width, height or depth' but in mathematics it can also mean 'reduce in length, width, height or depth'

substitute to replace something (e.g. a *variable*) with something else (e.g. a *value*). Let y = x + 1. If you *substitute* x = 2, you find that y = 2 + 1 = 3

subtend the angle *subtended* by an arc is the angle at the centre of the circle formed by the two radii

tangent a line that touches a curve but does not cross it

tend to move in a particular direction

term a separate part of a mathematical expression. The function $ax^2 + bx + c$ has three terms. The sequence 1, 3, 5, 7 has four terms

the binomial coefficients the numbers preceding the variables in a binomial expansion. The expansion of $(a + b)^4$ is $(1)a^4$, $4a^3b$, $6a^2b^2$, $4ab^3$, $(1)b^4$. Here the *binomial coefficients* are 1, 4, 6, 4, 1

the binomial theorem shows how to expand expressions of the form $(a + b)^n$

the generalised binomial theorem the version (discovered by Newton) used nowadays.

to round (a number) **up** to increase a decimal fraction to the nearest whole number

transformation altering a curve or shape by using arithmetic to change its mathematical definition. Examples are **translation**, rotation, reflection and enlargement

translation (in mathematics) movement from one position to another

triangle pattern Pascal's triangle

variable a mathematical quantity that can represent different (varying) amounts, usually represented by a single letter such as x

vector a quantity that has both size and direction

vertex (plural **vertices**) where two lines meet at an angle, especially in a shape such as a triangle

visual realisation something, e.g. a mathematical expression, represented in a form like something that can be seen in the real world

vital very important

volume of revolution the volume of a solid formed by rotating a curve around a line (usually an axis)

x-intercept/y-intercept where the line or curve crosses the x-axis/y-axis

ANSWERS

CHAPTER 1 – SURDS AND LOGARITHMIC FUNCTIONS

STARTER ACTIVITIES

- 1 ▶ a y¹¹ dx^8
- 2 > a $m^{\frac{3}{2}}$
 - d 18
- 3 ▶ a 4
 - **d** $-\frac{1}{8}$
- **b** $8q^{7}$ e a⁵
- c 27 k⁶ $f 16x^3y^4$
- - $c 4 c^{\frac{1}{3}}$ f $15y^{-1}$
- $c \frac{1}{64}$
- **b** 5

b $3p^{\frac{7}{2}}$

 $p^{\frac{3}{2}}$

 $\frac{27}{64}$

EXERCISE 1

- 1 > a $3\sqrt{2}$
- **b** $5\sqrt{2}$
- c 5√5

- d $8\sqrt{2}$
- e 2√33
- f 5√435

- 2 ▶ a √15
- c √2

- d $\frac{\sqrt{17}}{}$
- **b** $\frac{3}{2}\sqrt{15}$ **e** $\frac{2\sqrt{6}}{3}$
- 3 ▶ a $4\sqrt{3}$
- **b** 9√3
- $c \sqrt{7} + 4\sqrt{43}$

- **a** $4\sqrt{3}$ **d** $24\sqrt{3}$
- e 58√2
- f 30

- g 168
- h 192
- **4 a** $24 12\sqrt{3}$ **b** $54 27\sqrt{29}$ **c** $13 + 10\sqrt{3}$

- **d** $-8\sqrt{7} 2\sqrt{2}$ **e** 13 **f** $26 5\sqrt{42}$

- **g** 3
- **5 22.4%**
- 6 > 34 + $9\sqrt{2}$


EXERCISE 2


- $d\sqrt{2}$
- **e** 4√3
- f √15

- h 2 + $\sqrt{3}$
- **2 a** $2-\sqrt{2}$ **b** $7+4\sqrt{3}$
- $-5 + 2\sqrt{6}$


- **d** $6\sqrt{6} 14$ **e** $4 + \sqrt{10}$ **f** $\frac{5 + \sqrt{2} \, 1}{2}$
- g $\frac{19 + \sqrt{55}}{}$
- h $10\sqrt{30} 8\sqrt{10} 15\sqrt{42} + 12\sqrt{14}$
- $\sqrt{2}$
- $\mathbf{j} \quad \frac{a\sqrt{b} + b\sqrt{a}}{a b} \qquad \mathbf{k} \quad \frac{a\sqrt{b} + b\sqrt{a}}{ab}$


EXERCISE 3







c Intersection occurs at (1, 0).













C 10-





20-15 10 5 0 5 10 15 20 25 30 x











EXERCISE 5

6-0 1 2 3 4 5 6 x

b
$$x = 1.3 \text{ or } 1.4$$


$$x = 4$$

*y*₄ 10--1-0.5 ⁰ 0.5 ¹ 1.5 ² 2.5 ³

b
$$x = 2.4 \text{ or } 2.5$$

$$x = 0.6 \text{ or } 0.7$$

3 a
$$x = 30, y = 2.60; x = 75, y = 1.98$$



c
$$2 + 2\cos x = 5\sin 2x$$

 $\Rightarrow 2 = 5\sin 2x - 2\cos x$
Draw $y = 2$
and intersections at ≈ 25.1 and 74.6
or 25 and 75 to 2 sf

EXERCISE 6

1 \triangleright a $\log_4 64 = 3$

b
$$\log_5 8\left(\frac{1}{25}\right) = -2$$

$$\log_{8}(262144) = 6$$

d
$$\log_3 9 = x$$

$$\log_8 1 = x$$

$$\frac{1}{27}$$

$$\frac{3}{2}$$

$$\frac{1}{2}$$

 $\frac{1}{5}$

- 1 ▶ a log₄64
- b log₉6
- c log₅81

- d log₄120
- log₆1
- f log₂45

- h 2
- **2 a** $4\log_a x + 3\log_a y + \log_a z$ **b** $\log_a x 3\log_a y$
 - $c 2\log_a x + 2\log_a z$
- $d \log_a x \log_a y \log_a z$
- e $\frac{1}{2}\log_a x + \frac{1}{2}\log_a y$ f $2\log_a x + \log_a y + \frac{3}{2}\log_a z$
- $\frac{3}{2}\log_a x + \frac{7}{2}\log_a y 3\log_a z$

EXERCISE 8

- 1 **a** 3.21
- **b** 1.68

- d 0.558
- e 0.719
- c 1.93

- 2 **a** 1.51
- **b** 3.01 **c** 0.406
 - d 2.59
- f 1 **e** 1.46
- g 0.861
- **3 a** $x = 512, x = \frac{1}{2}$
- **b** x = 216, x = 6
- $x = \frac{1}{10}, x = \frac{2}{100000}$
- **d** $x = \frac{14}{9}$

EXERCISE 9

- 1 \triangleright a x = 1.79
- **b** x = 1.86
- x = 1.91

- d x = -0.712
- x = 1.45h x = -58.4
- x = 0.421x = 4.99

- x = -1
- x = 0.542
- **b** x = 1.23 or -0.0733
- **2 a** x = 0.792x = 1.21
 - x = 0.161
 - x = 0.387
- $\mathbf{f} \quad x = 2$

EXAM PRACTICE QUESTIONS

- 1 > $7\sqrt{2}$
- $2 > 3\sqrt{2} 2$
- 3 ▶ a $16 4\sqrt{5}$
- **b** $4 \sqrt{5}$

- **4** ▶ 2√3
- 5 ► $5 \sqrt{5}$
- 6 -4 2
- (0.254, 0)7 > x = 0.739
- 8 > a Proof

b $x = 9 \text{ or } \frac{1}{3}$

- 9 > x = 6 or 3
- 10 > $y = 8 \text{ or } \frac{1}{8}$
- 11 $\triangleright y = \frac{1}{9}b^5$
- **12** $\rightarrow x = 1 \text{ or } 1.21$
- **13** \triangleright x = 1.43
- **14** \triangleright $b = (\frac{1}{9})(a 5)$
- **15** $\rightarrow x = 0.89$
- **16** \triangleright t = 0.68
- 17 \triangleright a 4 + p
- **b** 4p-1 **c** $x=2\sqrt{2}$
- 18 $\triangleright t = 1$
- 19 ▶ a 10



- b Accept 1.6–1.7 range of answers
- c Accept 0.1-0.2 and 3.1-3.2 as a range of answers

CHAPTER 2 – THE QUADRATIC FUNCTION

STARTER ACTIVITIES

- 1 **a** 3x(2x + 3)
- **b** 2b(b+4)
- \circ 9m(qm-3)
- d 9xy(y+4x)
- 8x(3 8x)
- **2** \triangleright **a** (x + 3)(x + 6)
- **b** (x-3)(x-4)
- (x + 1)(x 3)
- (x + 3)(x + 12)
- (x + 3)(x + 9)
- 3 \triangleright a (x + 3)(x 3)
- **b** (x + 5)(x 5)
- (3x+4)(3x-4)
- d (5x + 4)(5x 4)

- 1 **a** (x-3)(3x+2)
- **b** (2x + 1)(x + 5)
- (2x+1)(x-4)
- d (5x-1)(x-3)
- (3x-4)(2x+3)
- $(3x-1)^2$
- $\mathbf{g} (3x + 1)(3x 7)$
- **b** x = -4 or $x = -\frac{1}{4}$
- **2 a** $x = -2 \text{ or } -\frac{1}{2}$ **c** $x = 4 \text{ or } x = \frac{1}{3}$
- **d** $x = 1 \text{ or } \frac{2}{3}$
- **e** $x = -6 \text{ or } \frac{1}{3}$
- $x = \frac{1}{5} \text{ or } -3$
- **g** $x = \frac{2}{3}$

1 > a
$$(x + 2)^2 - 4$$

$$(x-8)^2-64$$

$$3(x-4)^2-48$$

$$(x-\frac{1}{2})^2-\frac{49}{4}$$

$$(x+\frac{1}{2})^2-\frac{5}{4}$$

f
$$(x-1)^2-\frac{2}{3}$$

$$(x+\frac{3}{4})^2-\frac{17}{16}$$

$$\left(x+\frac{3}{4}\right)^2-\frac{13}{16}$$

2 a
$$x = -\frac{2}{3}, x = \frac{5}{2}$$

b
$$x = \sqrt{3} - 1, -\sqrt{3} - 1$$

c
$$x = \pm \frac{\sqrt{7} + 3}{2}$$
 d $x = -\frac{3}{4} \pm \frac{\sqrt{57}}{4}$

d
$$x = -\frac{3}{4} \pm \frac{\sqrt{57}}{4}$$

e
$$x = 15, x = -\frac{1}{4}$$
 f $x = -1 \pm \frac{\sqrt{13}}{2}$

f
$$x = -1 \pm \frac{\sqrt{13}}{2}$$

g
$$x = -\frac{3}{2} \pm \frac{\sqrt{39}}{2}$$

h
$$x = \frac{1}{8} \pm \frac{\sqrt{129}}{8}$$

EXERCISE 3

1 > a
$$x = 0, 6$$

b
$$x = 0, x - 25$$

$$x = 3, -2$$

d
$$p = 2 \pm \sqrt{2}$$

e
$$m = \pm \sqrt{3} - 1$$

$$x = -5, -2$$

$$t = 6, -1$$

h
$$x = 5, -7$$

$$n=2$$

$$x = -3 \pm \sqrt{3}$$

2 a
$$x = \frac{1 + \sqrt{26}}{3}, x = \frac{1 - \sqrt{26}}{3}$$

b
$$x = \frac{3 + \sqrt{3}}{3}, x = \frac{3 - \sqrt{3}}{3}$$

$$x = \frac{3}{2}, x = -\frac{2}{3}$$

d
$$x = \frac{-3 + \sqrt{17}}{4}, x = \frac{-3 - \sqrt{17}}{4}$$

$$x = -\frac{1}{2}, x = -3$$

f
$$x = \frac{-7 + \sqrt{37}}{6}$$
, $x = \frac{-7 - \sqrt{37}}{6}$

g
$$x = \frac{5}{2}, x = \frac{3}{2}$$

h
$$x = \frac{-5 + \sqrt{109}}{14}, x = \frac{-5 - \sqrt{109}}{14}$$

i
$$x = \frac{15 + \sqrt{545}}{20}, x = \frac{15 - \sqrt{545}}{20}$$

j
$$x = \frac{\sqrt{17} - 1}{4}, x = -\frac{1 + \sqrt{17}}{4}$$

$$x = 3\sqrt{7} + 7$$
, $x = -3\sqrt{7} + 7$

$$x = \frac{-15 + \sqrt{17}}{8}, x = \frac{-15 - \sqrt{17}}{8}$$

EXERCISE 4

- 1 > a Equal roots
- b 2 roots
- c 2 roots
- d No roots

Equal roots

- e Equal roots
- f 2 roots
- g No roots
- h 2 roots
- No roots
- k No roots

$$2$$
 ▶ $p > -\frac{1}{7}$

$$3 \triangleright m = \frac{1}{3}$$

EXERCISE 5

1 > a
$$x^2 + 8x - 1 = 0$$

b
$$x^2 - 6x + 8 = 0$$

2 > a
$$x^2 - 8 = 0$$

b
$$x^2 - 34x + 1 = 0$$

3 > a
$$x^2 - x - 1 = 0$$

b
$$x^2 - x - 1 = 0$$

EXAM PRACTICE QUESTIONS

1 **a** i
$$3x^2 - 56x + 3 = 0$$

ii
$$9x^2 - 90x + 194 = 0$$

b
$$\frac{5}{3} \pm \frac{\sqrt{31}}{3}$$

$$2 > 5x^2 + 9x - 15 = 0$$

b
$$x^2 - 7x + 12 = 0$$

$$2x^2 - 25x + 12 = 0$$

b
$$p > 1$$
 or $p < -3$

6
$$\triangleright$$
 a $(x + 3)(5x + 1)$

b
$$(x-1)(3x-4)$$

7 a
$$p = -1, -2$$

b
$$x = \frac{2}{3}$$
 or -5

8 a
$$x = \frac{-3}{10} \pm \frac{\sqrt{29}}{10}$$
 b $x = 3.82$ or 1.17 (3 s.f.)

b
$$x = \frac{2}{3}$$
 or -5

8 a
$$x = \frac{-3}{10} \pm \frac{\sqrt{29}}{10}$$

b
$$x = 3.82 \text{ or } 1.17 \text{ (3)}$$

9 a
$$a = -2, b = -1$$
 b -4

$$-4$$

10
$$\rightarrow$$
 $x = 0, x = \frac{2}{3}$

11 > a
$$\frac{37}{4}$$

b
$$\pm \frac{5}{9}$$

$$\pm \frac{215}{8}$$

CHAPTER 3 – INEQUALITIES AND IDENTITIES

STARTER ACTIVITIES

- 1 **a** (9, 6)
- **b** (4, 2
- **c** (6, 1)
- (-3, 2)
- 3 ▶ a 17.4
- **b** 18.4
- **c** 17.03
- d 38.89

EXERCISE 1

- **1** ▶ **a** (1, 2) (-2, -1)
- **b** (2, 5) (1, 3)
- **c** (2, 1) (1, 2)
- **d** (2, 5) (-2, -3)

- e (2, -1)f (-3, -3)(-1, 1)g $(-3, 4)(\frac{7}{2}, -\frac{5}{2})$ h $(2, -1)(-\frac{7}{8}, \frac{19}{4})$ i $(-\frac{3}{2}, \frac{9}{4}), (4, -\frac{1}{2})$ j $(\frac{21}{4}, \frac{8}{3}), (2, 7)$
- $(0, \frac{7}{2}), (-3, -4)$ $(\frac{13}{4}, \frac{5}{8}), (\frac{7}{2}, \frac{1}{2})$
- m (6, 3) (2, 7)
- **2** ▶ **a** (0, 0), (-3, 3), (4, 108) **b** (0, 0), (2, 14), (-1, -4)
- - **c** (0, 0), (-2, 10), (1.5, 3) **d** $(0, 0), (-2, 10), (\frac{3}{2}, 3)$

EXERCISE 2

- **1 a** x < 2
- **b** $x > \frac{12}{7}$ **c** $x \ge \frac{1}{2}$

- **d** $x \ge \frac{1}{3}$ **e** $x > -\frac{4}{41}$ **f** $x < \frac{2}{13}$
- **g** $x \le -55$ **h** $x \ge -\frac{17}{5}$
- **2 a** $x \ge -\frac{7}{2}$ **b** x < -5 **c** x > -22 **d** $x \le -\frac{11}{5}$ **e** $x \ge \frac{3}{13}$ **f** $x > -\frac{7}{3}$ **g** $x \ge \frac{43}{25}$ **h** $x < \frac{1}{9}$

- **3 a** $x > \frac{5}{2}$ **b** $x > \frac{1}{5}$ **c** $\frac{1}{7} \le x \le \frac{21}{11}$
 - d No solution
- **e** $x \ge \frac{31}{18}$ **f** $-10 < x < -\frac{1}{13}$

EXERCISE 3

- **1 a** -3 < x < 2
- **b** -3 ≤ x ≤ 4


- c $-4 < x < -\frac{3}{2}$ e $x \le -\frac{1}{2}$ or $x \ge \frac{3}{4}$ f $\frac{3}{4} \le x \le \frac{7}{5}$


- **2** \triangleright **a** -2 < x < 9
 - **c** $x \le 12, x \ge -1$ **d** 4 < x < 10


b x = -3

- **e** x < -4, x > -3 **f** $-\frac{4}{7} \le x \le 5$ **g** $x < -7, x > \frac{3}{2}$ **h** $-\frac{9}{5} < x < 2$
- i $x \le -3, x \ge \frac{2}{3}$ $\mathbf{k} \ x \leq -4, x \geq 4$
 - \mathbf{j} $-5 \leqslant x \leqslant 5$


- > y < 3</p>
- $y \ge 2, y < x$
- $x \le -3, y \ge -x$
- $y \le x, y > -6, x \le 5$
- y < 3x + 6, y > 3x 10
- $y \le \frac{1}{2}x 6, y \le -2x 6$







11 ▶



12 ▶ **a** $c + l \ge 10$

$$120c + 200l \le 1800$$

- b 3 chickens and 7 lambs for a profit of £3600
- 13 ➤ She can earn a maximum of 184 JOD by working 12 hours as a wedding photographer and 8 hours as a studio photographer.
- 14 ► Let type A be x and type B be y

Maximum profit is 15x + 12y

Maximise P = 15x + 12y

Subject to

$$x \ge 50$$


$$\frac{1}{5}(x+y) \le x \Rightarrow y \le 4x$$

$$x \le \frac{2}{5}(x+y) \Rightarrow y \le \frac{3}{2}x$$

$$3x + 2y \le 200$$

$$y \ge 0$$

Representing these equations gives:



Profit is maximised at the point (50, 25). Substituting (50, 25) into P=15x+12y gives P=1050 So the maximum profit is £1050.

EXERCISE 5

- 1 **a** $x^2 2x 15$
- **b** $x^2 + 5x 6$
- $x^2 + 5x + 3$
- d $x^2 + 7x + 10$
- $x^2 3x + 7$
- $x^2 3x + 2$
- $x^2 + 1$
- h $x^2 + 2 \frac{9}{x+5}$

- **2 a** $-2x^2 x + 5$
- **b** $4x^2 + 8x 32$
- $6x^2 + 3x + 2$
- d $3x^2 + 2x 2$
- $= -3x^2 3x + 90$
- $f -5x^2 + 3x + 5$
- $2x^2 + x 1$
- 3 > a $x^2 + 3x 1$
- **b** $2x^2 5x 3$
- $x^2 3x 10$
- d $x^2 x 6$
- $2x^2 + x 2$
- $x^2 + 3x + 2$

EXERCISE 6

- 1 > a Proof
- b Proof
- Proof
- d Proof
- Proof
- 2 Proof
- $3 \triangleright (x-1)(x^2-x-4)$
- 4 \triangleright (x-1)(x+2)(x-3)
- (x-2)(x+4)(2x-1)
- **6** $\triangleright a = 4$
- 7 \triangleright a b = 2
- **b** (x + 1)(x 1)(3x + 2)
- 8 p = 6, q = -13
- 9 \triangleright a a = 3, b = -1
- **b** (x + 2)(x 2)(3x 1)

EXERCISE 7

- 1 ba 2
- **b** 27
- **c** 2
- **d** −6

- **b** 1
- c 171

- **3** ▶ −80
- **4** \triangleright *b* = 7
- **5** ▶ 16*p* − 176
- **6** ▶ *b* = 32
- 7 $\triangleright a = \frac{1}{3}, b = \frac{4}{3}$
- **8 a** $\alpha = 2, b = 4$
- $\frac{13}{27}$
- 9 $\triangleright a = 5, b = -8$
- **10** \triangleright p = 8, q = 3

EXAM PRACTICE QUESTIONS

- 1 $x = -3, y = 5 \text{ or } x = \frac{5}{3}, y = \frac{1}{3}$
- **2** \blacktriangleright (4, 10) $\left(\frac{1}{3}, -1\right)$
- 3 \blacktriangleright (1, 3) $\left(-\frac{19}{5}, -\frac{9}{5}\right)$
- 4 > a x > 4
- **b** x < -2, x > 6

b
$$x < \frac{14}{5}$$

$$-1 < x < 2$$

6
$$x < -\frac{3}{2}, x > 4$$

7
$$\blacktriangleright$$
 $\left(4,\frac{1}{2}\right)(-5,-1)$

8 a
$$p < -5$$
 or $p > \frac{1}{2}$ **b** $-5 < n < -2$

b
$$-5 < n < -2$$

b
$$(x+2)(x-4)(2x-3)$$

b
$$x^2 + 3x + 2 + \frac{6}{x+5}$$

b
$$-(x-4)(3x^2-x+2)$$

12
$$\triangleright$$
 $(x + 1)(x - 4)(2x - 1)$

13 a
$$p = -20$$

b
$$q = -6$$

b
$$(x+3)(x^2-3x+7)$$

15
$$\triangleright$$
 $p = -20$


16 a
$$\alpha = -20$$

b
$$(x-4)(x^2+3x+5)$$

17 a
$$(x-1)(x-1)(2x+3)$$
 b $x=1$ and $x=-\frac{3}{2}$

$$x = 1 \text{ and } x = -\frac{3}{2}$$





19 > a Here is the summary.

Let x be the number of desk top diaries bought. Let y be the number of pocket diaries bought.


Minimise c = 6x + 3y

$$y \geqslant 80$$

$$2x \leqslant y$$


$$x + y \ge 400$$






CHAPTER 4 – SKETCHING POLYNOMIALS


STARTER ACTIVITIES









h-i















































b





C




























f y 10 10 5 1 5 1 5 1 6 3 1 6































b (0, -1), (1, 0), (3, 8)








There is no intersection.




10 5

-5 -10 5 10 x


-10 -5 0



One solution







20 -


10


5 0

-10 -20


5 10 15 x














EXAM PRACTICE QUESTIONS




(0, 5) and (-2.5, 0)






b (-1, 0), (-0.5, 0), (2, 0)



b Two, because the curves only intersect twice.





 $\left(\frac{7}{2} + \frac{\sqrt{33}}{2}, \frac{1}{2} - \frac{\sqrt{33}}{2}\right)$

and


$$\left(\frac{7}{2} - \frac{\sqrt{33}}{2}, \frac{1}{2} + \frac{\sqrt{33}}{2}\right)$$

5 **a**
$$x(x^2-3)$$








7 \triangleright *x* asymptotes are $x = \pm 3$, *y* asymptote remains the same




8 a a = 1 and b = 4









b There are no intersections.

CHAPTER 5 – SEQUENCES AND

- b Yes
- Yes

- No
- Yes

- h Yes
- Yes Yes

- a 21
- **b** 11.5
- c 105

- d 10x
- e 22 + 10a

- 3 ▶ a \$5800
- \$(3800 + 200n)
- **a** 33
- **b** 17
- c 21

- d 13
- e 18
- \mathbf{f} n

- 1 \triangleright a 60, 3n+1
- **b** 74, 3n + 14
- \circ 26, 106 -4n
- d -98, 22 6n
- e $\frac{113}{15}$, $\frac{1}{3}n \frac{13}{5}$
- f 39q, q(2n-1)
- $\mathbf{g} = -71k, k(9-4n)$

- 2 ▶ a 20 b 120 c 33 d 51 e 27 f 42
- **3** ▶ 4
- **4 a** A = 42, d = -15 **b** 4th term

- **5 22.2**
- **6** \triangleright P = 5; 25, 20, 15
- **7** > $7 \times \frac{1}{2}$, n = 8

EXERCISE 3

- 1 **a** 650
- **b** 960

- d 2312.5
- e 21(11x + 1)
- **2** ▶ **a** 50 **b** 20
- **c** 32
- d 321

- **3** > 900
- **4 >** \$652000
- **5 2**0100
- 6 > 36 168 euros
- **7** ▶ 3850
- **8** > 33
- 9 Proof
- **10** > 1800
- **11 a** $\sum_{1}^{10} (2r+1)$ **b** $\sum_{1}^{12} (8r+1)$

 - c $\sum_{10}^{10} 4(10-r)$ d $\sum_{10}^{8} 4r$

- 12 Proof
- **13** ▶ **a** 465
- **b** 630
- c −432
- d 511

- **14** > 49
- **15** > 32

EXERCISE 4

- 1 > a Yes, r = 5
- b No
- Yes, r = 0.5

- d No
- **e** Yes, r = -1 **f** Yes, r = 1

- g Yes, $r=\frac{1}{4}$
- h No

- 2 ▶ a -3051, 9153, -27 459
 - **b** 1280, 5120, 20 480
 - **c** 6, 18, 54
 - d 375, 1875, 9375
 - e −0.015625, 0.00390625, −0.00009765625
 - $f x^3, x^4, x^5$
- 3 > a 8

b 64

EXERCISE 5

- **1 a** 48, 192, 3(2ⁿ⁻¹)
- **b** 10, 2.5, $\frac{160}{2^{n-1}}$
- c 162, 1458, $2(-3)^{n-1}$ d $-16, -64, -(2^{n-1})$
- **2** \triangleright **a** a = 3, r = 2
- **b** a = 15, r = -2
- **c** a = -30, r = -0.5 **d** a = 0.5, r = 2
- 3 > a 3

b 1875

- 4 \ ±\frac{1}{4}
- 5 ► -6 (from p = 0), 4 (from x = 10)

EXERCISE 6

- 1 **a** 1093
- **b** 33.33...
- c 28697812

f 4092

- d 426.5625
- 97655
- g 544 (3 s.f.) h 308829
- **2 18446744073709551615**
- 3 ▶ 25 years
- 4 ▶ 10644 rupees
- 5 > The second option.
- 6 > 35 days
- **7** ▶ 19
- 8 \triangleright n = 6, r = 4

EXERCISE 7

- b Doesn't exist
- c Doesn't exist
- d $\frac{1}{1+2y}$. If $|y| < \frac{1}{2}$
- **2 a** $r = \frac{1}{2}$

b 1000

- **4 >** 3
- **5 2**0.25
- 6 > 18 m
- 7 ▶ 625πcm²
- 8 > £4 million

EXAM PRACTICE QUESTIONS

1 a
$$\alpha = 13$$

2
$$\triangleright$$
 a $q = 4.9$

4 a
$$a = 6$$
 and $d = 3$ **b** $n = 12$

b
$$n = 12$$

5 a
$$d = -7$$
 and $a = 312$ **b** $n = 45$

8 a Common ratio =
$$1 - \sqrt{2}$$
 first term is $\sqrt{2}$

b i
$$r = 1 - \sqrt{2} \approx -0.41$$
 since $|r| < 1$, series is convergent

ii
$$S_{\infty} = \frac{\sqrt{2}}{1} - (1 - \sqrt{2}) = \frac{\sqrt{2}}{\sqrt{2}} = 1$$

c
$$333\frac{1}{3}$$

d
$$8.95 \times 10^{-4}$$

c Common ratio =
$$\frac{3}{4}$$
 d 64

CHAPTER 6 – THE BINOMIAL **SERIES**

EXERCISE 1

1 > a
$$27x^3 + 27x^2y + 9xy^2 + y^3$$

b
$$m^4 - 4m^3n + 6m^2n^2 - 4mn^3 + n^4$$

$$\mathbf{c}$$
 1 + 9x + 27x² + 27x³

d
$$1+y+\frac{3y^2}{8}+\frac{y^3}{16}+\frac{y^4}{256}$$

$$= 27y^3 + 108y^2x + 144yx^2 + 64x$$

$$d + 27405$$

3 > a
$$1 + 12x + 54x^2 + 108x^3$$

b
$$1 - \frac{7y}{4} + \frac{21y^2}{16} - \frac{35y^3}{64}$$

$$\circ$$
 343 - 147b + 21b² - b³

$$d + 414720x^6 + 103680x^4 + 15360x^2 + 1024$$

4
$$\triangleright$$
 a $n = 5$

$$+\frac{16x^4}{125}$$

EXERCISE 2

1 > a 1 + 9
$$x$$
 + 36 x ² + 84 x ³

b
$$1 - 15y + 90y^2 - 270y^3$$

$$1 + 28x + 336x^2 + 2240x^3$$

d
$$1 + \frac{9x}{2} + 9x^2 + \frac{21x^3}{2}$$

$$= 1 - 21x + 189x^2 - 945x^3$$

$$f \quad 1 - 2x + \frac{5x^2}{3} - \frac{20x^3}{27}$$

g
$$1 - \frac{3x}{2} + \frac{15x^2}{16} - \frac{5x^3}{16}$$

3
$$\triangleright$$
 a 1 - 8x + 28x² - 56x³

4 a
$$1 + x + \frac{7}{16}x^2 + \frac{7}{64}x$$

6 a
$$p = 6$$
 b $\frac{20}{3}$

$$\frac{20}{3}$$

$$\frac{160}{27}$$

EXERCISE 3

1 > a
$$1 - 6x + 24x^2 - 80x^3$$
, $|x| < \frac{1}{2}$

b
$$1 + 3x + 9x^2 + 27x^3$$
, $|x| < \frac{1}{3}$

c
$$\frac{1}{2} - \frac{3}{8}x + \frac{27}{64}x^2 - \frac{135}{256}x^3$$
, $|x| < \frac{2}{3}$

2 ▶ **a**
$$-1$$
, $|x| < 1$

b
$$\frac{1}{2}$$
, $|x| < \frac{1}{2}$

27,
$$|x| < \frac{1}{3}$$

b
$$\frac{1}{2}$$
, $|x| < \frac{1}{2}$ **c** 27, $|x| < \frac{1}{3}$ **d** $-\frac{5}{81}$, $|x| < 1$

3
$$\triangleright$$
 a 1 - 3 x + 9 x^2 - 27 x^3

$$1-4x+12x^2-36x^3$$

4
$$\triangleright$$
 $b = \pm 8, \pm 160$

5 a
$$1 + 2x - 4x^2 + \frac{40}{3}x^3$$

b when
$$x = 0.004$$
,

$$(1 + 6x)^{\frac{1}{3}} \approx 1 + 2(0.004) - 4(0.004)^2 + \frac{40}{3}(0.004)^3$$

= 1.007936853

$$(1 + 0.024)^{\frac{1}{3}} = \sqrt[3]{1.024} = \sqrt[3]{\frac{512 \times 2}{1000}} = \frac{4}{5}\sqrt[3]{2}$$

$$\sqrt[3]{2} = \frac{5}{4} \times 1.007936853 = 1.259921$$
 (7sf)

EXAM PRACTICE QUESTIONS

1
$$> 1 + \frac{7x}{4} + \frac{21x^2}{16} + \frac{35x^3}{64}$$


2 a
$$243 + 405ax + 270a^2x^2 + 90a^3x^3$$


b
$$a = \frac{3}{4}$$


4
$$-\frac{21}{2}$$

- **5 a** $=\frac{1}{2}-\frac{1}{16}x$, $+\frac{3}{256}x^2-\frac{5}{2048}x^3+\dots$
 - **b** |x| < 4
- **6 a** $1 + \frac{3}{2}x \frac{9}{8}x^2 + \frac{27}{16}x^3$
 - b 10.14889188, accurate to 8 d.p.
 - c Students should be writing something similar to 'accuracy is over 98%'.
- **7** ▶ 2
- **8 a** $1-4x-8x^2-32x^3$
 - **b** $|x| < \frac{1}{2}$
 - c = 4.79584 (5 d.p.)
- 9 **a** $1 \frac{9}{2}x + \frac{27}{8}x^2 + \frac{27}{16}x^3$
 - **b** x = 0.01, 9.553358 (6 d.p.)
- **10 a** $5 + 5x + \frac{15}{2}x^2 + \frac{25}{2}x^3$
 - **b** $|x| < \frac{1}{2}$
- **11 a** $4 \frac{4}{3}x + \frac{2}{3}x^2$
 - **b** $3\frac{131}{150}$
- **12 a** $2-2x+3x^2-5x^3$
 - **b** $|x| < \frac{1}{2}$

-2**a**







- **3** ▶ 31.43
- **4 25.6**
- **5** \triangleright **a a** + **b b** -**c b a c** -**d** + **a** + **b d d a**
- **6 a** $\frac{1}{2}$ **a b** $-\frac{1}{2}$ **a** + **b** $-b + \frac{1}{2}a$
- 7 Proof

EXERCISE 2

1 a 2a + b **b** $\frac{1}{2}(2a + b)$ **c** $\frac{1}{2}(-2a - b)$

a b **b** -b **d** b + $\frac{1}{4}$ **a e** DE = b - $\frac{3}{4}$ **a**

3 ▶ a Yes b No c Yes d No

4 a $\lambda = \frac{1}{2}, \mu = -4$ **b** $\lambda = -2, \mu = 3$ **c** $\lambda = \frac{2}{5}, \mu = 4$ **d** $\lambda = -\frac{4}{3}, \mu = -\frac{1}{3}$

e $\mu = \frac{7}{3}, \lambda = 4$ g $\lambda = -1, \mu = 7$ f $\lambda = \frac{1}{3}, \mu = \frac{16}{3}$

5 a $\frac{1}{2}$ **b b b** - **a c** $\frac{1}{5}$ **b**

 $-\frac{4}{5}$ **b e** $-\mathbf{a} + \frac{1}{5}$ **b**

7 ▶ **a** 6**b** + 2**a**

b 6**b** – 6**a**

8 > p = $\frac{2}{3}a + \frac{1}{3}b$

EXERCISE 3

1 > $\frac{5}{6}$ a + $\frac{1}{6}$ b

 $2 - \frac{1}{2}a - \frac{1}{2}b + c$

 $\overrightarrow{OC} = -2a + 2b$, $\overrightarrow{OD} = 3a + 2b$, $\overrightarrow{OE} = -2a + b$

- 1 \triangleright a $\overrightarrow{AB} = \mathbf{y} \mathbf{x}$ b Proof $\overrightarrow{OM} = \frac{1}{2}(\mathbf{x} + \mathbf{y})$ $\overrightarrow{DC} = 2(\mathbf{y} - \mathbf{x})$
- 2 Proof
- **3** \triangleright **a** $\overrightarrow{AB} = \mathbf{y} \mathbf{x}$ **b** Proof $\overrightarrow{OC} = -2x$ $\overrightarrow{OD} = -2\mathbf{y}$ $\overrightarrow{DC} = 2\mathbf{v} - 2\mathbf{x}$
- 4 Proof
- **5 a** \overrightarrow{AB} **b a b** \overrightarrow{OM} = $\frac{1}{3}$ **a** + $\frac{2}{3}$ **b c** Proof

EXERCISE 5

- **b** 6i 6i 1 > a 6i + 8j
 - d -23i 12j
- 2 ▶ a 9i + 14j
- b 2√34 c i + 8j
- d √65
- e -2i + 8j
- **3 a** A = 4i 2j, B = 6i + 4j, C = -2i + 6j
 - **b** 2i + 6i
- c -6i + 8i d $2\sqrt{10}$

c i + 6j

- e 2√10 4 > a -i + 5j
- f 10
- **b** 3i 4j
- c -2i j

- 5 ▶ 5i + 6j
- **6 a** $=\frac{6\mathbf{i}+3\mathbf{j}}{3\sqrt{5}}$, **b** $=\frac{1}{10}(6\mathbf{i}-8\mathbf{j})$, **c** $=\frac{-5\mathbf{i}+14\mathbf{j}}{\sqrt{221}}$
- $7 \triangleright m = 3, n = 1$
- 8 p = -2, q = 5
- **9 a** $\binom{8}{-15}$ **b** 17 **c** $\binom{12}{0}$ **d** 12

EXAM PRACTICE QUESTIONS

- 1 Proof
- 2 Proof
- $3 > \frac{4}{3}$
- $4 > -1\frac{1}{2}$
- **5 a** 9i + 3j **b** Proof **c** $2\sqrt{10}$
 - d unit vector is $\frac{1}{2\sqrt{10}}(6\mathbf{i} 2\mathbf{j})$ or $\frac{-1}{2\sqrt{10}}(6\mathbf{i} 2\mathbf{j})$
 - e Proof
- **6 a** Zaynab: $\begin{pmatrix} 10 \\ 14 \end{pmatrix}$ Asaad: $\begin{pmatrix} 8 \\ 10 \end{pmatrix}$ Ali: $\begin{pmatrix} 6 \\ 4 \end{pmatrix}$
- - **b** i Zaynab: 17.2 km ii 2.9 km⁻¹ ii 2.1 km⁻¹ ii 1.2 km⁻¹ Asaad: 12.8 km
 - i Ali: 7.2 km

- 7 \triangleright a $\alpha + b$
- b Proof
- **a** -4a + 4b
- **b** 4a **c** -2a + 8b **d** Proof

- Proof
- 11 Proof
- **12** ▶ a μa
- **b** $\lambda(\mathbf{a} + \frac{3}{4}\mathbf{b})$ **c** $\lambda = \frac{1}{3}, \mu = \frac{1}{3}$

CHAPTER 8 – RECTANGULAR CARTESIAN COORDINATES

STARTER ACTIVITIES

- **1 a** m = 2, c = 5
- **b** m = 2, c = -3
- $m = \frac{1}{3}, c = 0$
- **d** m = 0, c = 6
- $m = \frac{2}{3}, c = 9$

EXERCISE 1

- 1 **a** 0.5
- **b** 4

- 2 > d = 5
- $3 \triangleright a = 3$
- $4 \triangleright a = -4$
- 5 > a = 3.3
- **6** ▶ −10.1
- **7 a** $AB = \frac{3}{2}, BC = -\frac{2}{3}, CD = \frac{3}{2}, DA = -\frac{2}{3}$
 - **b** The lines are parallel.

- **1 a** y = 3x **b** y = 24 6x **c** $y = \frac{1}{2}x 7$
- 2 > 19y 7x 219 = 0
- $y = \frac{1}{16}x + \frac{61}{8}$
- $4 \triangleright y = -\frac{3}{7}x + \frac{96}{7}$
- $5 \triangleright 2x + 5y + 20 = 0$
- **6** \triangleright 3y = 2x + 9
- **7 (3, 0)**
- 8 $y = -\frac{2}{3}x + 7$

- 1 > a Perpendicular
- b Parallel
- Parallel
- d Perpendicular e Neither
- f Neither

- 2 > 5y = -x + 37
- 3 > a y = -2x + 3
- **b** 4y = 22 x
- 2y = x + 1
- d 2y + 3x + 1 = 0
- 4 > -3x + 4y + 25 = 0
- 5 > 5x + 4y 42 = 0
- **6** \triangleright 3x + 2y 27 = 0
- 7 > 4y = 3x 33

EXERCISE 4

- 1 > a 10
- **b** 13
- **c** 5

- d $4\sqrt{2}$
- e 2√34
- f $a\sqrt{53}$

- **g** $b\sqrt{181}$
- h $c\sqrt{10}$

EXERCISE 5

- 1 **a** (0, 6)
- **b** (0, 2)
- c $(2, 3\frac{2}{7})$
- **d** (5, 5)

- $2 \triangleright \left(-\frac{2}{7}, \frac{13}{7}\right)$
- $3 \triangleright (-1, 3)$

EXAM PRACTICE QUESTIONS

- 1 No
- $2 \triangleright 6y 5x 15 = 0$
- 3 **a** $\sqrt{41}$ **b** 8
- c 20

- 4 > x + 2y 21 = 0
- 5 > x + 5y 170 = 0
- 6 > 3x y 22 = 0
- 7 > a $(5, \frac{7}{2})$
- **b** y = 0.5x + 1
- 8 Distances between points are parallel and equal.
- 9 > 20 square units
- 10 > 20.79 units
- **11** \triangleright **a** 2y = x + 5
- **b** (7, 6)
- c 4√5

- d 30 square units
- **12** \triangleright $\left(1, \frac{7}{2}\right)$ or (1, 3.5)
- **13** \triangleright $-\frac{3}{2}, \frac{3}{2}$
- **14 a** x + 2y 5 = 0
- **b** 7√5
- **15** \triangleright **a** y = -x + 6
- b 18 square units

CHAPTER 9 – DIFFEREN

STARTER ACTIVITIES

- 1 > a a 13
- a^{10}
- a^{-14}

- a^{-6}
- a^{-2}
- a^{-6}

- g a8
- $h a^{\frac{5}{2}}$
- $a^{-\frac{3}{2}}$ a^3

- a^2
- $k a^{-3}$
- o a12b2

- $\frac{a^6}{b^{12}}$

- $a^{-4}b^4$
- $~~ {\scriptstyle {\rm U}} ~~ {\displaystyle \frac{a^{-4}b^{-1}}{c^{-3}}}$

- $\frac{m^{-2}}{x^{-1}}$
- $\frac{x^{-8}y^{-4}}{z^{-12}}$
- $x a^{-2}b^{\frac{16}{3}}$

EXERCISE 1

- 1 > a $8x^7$
- **b** $3x^2$
- $\frac{1}{2}x^{-\frac{1}{2}}$
- **d** $\frac{3}{2}x^{\frac{1}{2}}$ **e** $-3x^{-4}$ **f** $\frac{1}{2}x^{-\frac{1}{2}}$

- $\frac{1}{3}x^{-\frac{2}{3}}$ $-\frac{1}{3}x^{-\frac{4}{3}}$
- k 1

- $-\frac{7}{2}x^{-\frac{9}{2}}$
- 2 ▶ a 6x
- $-6x^{-3}$
- **a** 6x **b** $28x^3$ **d** $-3x^{-2}$ **e** $25x^4$
- $\int \frac{3}{2}x^{-\frac{1}{2}}$
- g $-\frac{3}{2}x^{-\frac{3}{2}}$ h $-\frac{5}{3}x^{-\frac{4}{3}}$
- 3 > a $5x^4 2x^{-3}$
- $-x^{-3} + 8x$

h $-4x^{-5}$ i $-2x^{-3}$

- $\frac{1}{8}x^{-\frac{7}{8}} 6x^{-3} + 3x^{-\frac{2}{3}}$
- 4 > a $6x^5 + 2x + 2$
- **b** $3x^2$
- **5 a** $3 + \frac{2}{9}x^{-\frac{2}{3}}$ **b** $\frac{1}{2}x^{-\frac{5}{6}} + \frac{3}{2}x^{-\frac{1}{4}}$

 - c $8x^{-2} \frac{2}{3}x^{-\frac{5}{3}}$
- **6** ▶ **a** 6
- **b** $\frac{23}{2}$ **c** $\frac{513}{16}$
- **7 a** $\frac{4}{3}x^3 2x^2$
 - **b** $\frac{3}{2}x^{-\frac{1}{2}} \frac{1}{2}x^{-2}$
 - $-3x^{-2}$
- d $3 + 6x^{-2}$
- $= 3x^2 2x + 2$
- $\mathbf{f} = 24x 8 + 2x^{-2}$

- 1 **a** $12(3x + 1)^3$
- **b** $12(2x-4)^5$
- $c 6x(1 + x^2)^2$

- d $-35(1-5x)^6$ e $-3(2+3x)^{-2}$ f $3(6x+2)^{-\frac{1}{2}}$

- **g** $28x^3(x^4+3)^6$ **h** $-\frac{1}{2}(7-x)^{-\frac{1}{2}}$ **i** $-112(2-7x)^3$
- $54(7-3x)^{-4}$
- 2 > a 12e^{2x}
- **b** $-10e^{4-x}$ **c** $-15e^{-5x}$
- d $(2x-3)e^{x^2+3x}$ e $10e^{x^2+4x}(x+2)$
- $3 \triangleright a 2\cos(2x + 1)$
- $-4x\sin(2x^2+2)$
- $c -3\cos^2 x \sin x$
- d $8\sin 2x\cos 2x$
- $4 \triangleright -\frac{9}{256}$
- **5 a** y = 128x 240
- **b** y = 2x + 6

EXERCISE 3

- 1 ▶ a 3 cos x
- $b \sin x$
- $-6\cos 2x$
- d $2e^{2x}$
- $= 3\cos x + 6\sin 3x$
- $4e^{2x} + 4$
- $2 \triangleright a 6\cos 6x$
- **b** $\frac{3}{2}\sin{\frac{1}{2}x}$
- $c 4\cos 8x 24\sin 8x$
- d 12e4x
- $-\frac{3}{9}e^{3x}+15e^{5x}-48$
- **3** ▶ -3
- $4 > \frac{3}{2}$
- **5** ▶ -10

EXERCISE 4

- 1 > a $16x(1+4x)^3+(1+4x)^4$
 - **b** $24x^2(1+3x^2)+2(1+3x^2)^2$
 - $\frac{1}{2}x^2(x-2)^{-\frac{1}{2}}+(x-2)^{\frac{1}{2}}$
 - d $6x^4(5+x)^2+8x^3(5+x)^3$
 - $-20x^2(5x-2)^{-2}+8x(5x-2)^{-1}$
 - $\int_{0}^{1} \frac{3}{2}x^{3}(3x+1)^{-\frac{1}{2}} + 3x^{2}(3x+1)^{\frac{1}{2}}$
- 2 > a $(3x + 1)e^{3x}$
- $-2(x^2-x-4)e^{-2x}$
- $c 2e^{-3x}(1-3x)$
- d $4e^{x^3}(3x^3+1)$
- e $2(6x^2 + x + 3)e^{x^2}$ f $-(10x^2 + 4x 5)e^{-x^2}$
- $3 \triangleright a \cos x x \sin x$
- $\mathbf{b} \cos^3 x \sin 2x \sin x$
- c $4(\sin 6x + 6x \cos 6x)$ d $e^{x}(\cos x \sin x)$
- $e^{x}(\cos x + \sin x)$

- 4 **a** 1700
- **b** 14
- c 224

5 $y = \frac{8}{\sqrt{3}}x - \frac{5}{\sqrt{3}}$

EXERCISE 5

- 1 > a $-\frac{6}{(x-1)^2}$
- **b** $\frac{1}{(2x+1)^2}$ **c** $-\frac{6x(x+3)}{(2x-3)^4}$
- **d** $\frac{24x}{(x+2)^3}$ **e** $-\frac{3(x+4)}{8x^3\sqrt{x+3}}$ **f** $\frac{2x-9}{(4x+3)^3}$

- 2 > a $\frac{1-3x}{8^{3x}}$
- c $2e^{x^2} \frac{e^{x^2}}{x^2}$
- $3 \triangleright a \frac{x \sin x + \cos x}{x^2}$
- $e^x \sin x e^x \cos x$
- c $\frac{3x^2\sin x + \sin x + 6x\cos x}{\cos^2 x}$ d $\frac{2}{\sin x} \frac{(2x-1)\cos x}{\sin^2 x}$
- **4 a** $y = \frac{1}{4}x + 1$
- **b** $y = -\frac{x}{\pi} + 1$

EXERCISE 6

- **1 a** Tangent: y = -2x + 2
 - Normal: $y = \frac{1}{2}x = -\frac{11}{2}$
 - **b** Tangent: $y = 8x \frac{5}{2}$
 - Normal: $y = -\frac{1}{8}x + \frac{45}{8}$
 - **c** Tangent: $y = \frac{13}{2}x \frac{17}{2}$
 - Normal: $y = -\frac{2}{13}x \frac{24}{13}$
 - d Tangent: $y = 2e^2x e^2$
 - Normal: $y = -\frac{1}{2e^2}x + \frac{2e^4 + 1}{2e^2}$
 - e Tangent: $y = -\frac{3}{2}(x-1) + \frac{1}{2}$
 - Normal: $y = \frac{3}{2}(x-1) + \frac{1}{2}$

- **1 a** x = -6
- **b** $x = -\frac{3}{10}$
- $x = \frac{3}{9}$
- d −3 or 3
- $2 \triangleright a \quad x = -\frac{1}{6}$, minimum
 - **b** $x = \frac{7}{12}$, maximum
 - $x = \frac{1}{4}$, minimum
 - **d** x = 1, minimum and $x = -\frac{1}{3}$, maximum
 - e x = 1, minimum and x = -1 maximum
 - f x = 3, minimum

- 1 > 20 m × 40 m = 800 m²
- 2 ▶ 2000 π cm³
- $3 \triangleright \frac{800}{4 + \pi} \text{ cm}^2$
- **4 a** $y = 1 \frac{x}{2} \frac{\pi x}{4}$
 - b Proof
- 5 > 27 216 mm²

EXAM PRACTICE QUESTIONS

- 1 **a** $f'(x) = 3x^2 + 3$
- 2 $\Rightarrow \frac{d}{dx}(5x-7+\frac{2\sqrt{x}+3}{x})=\frac{-\sqrt{x}-3}{x^2}+5$
- **3 a** P = 16, Q = -24 **b** $= \frac{16x 9}{2x^{\frac{3}{2}}}$ **c** $\frac{5}{2}$
- **4 a** $6x^2 + 2kx$ **b** $k = \frac{41}{9}$ **c** P = 0.5

- **5 a** 7 **b** y = 3x + 1 **c** $\frac{1}{3}(2 + \sqrt{6})$
- **6 a** = 1 + $\frac{24}{3}$
- **b** y = 7x 29
- $7 > 15\cos 5x 12\sin 3x$
- $8 \rightarrow -\frac{1}{2}\sin(\frac{x}{2})$
- $9 \triangleright 2e^{2x} + 6x + 16\cos 4x$
- **10** ▶ -4
- **11** ▶ -8
- 12 > a $\frac{-3k}{(x-2k)^2}$
 - **b** Decreasing as $\frac{\text{negative}}{\text{positive}} = \text{negative}$
- **13** $\rightarrow y = 2x + 10$
- 14 ► 36e^{2x³}x²
- 15 \triangleright 12 sin 2x cos 2x
- **16 a** $-\frac{e^x(e^x+2)}{(e^x-2)^3}$
 - -34.6 (3 s.f.)
- $17 > 4x \cos 4x 3 \sin 4x$
- 18 \triangleright 2x cos 3x 3x² sin 3x
- **19** \triangleright 12 $x^3 + 45x^2 + 10x 5$
- **20** \triangleright y = -x
- **21 a** (4, 6)
 - **b** Maximum

- 22 > a Proof
- **b** k > 32
- 23 > a Proof
- **b** 40 cm

STARTER ACTIVITIES

- 1 > a i 2x 3
- ii 1 ii 5
- iv -3iv 1
- **Ⅲ** −5 iii -1
- **iii** 8
 - iv 0

 $d = 4x^3$

c i $6x^5 - 14x$ ii 164

b i 2x + 1

- ii 32
- **Ⅲ** −4
- iv 0 iv $\frac{1}{4}$
- e i $\frac{1}{4}$ iii $\frac{1}{4}$ **f** i -2x - 2 ii -6
- 0
- iv -2

- 1 > a $\frac{x^4}{4} + \frac{3x^2}{2} + c$
- **b** $x^4 x^2 + c$
- **c** $4x \frac{3}{x} + c$ **d** $-x^3 + \frac{10x^{\frac{3}{2}}}{2} + c$
- e $10x^3 + \frac{1}{2}x^{-2} + c$ f $\frac{x^3}{3} \frac{2x^{\frac{5}{2}}}{5} + 4x + 2\sqrt{x} + c$
- g $\frac{4}{3}x^4 + x^{-3} + rx + c$ h $\frac{p}{5}x^5 + 2tx 3x^{-1} + c$
- $-\frac{x}{2} \frac{3}{4x^2} + c$
- **2 a** $\frac{1}{4}x^4 + x^3 + c$ **b** $2x \frac{3}{x} + c$

 - **c** $\frac{4}{3}x^3 + 6x^2 + 9x + c$ **d** $\frac{4}{3}x^{\frac{3}{2}} + 6x^{\frac{1}{2}} + c$
- - e $\frac{3}{x} \frac{1}{8x^2} + c$ f $\frac{3x^{\frac{8}{3}}}{8} \frac{4}{3x^{\frac{2}{3}}} + c$
- **3 a** $2\sin x + c$ **b** $-\frac{2\cos 3x}{3} + c$

 - **c** $4e^x + c$ **d** $\frac{3e^{4x}}{4} + c$

 - $e = -3e^{-x} + c$ $f \frac{5e^{-3x}}{3} + c$
 - g $-2\cos\frac{1}{2}x + c$
- h $\frac{5}{3}$ sin 3x + c
- $-\frac{4}{3}\sin 3x + c$
- 4 > a $2(\sin x + \cos x) + 3x^2 + c$
 - **b** $-3\cos x + 7e^x + \frac{2}{x} + c$
 - c $2\sin x \cos x + e^x + c$
 - d $-\cos 2x + \frac{3e^{4x}}{4} + x^3 + c$
 - **e** $4(\sin x + e^x + \frac{x^3}{3}) + c$

- v = 20t
- v = 48 32t
- **a** 30 10t
- b 0 m s 1
- $a = -32 \, \text{ms}^{-2}$
- i $s = 384 + 160t 16t^2$ ii t = 12
- **5 a** a = 6 2t
- $b 2 m s^{-2}$
- c 57.3 m

EXERCISE 3

- 1 \triangleright a $\frac{4}{3}$ square units
- **b** $20\frac{5}{6}$ square units
- c $20\frac{5}{6}$ square units
- d $21\frac{1}{12}$ square units
- 2 > a 33 square units
- b 1.5 square units
- $\frac{9}{4}$ square units
- d 6 square units
- e $8\frac{5}{12}$ square units
- 3 ▶ a $\frac{4}{3}$ square units
- **b** $10\frac{2}{3}$ square units
- $\frac{1}{6}$ square units
- d $5\frac{1}{3}$ square units
- e $\frac{9}{2}$ square units
- 4 ▶ a 7

b 0.5

c 2

- d 32
- $5 \triangleright 20\frac{5}{6}$ square units
- 6 > 3 square units
- **7** ▶ **a** (-3, 9) (3, 9)
- b 36 square units
- 8 **a** (0, 1) (3, 4)
- $\frac{9}{2}$ square units
- $9 \triangleright \frac{59}{6}$ square units
- 10 ▶ a (2, 12)
- b Proof

11 Proof

EXERCISE 4

- 1 $\triangleright \frac{32}{5}\pi$ cubed units
- $\frac{2}{7} = \frac{128}{7} \pi$ cubed units
- $3 \triangleright \frac{198}{7} \pi$ cubed units
- 4 > a i 2 ln 2
- 112π
- **b** i 2ln2 1
- ii $[2(\ln 2)^2 4 \ln 2 + 2]\pi$

c i $\frac{8}{3}$

- $\frac{64}{15}\pi$
- 5 $\blacktriangleright \frac{\pi^2}{4}$ cubed units

EXERCISE 5

- 3 > 15e²
- **5** ▶ 2%
- 6 ▶ 2%
- $7 \triangleright \frac{dM}{dt} = km$
- 8 Proof
- 9 Proof
- **10** $\triangleright \frac{dh}{dt} = \frac{18}{\pi h^2}$

EXAM PRACTICE QUESTIONS

- 1 > $y = \frac{1}{5}x^5 + 2x^3 + 9x \frac{548}{5}$
- **2** $y = -\frac{1}{4}x^4 2x^{-1} + \frac{5}{4}x^{-2} + 8$
- $3 \rightarrow \frac{5x^4}{4} + \frac{5}{x} + c$
- 4 $\triangleright \frac{2\sqrt{x}(x+27)}{3} + c$
- 5 > $-4 \sin x + 5 e^x + \frac{x^4}{2} + c$
- 6 ► $12\frac{2}{3}$ square units
- 7 **a** (2, 8) and (9, 1)
- **b** $57\frac{1}{6}$ square units
- **8 a** x = -1 and x = 4 **b** $\frac{125}{6}$ square units
- 9 > 37.7 (3 s.f.)
- 10 $\rightarrow \frac{148}{3}$ square units
- 11 $\triangleright \frac{11}{6}$ square units
- 12 \triangleright $\frac{243}{2}$ square units
- 13 $\triangleright \frac{128}{7} \pi$ cubed units
- 14 $\triangleright \frac{23}{14}\pi$ cubed units
- 15 $\triangleright \frac{146}{5} \pi$ cubed units

16
$$\blacktriangleright \frac{\mathrm{d}x}{\mathrm{d}t} = \frac{\mathrm{d}x}{\mathrm{d}V} \times \frac{\mathrm{d}V}{\mathrm{d}t} = \frac{0.048}{3x^2}$$

At
$$x = 8$$

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{0.048}{3(8^2)} = 0.00025 \; \mathrm{cm} \, \mathrm{s}^{-1}$$

17 For a sphere:
$$V = \frac{4}{3}\pi r^3 = \frac{\mathrm{d}v}{\mathrm{d}r} = 4\pi r^2$$

$$\delta r = 0.02\,\mathrm{cm}$$

$$\delta v = 4\pi r^2 \, \delta r$$

Use
$$r = 1 \Rightarrow$$

$$\delta v = 4\pi \times 0.02$$
$$= 0.25 \,\mathrm{cm}^3$$

STARTER ACTIVITIES

- 1 > Sin
 - Opposite
 - Hypotenuse
 - Cosine
 - Adjacent
 - Hypotenuse
 - Tan
 - Opposite
 - Adjacent
- 2 > 1500 m
- 3 ▶ 41.2 m

EXERCISE 1

- 1 > a 18°
- b 9°
- c 75°

- d 90°
- e 180°
- f 540°

- 900°
- h 140°
- 225°

- j 270°
- 2 > a 32.7°
- **b** 57.3°
- c 114.6°

- d 78.5°
- e 81.0°
- f 179.9°

- g 257.3°
- **3** ▶ **a** 0.247
- **b** 6.334
- c 0.847

- d 0.998
- e 0.058

18

f 0.0585

- 45
- 8

- π 6
- 5π 8
- 2π 3

- 3π 4
- 4π 3
- 3π 2

- 7π
- 11π
- 5 **a** 0.873
- **b** 1.31
- c 1.75

- d 2.09
- e 4.10
- f 5.67

EXERCISE 2

- 1 > a i 3.6 cm
- ii 25.92 cm
- 15.4 cm
- b i 20 cm
- ii 3.34 cm
- iii 6cm
- - ii 0.978 rad
- iii 2.45 rad
- c i 1.19 rad d i 13.8 cm²
- ii 28.9 cm²
- iii 169 cm²
- 2 > a Area = 24.3 cm² and arc length = 5.4 cm
 - **b** Area = $6.73 \, \text{cm}^2$ and arc length = $2.01 \, \text{cm}$
 - c Area = 88.1 cm² and arc length = 26.3 cm
- 3 > a 5.6 cm²
- b 4.87 cm²
- $4 > \frac{16}{3} \pi \text{cm}^2$
- 5 > a 4.13 radians
- 45.5 cm
- 6 Area = 39.6 cm² and perimeter = 25.2 cm
- 7 > a Perimeter = 10 cm
- b Area = $6 \, \text{cm}^2$
- 8 > a Perimeter = 20.4 cm
- b Area = 23.7 cm²
- 9 > a $r = 20 \, \text{cm}$
- b Proof
- c 0.519 cm²

- 10 > 42.3 cm²
- **11** \triangleright a (R-r) cm
- b Proof
- c 5.68

- 12 ▶ a Proof
- **b** 34.1 m²
- 13 ▶ a Proof
 - **b** i 80.9 m
- ii 26.7 m
- iii 847 m²

EXERCISE 3

- √2

- $m\sqrt{3}$
- $\frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}$

- **1 a** x = 37.7, y = 86.3, z = 6.86
 - **b** x = 48, y = 19.5, z = 14.6

 - x = 30, y = 11.5, z = 11.5
 - **d** x = 21.0, y = 29.0, z = 8.09x = 93.8, y = 56.3, z = 29.9
 - x = 97.2, y = 41.4, z = 41.4
 - $\mathbf{g} \ \ x = 45.3, \ y = 94.7, \ z = 14.7$
 - or x = 134.7, y = 5.27, z = 1.36

- **h** x = 7.07, y = 73.7, z = 61.2or x = 7.07, y = 106, z = 28.7
- x = 49.8, y = 9.39, z = 37.0
- 2 > a 23.7 cm²
- b 4.31 cm²
- c 20.2 cm²

- 3 > 42.4 cm²
- 4 Proof
- 5 > 29.8 cm
- 6 ▶ Proof
- 7 Proof
- 8 > a x = 4
- 6.50 cm²

- 1 > a 11.3 cm
- **b** 288.7 m
- c 51.1°
- d 32.2 cm

- 2 > a 10.4 cm
- b 35.2°
- 3 > a 13.4 cm
- b 17.3°

- **4** ▶ 22.2
- 5 > a 7.5 cm

- b 143°
- 6 > a 195.8 m
- **b** 325 m

EXERCISE 6

- 1 \triangleright a $\cos^2\theta$
- b $\sin^2\left(\frac{\theta}{2}\right)$
- **c** 6

- $d \sin^2 A$
- $e \cos 4\theta$
- f $tan \theta$

- α tan 3A
- h 4
- $i sin^2x$

- i 1
- 2 > a Proof
- **b** Proof
- Proof

- d Proof
- Proof
- f Proof

- g Proof
- h Proof
- 3 ▶ a sin 35°
- b cos 210°
- c tan31° f tan 5x

- $d \cos 3x$ $\mathbf{g} \sin A$
- $e \cos 7\theta$ $h \cos 3x$
- 4 ▶ a 1
- **b** 0
- $c \frac{\sqrt{2}}{2}$

- d √3
- f 1

- 5 > a Proof
- **b** Proof
- Proof

EXERCISE 7

- 1 > a 210°, 330°
 - c 180°

 - e 140°, 220°
 - g 90°, 270°

- **b** 60°, 240°
- d 15°, 165°
- f 45°, 225°
- h 230°, 310°
- i 45.6°, 134.4°
- j 135°, 225°

- 2 ▶ a −120°, −60°
 - c -144°, 144°
- **b** -171.37°, -8.63° d -327.1°, -32.9°
- e 30°, 210°, 390°, 570°
- f 251°, 431°
- $\mathbf{g} \pi, 0, \pi, 2\pi, -2\pi$
- $-\frac{7\pi}{4}, -\frac{5\pi}{4}, \frac{\pi}{4}, \frac{3\pi}{4}$
- $-\frac{4\pi}{3}, -\frac{2\pi}{3}, \frac{2\pi}{3}$
- i −6.34°, 173.7°, 353.7°
- 3 > a 0°, 45°, 90°, 135°, 180°, 225°, 270°, 315°, 360°
 - **b** 60°, 180°, 300°
- c $22\frac{1}{2}^{\circ}$, $112\frac{1}{2}^{\circ}$, $202\frac{1}{2}^{\circ}$, $292\frac{1}{2}^{\circ}$

d 300°

- e 225°, 315°
- f 50°, 170°
- g 165°, 345°
- **4 a** -2π , $-\pi$, 0, π , 2π
- **b** 0. 2π
- **5 a** $x = -40^{\circ}, 0^{\circ}, 80^{\circ}$
 - **b** 4.78°, 6.22°
 - c 113°, 158°, 203°, 248°
 - d $-\frac{11}{12}\pi$, $-\frac{7}{12}\pi$, $\frac{\pi}{12}$, $\frac{5}{12}\pi$
 - $-\frac{29}{48}\pi$, $-\frac{13}{48}\pi$, $\frac{19}{48}\pi$, $-\frac{35}{48}\pi$

EXERCISE 8

- 1 > a 30°, 210°
- b 135°, 315°
- **2 0.59**, 3.73
- 3 > a 60°, 120°, 240°, 300°
 - **b** 60°, 300°
 - c 270°
 - d 194°, 270°, 346°
- $\frac{5\pi}{12}$, $\frac{11\pi}{12}$, $\frac{17\pi}{12}$, $\frac{23\pi}{12}$ **b** 0.841, $\frac{2\pi}{3}$, $\frac{4\pi}{3}$, 5.44

 - **c** 4.01, 5.41

EXAM PRACTICE QUESTIONS

- 1 > a 17.1 cm²
- **b** 5.7 cm
- c 11.7 cm
- d 4.5 cm²
- 2 > a Proof
- **b** 2.04 rad **c** 61 cm²
- 3 ▶ a 94°
- b 27.9 cm²
- 4 Proof
- **5** \triangleright a tan $\theta = 5$
- **b** 78.7°, 258.7°

- $6 > \frac{\pi}{6}, \frac{5\pi}{6}$
- 7 \triangleright a $\sin^4 3\theta$
- **b** 1


8 ▶ a 1

 $b \tan y = \frac{4 + \tan x}{2 \tan x - 3}$

9 a
$$x = \frac{23\pi}{12}, x = \frac{11\pi}{12}$$

b
$$x = \frac{2\pi}{3}, x = \frac{5\pi}{6}, x = \frac{11\pi}{6}, x = \frac{5\pi}{3}$$

- **10** ▶ 23.8°, 203.8°
- 11 ▶ Proof




- **13** > 40.9°, 139.1°, 220.9°, 319.1°
- 14 ▶ Proof

INDEX


| A | $\cos x$ | F |
|---|---|--------------------------------------|
| acceleration 164 | differentiation 141-3 | factor theorem for polynomials 52-4 |
| Al-Karaji 97 | graph 190–3 | Fibonacci sequence 75 |
| angles, in radians 141, 184–5 | identities 199-202 | fractions, rationalise the |
| arc length 185–90 | integration 162 | denominator 5 |
| area | cosine rule 193-6 | functions |
| circles 185–90 | 3D problems 197-9 | modulus 101 |
| under a curve 166-71 | cubic functions, sketching 62-6 | periodic 191 |
| sectors 187-90 | curves | reciprocal 65-6 |
| triangles 193-6 | area under 166-71 | solving equations by graphs |
| arithmetic sequences, identifying | gradient 138-41 | 12–14 |
| 76–7 | stationary points 151-5 | • |
| arithmetic series | tangent and normal 149-50 | G |
| nth term 78-7 | volume of revolution 171–3 | Gauss, Carl Friedrich 80 |
| sum to n terms 80-3 | D | geometric sequences |
| asymptotes 7 | D | common ratio 84, 85-7 |
| ,, | definite integral 166 | identifying 84 |
| В | derived functions 138 | nth term 85–7 |
| bearings 107 | difference of two squares 25 | geometric series |
| binomial expansion | differential equations 174-7 | sum to infinity 90-2 |
| coefficients 98–101 | differentiation | sum to n terms 87-90 |
| Pascal's triangle 97–8 | chain rule 143–6, 175 | gradient |
| true and valid range 101–3 | differential equations 174-7 | curves 138-41, 151-5 |
| binomial series 97–8 | exponential function 141-3 | perpendicular lines 128-30 |
| binomial theorem 98–9 | gradient function 138-41 | straight line 125-8 |
| binomial theorem 30–3 | kinematic problems 164-5 | tangents to a curve 149-50 |
| C | maximum and minimum points | graphs |
| | 151–5 | basic principles 61 |
| chain rule 143–5, 175 circles | multiple powers 138–41 | cubic functions 62-6 |
| arc length and area 185–90 | problem solving 151–5 | exponential functions 9-11 |
| area of sectors 187–90 | product rule 145-7 | of functions to solve equations |
| | quotient rule 147-8 | 12–14 |
| coefficients, binomial expansion 98–101 | rates of change 174-7 | linear inequalities 44-9 |
| column matrix 117 | second derivative 164 | logarithmic functions 7-9 |
| | $\sin ax$ and $\cos ax$ 141–3 | quadratic inequalities 41-3 |
| common difference 76, 78–80 common ratio 84, 85–7 | stationary points 151–5 | reciprocal functions 65-6 |
| | tangent and normal 149–50 | $\sin x$, $\cos x$, $\tan x$ 190–3 |
| completing the square 27–8 compound interest 10 | discriminant 29–31 | u . |
| constant of integration 162 | displacement 164-5 | Н |
| convergent geometric series 90–2 | distance between two points 131 | hemispheres, volume of 175 |
| coordinate geometry | division of polynomials 50-2 | 1 |
| distance between two points 131 | E | indiana lawa of 15 |
| equation of a straight line 127–8 | | indices, laws of 15 |
| gradient of straight line 127–8 | e (base of natural logarithms) 10 exponential functions | inequalities |
| perpendicular lines 128–30 | differentiation 141–3 | linear 39–41 |
| point that divides line in a given | | modulus function 101 |
| ratio 132 | graphs of 9–11 integration 162 | quadratic 41–3 infinite series 90–2 |
| I GLIO I OE | IIIEUIAUUII 107 | |

| integration | different functions, solving | stationary points 151-5 |
|-------------------------------------|-----------------------------------|---|
| area under a curve 166-71 | equations 67-8 | Stifel, Michael 97 |
| constant of integration 162 | reciprocal functions 65-6 | straight line |
| exponential function 162 | transformations 68-9 | coordinates of a point that divides |
| kinematic problems 164-5 | position vectors 113-14 | line in a given ratio 132 |
| kinematics 164-5 | product rule 145-7 | distance between two points 131 |
| as reverse of differentiation 162-4 | | equation 125, 127-8 |
| rules of 162 | Q | gradient 125-6 |
| $\sin x$, $\cos x$ 162 | quadratic equations | perpendicular lines 128-30 |
| volume of revolution 171-3 | α and β roots 31–3 | surds 1-6 |
| irrational numbers 4 | basic principles 25 | |
| | completing the square 27-8 | T |
| K | discriminant 29-31 | tan x |
| Khayyam, Omar 97 | factorising 26 | graph 190-3 |
| | one linear one quadratic 38-9 | identities 199–202 |
| L | quadratic formula 28-31 | ratios 191 |
| linear inequalities 39–41 | real and unreal roots 29-31 | tangent, to a curve 149–50 |
| graphs 44–9 | quadratic inequalities 41-3 | transformations 68–9 |
| | quotient rule 147-8 | triangles, area of 193-6 |
| Linear Programming 47–8 | quotients 50 | trigonometry |
| logarithmic functions 7–9 | | basic principles 183 |
| logarithms | R | cosine rule 193–9 |
| change the base 17–18 | radians 141, 185-5 | formulae to solve equations |
| laws of 15–17 | rates of change 174-7 | 206–7 |
| natural logarithm 10 | rationalise the denominator 5 | identities 199–202 |
| solving equations 18–20 | reciprocal functions, graphs 65-6 | radians 141, 184–5 |
| writing expressions as 14-15 | remainder theorem 53-5 | sine rule 193–9 |
| | | |
| M | S | $\sin x$, $\cos x$, $\tan x$ graphs 190–3 |
| mathematical modelling 37 | second derivative 164 | solve trigonometric equations 202-5 |
| maximum and minimum points | sectors, area of 185-90 | |
| 151–5 | sequences | turning points 151-5 |
| modulus function 101 | arithmetic 75-83 | V |
| | definition 75 | • |
| N | geometric 84, 85-7 | vectors |
| natural logarithm 10 | series | addition and subtraction 110–13 |
| Newton, Sir Isaac 97, 137 | definition 75 | basic principles 107 |
| Newton's Law of Cooling 174 | nth term 78-80 | Cartesian components 117–19 |
| normal to a curve 149-50 | sum 87-90 | modulus 109 |
| | sum to infinity 90-2 | notation and diagrams 108-10 |
| P | simultaneous equations | position of a point 113-14 |
| Pascal's triangle 97-8 | basic principles 37 | position vectors 113-14 |
| periodic functions 191 | one linear one quadratic 38-9 | properties of geometrical figures |
| perpendicular lines 128-30 | sine rule 193–6 | 114–17 |
| point of inflexion 64 | 3D problems 197-9 | unit vector 110-13 |
| polynomials | $\sin x$ | velocity 164–5 |
| division 50–2 | differentiation 141–3 | volume |
| factorising 52–4 | graph 190–3 | hemispheres 175 |
| remainder theorem 53–5 | identities 199–202 | of revolution 171–3 |
| polynomials, sketching 61–6 | integration 162 | W. |
| cubic functions 62–6 | sound 183 | Υ |
| | square roots 1–6 | Yang Hui 97 |

square roots 1-6



| | | | | | |
|-------|--|---|------|---|---|
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| +++++ | | | | | |
| | | | | | |
| | | | | | |
| | | +++++++++++++++++++++++++++++++++++++++ | | | +++++++++++++++++++++++++++++++++++++++ |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | + |
| | | | | | |
| | | | | | |
| | | | | | |
| ++++ | | | | | +++++++++++++++++++++++++++++++++++++++ |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |



| | | + |
|-------------|--|---------|
| | | \Box |
| | | |
| | | \perp |
| | | |
| | | +++ |
| | | |
| | | ш |
| | | |
| | | |
| | | + |
| | | +++ |
| | ╒┋╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒ | ++ |
| | | |
| | | |
| | | |
| | | +++ |
| | | +++ |
| | ╒┋╘┊╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒ | + |
| | | |
| | | |
| | | ш |
| | | ++ |
| | _ | ++ |
| | | |
| | | |
| | | |
| | | + |
| | | H |
| | ╎╂┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼ | + |
| | | |
| | | |
| | | |
| | | + |
| | , | + |
| | <u>┍╊╒┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼</u> | H |
| | | |
| | | \Box |
| | | H |
| | , | + |
| | | + |
| | | |
| | | |
| | | ш |
| | | ++ |
| | | + |
| | | + |
| | | |
| | | |
| | | + |
| | | - |
| | | \Box |
| | | |
| | | ш |
| | | + |
| | | +++ |
| | | \Box |
| | | |
| | | |
| | | +++ |
| | ╒┋╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒ | ++ |
| | | |
| | | |
| | | |
| | | + |
| | ╒┋╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒╒ | + |
| | | |
| | | |
| | | |
| | , | + |
| | <u>┍╊╒┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼</u> | |
| | | |
| | | Ш |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | + |
| | | + |
| | | + |
| | | + |
| | | + |
| | | + |

| | +++++++++++++++++++++++++++++++++++++++ | | |
|---|---|---|---|
| | | | |
| | | | |
| | | | |
| | | | |
| | +++++++++++++++++++++++++++++++++++++++ | | +++++++++++++++++++++++++++++++++++++++ |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |