

Mark Scheme (Results)

Summer 2024

Pearson Edexcel Adanced Level GCE In Physics (9PH0) Paper 2: Advanced Physics II

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson. Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2024

Question Paper Log Number: P74469A Publications Code: 9PH0_02_2406_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
 - i) ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
 - ii) select and use a form and style of writing appropriate to purpose and to complex subject matter
 - iii) organise information clearly and coherently, using specialist vocabulary when appropriate.

Mark scheme notes

Underlying principle

The mark scheme will clearly indicate the concept that is being rewarded, backed up by examples. It is not a set of model answers.

1. Mark scheme format

- 1.1 You will not see 'wtte' (words to that effect). Alternative correct wording should be credited in every answer unless the MS has specified specific words that must be present. Such words will be indicated by underlining e.g. 'resonance'
- 1.2 Bold lower case will be used for emphasis e.g. 'and' when two pieces of information are needed for 1 mark.
- 1.3 Round brackets () indicate words that are not essential e.g. "(hence) distance is increased".
- 1.4 Square brackets [] indicate advice to examiners or examples e.g. [Do not accept gravity] [ecf].

2. Unit error penalties

- 2.1 A separate mark is not usually given for a unit but a missing or incorrect unit will normally mean that the final calculation mark will not be awarded.
- 2.2 This does not apply in 'show that' questions or in any other question where the units to be used have been given, for example in a spreadsheet.
- 2.3 The mark will not be awarded for the same missing or incorrect unit only once within one clip in epen.
- 2.4 Occasionally, it may be decided not to insist on a unit e.g the candidate may be calculating the gradient of a graph, resulting in a unit that is not one that should be known and is complex.
- 2.5 The mark scheme will indicate if no unit error is to be applied by means of [no ue].

3. Significant figures

- 3.1 Use of too many significant figures in the theory questions will not be prevent a mark being awarded if the answer given rounds to the answer in the MS.
- 3.2 Too few significant figures will mean that the final mark cannot be awarded in 'show that' questions where one more significant figure than the value in the question is needed for the candidate to demonstrate the validity of the given answer.
- 3.3 The use of one significant figure might be inappropriate in the context of the question e.g. reading a value off a graph. If this is the case, there will be a clear indication in the MS.

- 3.4 The use of $g = 10 \text{ m s}^{-2}$ or 10 N kg⁻¹ instead of 9.81 m s⁻² or 9.81 N kg⁻¹ will mean that one mark will not be awarded. (but not more than once per clip). Accept 9.8 m s⁻² or 9.8 N kg⁻¹
- 3.5 In questions assessing practical skills, a specific number of significant figures will be required e.g. determining a constant from the gradient of a graph or in uncertainty calculations. The MS will clearly identify the number of significant figures required.

4. Calculations

- 4.1 Bald (i.e. no working shown) correct answers score full marks unless in a 'show that' question.
- 4.2 If a 'show that' question is worth 2 marks. then both marks will be available for a reverse working; if it is worth 3 marks then only 2 will be available.
- 4.3 **use** of the formula means that the candidate demonstrates substitution of physically correct values, although there may be conversion errors e.g. power of 10 error.
- 4.4 **recall** of the correct formula will be awarded when the formula is seen or implied by substitution.
- 4.5 The mark scheme will show a correctly worked answer for illustration only.

1. Graphs

- 1.1 A mark given for axes requires both axes to be labelled with quantities and units, and drawn the correct way round.
- 1.2 Sometimes a separate mark will be given for units or for each axis if the units are complex. This will be indicated on the mark scheme.
- 1.3 A mark given for choosing a scale requires that the chosen scale allows all points to be plotted, spreads plotted points over more than half of each axis and is not an awkward scale e.g. multiples of 3, 7 etc.
- 1.4 Points should be plotted to within 1 mm.
 - Check the two points furthest from the best line. If both OK award mark.
 - If either is 2 mm out do not award mark.
 - If both are 1 mm out do not award mark.
 - If either is 1 mm out then check another two and award mark if both of these OK, otherwise no mark.
 - For a line mark there must be a thin continuous line which is the best-fit line for the candidate's results.

Question Number	Answer	Mark				
1	The only correct answer is B					
	A is not correct because V , X and Z are all on the main sequence					
	C is not correct because Z is on the main sequence and the sun will become a white dwarf, which is position Y					
	\boldsymbol{D} is not correct because Z, X and V are all on the main sequence					
2	The only correct answer is B	1				
	A is not correct because gravitational force is equal to centripetal force, so the velocity is independent of the mass of the satellite					
	C is not correct because gravitational force is equal to centripetal force, so the velocity is independent of the mass of the satellite					
	D is not correct because gravitational force is equal to centripetal force, so the velocity is independent of the mass of the satellite					
3	The only correct answer is C	1				
	A is not correct because the atomic mass number must decrease by 16, which corresponds to 4 alpha decays					
	B is not correct because the atomic mass number must decrease by 16, which corresponds to 4 alpha decays					
	D is not correct because the atomic number would decrease by 8 due to alpha decay, then increase by 6 due to beta decay					
4	The only correct answer is B	1				
	A is not correct because this produces a beam of electrons with a shorter de Broglie wavelength, which produces a smaller angle					
	of diffraction					
	$\hat{m{C}}$ is not correct because increasing the filament temperature would have no effect on the wavelength of the electron beam					
	D is not correct because diffraction is maximised when the wavelength is similar to the gap size, increasing the gap size would					
	decrease the angle of diffraction					
5	The only correct answer is D	1				
	A is not correct because gravitational field strength is proportional to $1/r^2$					
	B is not correct because electric field strength is proportional to $1/r^2$					
	C is not correct because gravitational potential is always negative					
6	The only correct answer is C	1				
	A is not correct because smaller nuclei do not undergo fission					
	B is not correct because larger nuclei do not undergo fusion					
	D is not correct because larger nuclei do not undergo fusion					
7	The only correct answer is A	1				
	B is not correct because ray z has increased in speed, as shown by refracting away from the normal					
	C is not correct because ray z has increased in speed, as shown by refracting away from the normal					
	D is not correct because in medium y the ray has slowed down as it has refracted towards the normal					

8	The only correct answer is D	1
	A is not correct because elastic strain energy is equal to $\frac{1}{2}F^{2}/k$	
	B is not correct because elastic strain energy is equal to $\frac{1}{2}F^2/k$	
	C is not correct because elastic strain energy is equal to $\frac{1}{2}F^{2}/k$	
9	The only correct answer is B	1
	A is not correct because the initial gradient is not zero. The acceleration at the start is zero, so the velocity graph must have an	
	initial gradient of zero	
	C is not correct because the initial gradient is not zero	
	D is not correct because the velocity in the first quarter cycle is decreasing	
10	The only correct answer is D	1
	A is not correct because doubling the force and the length whilst halving the diameter leads to an increase by a factor of 16	
	B is not correct because doubling the force and the length whilst halving the diameter leads to an increase by a factor of 16	
	C is not correct because doubling the force and the length whilst halving the diameter leads to an increase by a factor of 16	

(Total for Multiple Choice Questions = 10 marks)

Question Number	Acceptable answers	Additional guidance	Mark
11	• Use of $n\lambda = d\sin\theta$ (1)	Example of calculation $3 \lambda = d \times 1.67 \times 10^{-6} \text{ m} \times \sin 73.3^{\circ}$	
	• $\lambda = 5.33 \times 10^{-7} \mathrm{m}$ (1)	$\lambda = 5.33 \times 10^{-7} \mathrm{m}$	2

Total for Question 11 = 2 marks

Question Number	Acceptable answers	Additional guidance		
12(a)(i)	The (minimum) frequency at/above which electrons are released	(1)	Accept frequency below which electrons are not released	1
12(a)(ii)	An explanation that makes reference to the following points:			
	With waves, energy would build up	(1)		
	A photoelectron would (eventually have enough energy to) be emitted with any frequency, (which is not observed)	(1)		2
12(b)	A <u>quantum</u> of (electromagnetic) energy/radiation Or A <u>discrete</u> packet of (electromagnetic) energy/radiation	(1)	Accept light for energy/radiation	1

(Total for Question 12 = 4 marks)

Question Number	Acceptable answers	Additional guidance	Mark
13	• Use of $\lambda_{\text{max}} T = 2.898 \times 10^{-3} \text{ m K}$ (1)	Example of calculation $7.43 \times 10^{-7} \text{ m} \times T = 2.898 \times 10^{-3} \text{ m K}$	
	• Use of $A = 4\pi r^2$ (1)	T = 3900 K	
	• Use of $L = \sigma A T^4$ (1)	$A = 4\pi (3.14 \times 10^{10} \mathrm{m})^2$ = 1.24 × 10 ²² m ²	
	• $L = 1.63 \times 10^{29} \mathrm{W}$ (1)	$L = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4} \times 1.24 \times 10^{22} \text{ m}^{2} \times (3900 \text{ K})^{4}$ $L = 1.63 \times 10^{29} \text{ W}$	4

(Total for Question 13 = 4 marks)

Question Number	Acceptable answers		Additional guidance	Mark
14(a)	• Use of $1/f = P$	(1)	Example of calculation $P_1 = 1/0.1 \text{ m} = 10 \text{ D}$	
	• Use of $P = P_1 + P_2$	(1)	$P_2 = 0.15 \text{ m} = 6.7 \text{ D}$ P = 10 D + 6.7 D	
	• $P = 17 (D)$	(1)	= 16.7 D	3
			P = 1/0.1 m + 1/0.15 m for MP1 and MP2	
14(b)	• Use of $1/v + 1/u = 1/f$ Or $1/v + 1/u = P$	(1)	Example of calculation $1/v + 1/(0.04 \text{ m}) = 16.7 \text{ D}$	
	• $v = (-)$ 12 cm (ecf from a)	(1)	v = -0.12 m	2
			(Show that answer gives 20 cm)	
14(c)	• Use of $M = v/u$	(1)	Example of calculation $M = 12 \text{ cm} / 4 \text{ cm} = 3$	
	• Use of $M = (image height) / (object height)$	(1)	$I = 3 \times 1.4 \text{ cm} = 4.2 \text{ cm}$	
	• Image height = 4.2 cm (ecf from b)	(1)	(Show that from (a) gives 7.0 cm)	3
14(d)	Virtual and erect/upright	(1)		1

(Total for Question 14 = 9 marks)

Question Number	Acceptable answers		Additional guidance	Mark
15(a)	EITHER	Accept answer in terms of multiple 'loops'		
	Waves are produced by the toothbrush and reflected (at the mass)	(1)		
	• Waves meet and superpose	(1)	Accept reference to phase difference in terms of	
	 Constructive interference takes place where the waves are in phase Or 	(1)	2π , or 0 (radians)	
	Destructive interference takes place where the waves are in antiphase		Accept reference to phase difference in terms of π (radians)	
	 The string vibrates at the centre with maximum amplitude, forming an antinode in the centre Or 	(1)	Do not credit out of phase	
	The string here has zero/minimum amplitude, forming nodes at the ends			
	Creating a standing/stationary wave	(1)		
	OR			
	 The toothbrush causes the string to vibrate at its natural frequency 	(1)		
	• There is a maximum/efficient transfer of energy from the toothbrush to the string	(1)		
	Resonance occurs	(1)		5
	• The string vibrates at the centre with maximum amplitude, forming an antinode in the centre	(1)		
	Creating a standing/stationary wave	(1)		

15(b)	• Use of $W = mg$	(1) Example of calculation $T = 0.010 \text{ kg} \times 9.81 \text{ N kg}^{-1} = 0.0981 \text{ N}$
	• Use of $v = \sqrt{T/\mu}$	(1) $ v = \sqrt{(0.0981 \text{ N} / 9.1 \times 10^{-4} \text{ kg m}^{-1})} = 10.4 \text{ m s}^{-1} $ $ \lambda = 0.69 \text{ m} / 2 = 0.345 \text{ m} $
	Determination of wavelength	(1) $ \begin{cases} \lambda = 0.69 \text{ m} / 2 = 0.343 \text{ m} \\ f = 10.4 \text{ m s}^{-1} / 0.345 \text{ m} \\ = 30 \text{ Hz} \end{cases} $
	• Use of $v = f\lambda$	$(1) \qquad -30 \text{ Hz}$
	• $f = 30 \text{ Hz}$	(1)

(Total for Question 15 = 10 marks)

Question Number	Acceptable answers	Additional guidance	Mark
16(a)	• Use of $f = 1/T$	Example of calculation $T = 1 / 0.34 \text{ Hz} = 2.94 \text{ s}$	
	• Use of $T = 2\pi \sqrt{(m/k)}$	1) $2.94 \text{ s} = 2\pi \sqrt{(m/350 \text{ N m}^{-1})}$ m = 76.6 kg	
	Correct for mass of platform	mass of astronaut = $76.6 \text{ kg} - 5.7 \text{ kg}$ = 70.9 kg	
	• Mass of astronaut = 71 kg		4
16(b)(i)	Determine amplitude	Example of calculation Amplitude = $0.29 \text{ m} / 2 = 0.145 \text{ m}$	
	• Use of $\omega = 2\pi f$	1) $\omega = 2\pi \times 0.34 \text{ Hz} = 2.14 \text{ rad s}^{-1}$ $\max a = (2.14 \text{ rad s}^{-1})^2 \times 0.145 \text{ m}$	
	• Use of max $a = (-) \omega^2 A$	$= 0.66 \text{ m s}^{-2}$	
	• Max acceleration = 0.66 m s^{-2}		4
16(b)(ii)	• Use of $v = (-) A\omega \sin \omega t$	Example of calculation $v = 0.145 \times 2.14 \text{ s}^{-1} \sin ((2.14 \text{ s}^{-1} \times 3.5 \text{ s}) \text{ radian})$	
	• Speed = 0.29 m s ⁻¹ (ecf for ω and A from (b)(i))	1) $ = 0.145 \times 2.14 \text{ s}^{-1} \sin ((2.14 \text{ s}^{-1} \times 3.5 \text{ s}) \times 360^{\circ} / 2\pi) $ $ \text{Speed} = 0.29 \text{ m s}^{-1} $	2

(Total for Question 16 = 10 marks)

II O II NIM	Acceptable answers			
Use of $pV = NkT$	(1)	Example of calculation $p_1 V_1 = NkT_1$		
Conversion of T in K	(1)	$ \begin{aligned} p_1 \ V_1 / T_1 &= NkT_1 = p_2 \ V_2 / T_2 \\ V_1 / 293 \ K &= 2800 \ m^3 / 393 \ K \end{aligned} $		
Use of $\rho = m/V$ to determine mass of air in the balloon	(1)	Volume of gas before heating, $V_1 = 2087 \text{ m}^3$ mass of air in balloon		
Calculation of total mass = mass of air at 120 °C + passengers + balloon	(1)	=2505 kg		
Use of $W = mg$	(1)	$= (2505 + 340 + 380) \text{ kg} = 3225 \text{ kg}$ $W = 3225 \text{ kg} \times 9.81 \text{ N kg}^{-1} = 31 637 \text{ N}$		
$W = 31\ 600\ \text{N}$, which is less than 33 000 N, so the balloon can take off	(1)	31 600 N < 33 000 N	6	
Any one from: Point particles Particles have negligible volume (compared to their container) Negligible forces between particles (except during a collision) Range of molecular forces small (in comparison to average separation) Duration of collisions negligible (in comparison to time between them) Newtonian mechanics applies Particles moves with uniform velocity between collisions Elastic collisions Particles move with (constant) random motion	(1)	Accept molecules for particles	1	
	Use of $\rho = m/V$ to determine mass of air in the balloon Calculation of total mass = mass of air at 120 °C + passengers + balloon Use of $W = mg$ $W = 31\ 600\ \text{N}$, which is less than 33 000 N, so the balloon can take off The one from: Point particles Particles have negligible volume (compared to their container) Negligible forces between particles (except during a collision) Range of molecular forces small (in comparison to average separation) Duration of collisions negligible (in comparison to time between them) Newtonian mechanics applies Particles moves with uniform velocity between collisions Elastic collisions	Use of $\rho = m/V$ to determine mass of air in the balloon Calculation of total mass = mass of air at 120 °C + passengers + balloon Use of $W = mg$ (1) $W = 31\ 600\ \text{N}$, which is less than 33 000 N, so the balloon can take off (1) The particles Particles have negligible volume (compared to their container) Negligible forces between particles (except during a collision) Range of molecular forces small (in comparison to average separation) Duration of collisions negligible (in comparison to time between them) Newtonian mechanics applies Particles moves with uniform velocity between collisions Elastic collisions	Use of $\rho = m/V$ to determine mass of air in the balloon Calculation of total mass = mass of air at 120 °C + passengers + balloon Use of $W = mg$ Use of $W = mg$ Use of $W = mg$ (1) $W = 31\ 600\ \text{N}$, which is less than 33 000 N, so the balloon can take off (1) The point particles Particles have negligible volume (compared to their container) Negligible forces between particles (except during a collision) Range of molecular forces small (in comparison to average separation) Duration of collisions negligible (in comparison to time between them) Newtonian mechanics applies Particles moves with uniform velocity between collisions Elastic collisions	

17(b)(ii)	•	States $pV = NkT$ or $pV = 1/3 Nm < c^2 >$	(1)		
	•	Suitable algebra to show $\frac{1}{2} m < c^2 > = 3/2 kT$	(1)		2
17(b)(iii)	•	Use of a.m.u. to determine particle mass	(1)	Example of calculation	
	•	Use of $\frac{1}{2} m < c^2 > = 3/2 kT$ with T in K	(1)	Mass = $28 \times 1.66 \times 10^{-27} \text{ kg} = 4.65 \times 10^{-26} \text{ kg}$ $\frac{1}{2} \times 4.65 \times 10^{-26} \text{ kg} < c^2 > = \frac{3}{2} \times 1.38 \times 10^{-23}$ J K ⁻¹ × 393 K c = $\sqrt{349900} = 590$ m s ⁻¹	
	•	rms speed = 590 m s^{-1}	(1)		3

(Total for Question 17 = 12 marks)

Question Number	Acceptable answers				Additional guidance	Mark
*18(a)	logically struct reasoning. Marks are awar answer is struct	essesses a student's ured answer with liner ded for indicative of tured and shows line table shows how the ontent.	econtent and for how es of reasoning.	ustained v the	The following table shows how the marks should be awarded Number of marks awarded for structure of answer and sustained line of reasoning Answer shows a coherent and logical structure with linkages and fully 2	
	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	Max linkage mark available	Max final mark	sustained lines of reasoning demonstrated throughout Answer is partially structured with some linkages and lines of reasoning Answer has no linkages between 0	
	6	4	2	6	points and is unstructured	
	5	3	2	5	for structure and lines of reasoning. Guidance on how the mark scheme should be applied: The mark for	
	4	3	1	4		
	3	2	1	3	indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking	
	2	2	0	2	points which is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1	
	1	1	0	1	mark for partial structure and some linkages and lines of reasoning). If there are no linkages between points, the same five indicative	
	0	0	0	0	marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).	

	Indicative content:			
	IC1 Electrons are in (discrete) energy levels			
	IC2 Absorption of (a single) photon causes an electron to move to a higher energy level			
	IC3 Photon energy = hf Or photon energy is proportional to frequency			
	IC4 Energy of (absorbed) photon must equal difference in energy levels			
	IC5 The (changes in) energy levels are specific to each element			
	IC6 Different wavelengths/frequencies of light are absorbed			6
18(b)	• Use of $c = f\lambda$	(1)	Example of calculation $f = 3.00 \times 10^8 \text{ m s}^{-1} / 656.46 \times 10^{-9} \text{ m}$	
	• Use of $\Delta E = hf$	(1)	$= 4.57 \times 10^{14} \text{ Hz}$ $= 6.63 \times 10^{-34} \text{ J s} \times 4.57 \times 10^{14} \text{ Hz}$	
	• Use of 1 eV = $1.6 \times 10^{-19} \text{ J}$	(1)	$=3.03 \times 10^{-19} \mathrm{J}$	
	• Change in energy equal to 1.89 eV and transition from energy level 2 to energy level 3	(1)	= $3.03 \times 10^{-19} \text{ J} / 1.60 \times 10^{-19} = 1.89 \text{ eV}$ 1.89 eV - 3.40 eV = -1.51 eV = level 3	4

18(c)(i)	(According to the Doppler effect) if the source is moving away from the observer the wavelength will increase	(1)	Allow converse statement i.e. If the source is moving towards the observer the wavelength will decrease	
	(According to the Doppler effect) there is a change in wavelength proportional to velocity	(1) (1)		
	He is correct (as speed and direction can be determined). MP3 dependent on MP1 and MP2			3
18(c)(ii)	• Use of $\Delta \lambda / \lambda_0 = v/c$	(1)	Must see_656.46 ×10 ⁻⁹ m in denominator	
	• $\lambda = 656 \times 10^{-9} \mathrm{m}$	(1)	Example of calculation $\Delta \lambda = 656.46 \times 10^{-9} \text{ m} \times 5.5 \times 10^{3} \text{ m s}^{-1}/3.00 \times 10^{8} \text{ m s}^{-1}$ = 0.012 ×10 ⁻⁹ m Moving away, so add $\Delta \lambda$ $\lambda = 656.46 \times 10^{-9} \text{ m} + 0.012 \times 10^{-9} \text{ m}$ $\lambda = 656.47 \times 10^{-9} \text{ m}$	2

(Total for Question 18 = 15 marks)

Question Number	Acceptable answers		Additional guidance	Mark
19(a)(i)	(Spontaneous) – independent of external conditions	(1)		
	(Random) – cannot predict when a decay will occur	(1)	Accept cannot tell which nucleus will decay next	2
19(a)(ii)	The probability of decay (in a specified time) may be determined	(1)	For MP1 accept reference to the proportion of nuclei decaying at a specified time	
	(Probability may be applied accurately because) very large numbers are involved	(1)		2
19(b)	A description that makes reference to the following points:			
	Measure the background count (rate)	(1)		
	Measure count (rate) from source	(1)		
	Use paper to absorb alpha radiation	(1)		
	• Percentage = $\frac{\text{count with paper}(-\text{background rate})}{\text{count from source }(-\text{background rate})} \times 100$	(1)		4
19(c)(i)	• Use of $\ln 2 = \lambda t_{1/2}$	(1)	Example of calculation $\lambda = \ln 2 / 432 \text{ years} = 1.6 \times 10^{-3} \text{ y}^{-1}$	
	• Use of $A = A_0 e^{-\lambda t}$	(1)	$A / A_0 = e^{-(1.6 \times 10^{-3} \text{ y} - 1 \times 34 \text{ y})}$	
	• Expected percentage of $A_0 = 95 \%$	(1)	$(A/A_0) \times 100 \% = 94.7 \%$	3

19(c)(ii)	Beta is only emitted after the decay of neptunium to protactinium Or Americium and neptunium only emit alpha	(1) (1)		
	• The half-life of <u>neptunium</u> is very long	(1)	Accept amount for proportion	
	 There will only be a very small proportion of protactinium (relative to americium and neptunium), so the suggestion is incorrect Or Only 5% of the americium has decayed so the proportion of protactinium is very small, so the suggestion is incorrect 			2

(Total for Question 19 = 14 marks) TOAL FOR PAPER = 90 MARKS