

#### © International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

#### © Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

#### © Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.





### Chemistry **Higher level** Paper 2

12 May 2023

Zone A afternoon	Zone B morning	Zone C	afternoon	
,				Candidate session number

2 hours 15 minutes

#### Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all questions.
- Answers must be written within the answer boxes provided.
- A calculator is required for this paper.
- A clean copy of the **chemistry data booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



Answer all questions. Answers must be written within the answer boxes provided. 1. This question is about acid-base properties. (a) Deduce the ionic equation, including state symbols, for the reaction of hydrogen chloride gas with water. [2] (ii) Calculate the pH of 0.50 mol dm<sup>-3</sup> hydrochloric acid. [1] (iii) Explain why a solution of ethanoic acid has a higher pH than hydrochloric acid of the same concentration. [1] (iv) A pH probe can be used to distinguish between the acids in part (a)(iii). Identify another simple instrumental method that could be used in a school laboratory to distinguish between the two acids. [1] Outline how the instrumental method identified in part (a)(iv) distinguishes (v) between the acids in part (a)(iii). [1]



(b)	Outline one chemical test, other than an indicator, that can distinguish between the two
	acids in part (a)(iii), and the expected result.

[1]


Chemical test:		 	 	
Expected result	t:	 	 	



Turn over

(ii)

(c) A neutralization curve for a weak acid, HA, and a strong base is given.



(i)	Estimate the $pK_a$ of HA.	[1]

Explain, using an equation, why adding a strong base to the weak acid, HA,

leads to very little change in pH in the buffer zone of the graph.	



In a separate experiment,  $80\,\mathrm{cm^3}$  of  $0.1\,\mathrm{mol}$  dm<sup>-3</sup> ammonia,  $\mathrm{NH_3}(\mathrm{aq})$ , was added to  $40\,\mathrm{cm^3}$  of  $0.1\,\mathrm{mol}$  dm<sup>-3</sup> hydrochloric acid, HCl (aq).

 (111)	Determine the final pH of the solution. Use section 21 of the data booklet.	[4]



**-6-** 2223-6108


Please do not write on this page.

Answers written on this page will not be marked.




- 2. The periodic table is a useful tool in explaining trends of chemical behaviour.
  - (a) (i) Annotate and label the ground state orbital diagram of boron, using arrows to represent electrons.

[1]



(ii) Sketch the shapes of the occupied orbitals identified in part (a)(i).

[2]



(b)	Expl	ain the decrease in first ionization energy from Li to Cs, group 1.	[2]
(c)	(i)	State the electron domain geometry of the ammonia molecule.	[1]
	(ii)	Deduce the Lewis (electron dot) structure of ammonia and sketch its 3D molecular shape.	[2]
Lev	vis stru	icture:	
3D	molec	ular shape:	



	(iii)	Explain, with reference to the forces between molecules, why ammonia has a higher boiling point than phosphine (PH <sub>3</sub> ).	[3]



Turn over

(d)	(i)	Ammonia	is manufactured	by the	Haber pr	rocess
-----	-----	---------	-----------------	--------	----------	--------

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g) \quad \Delta H_r^{\ominus} = -92.0 \, kJ \, mol^{-1}$$

Outline what is meant by dynamic equilibrium.

[1]


(ii) Deduce the $K_c$ expression for the reaction in part (d)(i).	[1]

(iii) Determine the entropy change,  $\Delta S^{\ominus}$  for the forward reaction to **four** significant figures, using the data given.

[2]

Substance	Entropy (S <sup>⊖</sup> ) J K <sup>-1</sup> mol <sup>-1</sup>
H <sub>2</sub>	130.7
N <sub>2</sub>	191.6
NH <sub>3</sub>	192.8




(iv)	Calculate the temperature, in K, below which this reaction becomes spontaneous. Use section 1 of the data booklet. (If you were unable to obtain an answer for part (d)(iii) use -210.0 J K <sup>-1</sup> mol <sup>-1</sup> , but this is not the correct value.)	[2]
(v)	The value of $K_c$ for this reaction is $6.84 \times 10^{-5}$ at $500^{\circ}$ C. Suggest, with a reason, how lowering the temperature affects the value of $K_c$ .	[1]
(vi)	Calculate the standard Gibbs free energy change, $\Delta G^{\ominus}$ , in kJ mol <sup>-1</sup> , for this reaction. Use sections 1 and 2 of the data booklet.	[2]

(This question continues on page 13)



Turn over

**- 12 -** 2223-6108

Please do not write on this page.

Answers written on this page will not be marked.



28FP12

(e) (i)	) T	he Haber process requires a catalyst. State how a catalyst functions.	[′
(ii		ketch a Maxwell–Boltzmann distribution curve showing the activation energinal vith and without a catalyst.	es [
		Fraction of particles	
		Kinetic energy (KE)	
	ii) S	uggest how the progress of the reaction could be monitored.	[




3. Alkanes form a homologous series.

1	(a)	) (i	(	Outline	the	meaning	٥f	homo	logous	series
١	(a	<i>)</i> (i	)	Outilite	uic	meaning	ΟI	HOHIO	logous	301103.

[1]


(ii) State the preferred IUPAC name for the following compounds.

[2]



(iii) Identify **one** chiral carbon atom present in one of the following structures with an asterisk (\*).

[1]



(iv)	But-2-ene can be polymerized. Draw a section of the resulting polymer showing <b>two</b> repeating units.	[1]

(b) Chloroethane can be converted into ethanoic acid in a two-step process.

Chloroethane	Step 1	Ethanol	Step 2	Ethanoic acid

Identify reagents for each step.

[2]

Step 1:	 
·	



**Turn over** 

Ques	tion 3	3 со	ntinued)	
(	(c)	(i)	Identify the type of reaction that takes place in step 1 of part (b).	[1]
	(	(ii)	Sketch the mechanism of the reaction for step 1 in part (b), using curly arrows to show the movement of electron pairs.	[4]
	(	(iii)	Identify the products formed from the reaction of ethanol and ethanoic acid in the presence of an acid catalyst.	[1]



28FP16

4.	Redox reactions can be used to produce electricity.							
	(a)	State	e the oxidation state of sulfur in copper(II) sulfate.	[1				
	(b)		Itaic cell was constructed using a copper $(\mathrm{II})$ sulfate/copper half-cell and a sulfate/zinc half-cell.					
		(i)	Outline why electrons flow from zinc to copper when these half cells are connected with a wire. Use section 25 of the data booklet.	[′				
		(ii)	Formulate equations for the reactions taking place at each electrode.	[2				
	Ano	de (ne	egative electrode):					
	Cat	hode (	positive electrode):					
	(c)	(i)	Calculate the standard cell potential for the voltaic cell in part (b). Use section 24 of the data booklet.	[1				
		(ii)	Calculate the standard Gibbs free energy change, $\Delta G^{\ominus}$ , in kJ mol <sup>-1</sup> , for this reaction. Use section 1 of the data booklet. (If you did not answer part (c)(i) use 1.05 V, but this is not the correct value.)	[2				



**5.** Double salts are substances with two cations and one anion. A hydrated sulfate containing two cations has this percentage composition.

Element	Percentage (%)
Nitrogen (N)	7.09
Hydrogen (H)	5.11
Sulfur (S)	16.22
Cobalt (Co)	14.91
Oxygen (O)	_

(a)	(i)	Draw <b>one</b> Lewis (electron dot) structure of the sulfate ion.	[1]
	(ii)	Calculate the percentage of oxygen present in the double salt.	[1]



(iii)	Determine the empirical formula of the double salt. Use section 6 of the data booklet.	[3]
(iv)	The molar mass of the empirical formula is the same as the molar mass of the formula unit. Deduce the formula unit of the hydrated double salt.	[1]
		(iv) The molar mass of the empirical formula is the same as the molar mass of the



Turn over

(b)		g of the double salt was dissolved in water and an excess of aqueous barium ide was added, precipitating all the sulfate ions as barium sulfate.	
	(i)	Formulate an ionic equation, including state symbols, for the reaction of barium ions with sulfate ions.	[1]
	(ii)	Calculate the mass of barium sulfate precipitate. Use your answer to part (a)(iii) and section 6 of the data booklet. (If you did not obtain an answer for part (a)(iii), use $400.0 \mathrm{g} \mathrm{mol}^{-1}$ as $M_{_{\mathrm{f}}}$ for the double salt, but this is not the correct value.)	[2]



[1]

- The element sulfur has many industrial uses. 6.
  - Determine the standard enthalpy of reaction ( $\Delta H_{\rm r}^{\ominus}$ ), in kJ mol<sup>-1</sup>, for the oxidation (a) of  $SO_2$  to  $SO_3$ .


Substance	Enthalpy of formation, $\Delta H_{\rm f}^{\ominus}$ (kJ mol <sup>-1</sup> )
SO <sub>2</sub>	-296.8
SO <sub>3</sub>	-395.8

(ii) Formulate equations showing how SO <sub>2</sub> and SO <sub>3</sub> lead to acid deposition.	[1]
SO <sub>2</sub> :	
(iii) Explain the polarity of the S—O bond. Use section 8 of the data booklet.	[2]



**Turn over** 

(b) The combustion of 0.1 moles of sulfur (S) was demonstrated in a school laboratory using the following apparatus in a fume cupboard.



(i) Calculate the enthalpy of combustion of sulfur,  $\Delta H_{\rm c}$ , in kJ mol<sup>-1</sup> from this data. Use sections 1 and 2 of the data booklet.

[2]

Mass of water (g) ±0.01	50.00
Initial temperature of water (°C) ±0.5	20.0
Final temperature of water (°C) ±0.5	35.0

								 								 	 								 	 	 				٠								 																				
							_			_	_		_			 	 								 		 											_								 										_		_	 _
•	•	•	•	•	•	•					•	•	•	•	•		•	•	•	•	•	•	•	•				•	•	•	•	•	•	•	•	•	•			•	•	•	•	•	•	 •	•	•	•	•	•	•	•	•	•		•	•	
	٠	•	•	•	•											 	 			•	•	•	•	٠		 		•	•	•	٠	٠	٠	٠	•				 	-	٠	•	•	•		 -	٠	•	٠	•	٠	•							



(ii)	Suggest the major source of systematic error in this experiment and an improvement to reduce this error.	[2]
Source of	systematic error:systematic error:	
Improvem	nent:	
(iii)	Calculate the percentage uncertainty in the temperature change to <b>two</b> significant figures.	[1]
(iv)	Suggest <b>one</b> way of reducing the percentage uncertainty in this experiment.	[1]
(v)	Calculate the overall percentage error of this experiment. Use part (b)(i) and section 13 of the data booklet. (If you did not obtain an answer for part (b)(i) use $-50.0  \text{kJ mol}^{-1}$ , but this is not the correct value.)	[1]



7. The structural formulae of two esters of formula  $C_3H_6O_2$  are shown.

#### **Ethyl methanoate**

## Methyl ethanoate

(a) (i) Deduce the number of signals you would expect to find in the <sup>1</sup>H NMR spectrum of each compound.


[1]

Name	Number of signals
Ethyl methanoate	
Methyl ethanoate	

(ii)	Outline why infrared spectroscopy is not used to differentiate between the two esters.	[1]



8. A series of experiments were carried out at different temperatures and the rate of reaction, in mol dm $^{-3}$  s $^{-1}$ , was determined for each. The rate constant for the reaction of propanone (CH $_3$ COCH $_3$ ) with iodine ( $I_2$ ) was calculated and the processed data is represented in the following graph.



Determine the activation energy for this reaction, stating the units. Use sections 1 and 2 of the data booklet.


[3]

(a)	rea	ctions of Zn <sup>2+</sup> compounds. Use section 3 of the data booklet.	
(b)	Nitı	rogen (II) oxide radicals (NO $\bullet$ ) catalyse the decomposition of ozone (O $_3$ ).	
	(i)	Formulate equations showing how NO• acts as a catalyst in this reaction.	
	Ch	orine also forms free radicals; the bond enthalpy for $\text{Cl}_2$ is $4.02 \times 10^{-19}  \text{J}$ .	
	(ii)	Calculate the minimum frequency of light needed to break this bond. Use sections 1 and 2 of the data booklet.	



(iii) Calculate the formal charge on each atom in the two Lewis structures of the  $NO_2^{\bullet}(g)$  radical.

[1]

	Structure A	Structure B
	: 0 N 1 0:	: 0 0 :
Oxygen 1		
Nitrogen		
Oxygen 2		

(iv)	) Lewis structure A is more stable. Suggest, giving <b>one</b> reason, whether the formal charge model supports this.	[1]



#### References:

© International Baccalaureate Organization 2023

