

Proteins and enzymes 3

Level: Edexcel A Level 9BN0 Subject: Biology Exam Board: Suitable for all boards Topic: Proteins and enzymes 3 Type: Mark Scheme

To be used by all students preparing for Edexcel Biology A Level 9BN0 foundation or higher tier but also suitable for students of other boards.



Mark schemes

	(a)	Lilac / purple / mauve / violet;		
1		Xanthine oxidase is a protein; Reject pink or blue as the resulting colour with biuret.	2	
	(b)	Substrate has specific shape;		
		Allows binding / fitting / forms ES complex with active site;		
		Or		
		Active site has specific shape;		
		Allows binding / fitting / forms ES complex with substrate; Accept structure ≡ shape	2	
	(c)	Xanthine <u>similar</u> shape to drug;	-	
		Drug fits active site / competes for active site / is a competitive inhibitor;		
		Less / no uric acid formed;		
			3	[7]
2	USE	of water; must be above arrowhead OH drawn correctly in place of glycosidic bond on each monosaccharide;		[2]
3	(a)	Stimulates memory cells;		
3		Secondary response, so antivenom / antibodies produced quicker;	2	
	(b)	Passive immunity; so no memory cells produced;		
		Antivenom breaks down / destroyed;	2	
	(c)	Could transfer disease / Allergy / Immune response to antibodies from animal;	_	
			1	[5]



4

(a)

	Glycolysis	Link reaction	Krebs cycle
Occurs in mitochondria		\checkmark	\checkmark
Carbon dioxide produced		\checkmark	\checkmark
NAD is reduced	\checkmark	\checkmark	\checkmark

Mark horizontally

3

2

(b) (i) 1. Glucose is used / broken down during glycolysis / in cytoplasm;
 1. Accept: glucose to pyruvate or glucose not converted to pyruvate for one mark

- Glucose cannot cross mitochondrial <u>membrane(s)</u> / pyruvate can cross mitochondrial <u>membrane(s);</u>
- (ii) 1. Is a competitive inhibitor / attaches to active site;
 1 Accept: inhibitor / malonate attaches to active site to form an enzyme-substrate complex
 - Reduces / prevents enzyme-substrate / E-S complex forming;
 2 Accept: substrate / succinate cannot bind to enzyme
 2 Accept mark point 2, but not mp1 in context of non-competitive inhibition
- (iii) 1. Krebs cycle inhibited as NAD / Coenzyme / FAD not / less reduced;
 - 2. Hydrogens not passed to ETC therefore oxygen not used as (much as a) final / terminal (electron) acceptor;

2



- (a) (i) Glucose <u>and</u> fructose;
 Ignore reference to alpha and beta
 Either way around
 - (ii) Glucose <u>and</u> galactose;
 Ignore reference to alpha and beta
 Either way around
- (b) 1. (Amylase) pancreas, produces maltose;
 Place <u>and</u> product = 1 mark (mark horizontally)
 - (Maltase) in / on epithelium (of small intestine), produces glucose; Ignore references to salivary glands or saliva Accept wall / lining of small intestine Ignore reference to cells alone Ignore reference to ribosomes / rER

[4]

2

1

1



6	(a)	1.	(Phosphate) changes shape of TK / changes shape of enzyme / changes the active site;		
			lt = phosphate		
			Accept 'alters' for changes		
			Reject that phosphate is an inhibitor		
			Accept adding energy / affecting charged / affects polar groups (on amino acids)		
		2.	Active site forms / becomes the right shape / can bind to substrate / complementary to substrate / E-S complex can form;		
			Reject similar / same shape as substrate		
				2	
	(b)	1.	Faulty TK has functional active site without phosphate;		
	. ,		Accept 'works without phosphate'		
		0			
		2.	(So, faulty) TK functional all the time / TK not controlled (by phosphate);	2	
				2	
	(c)	1.	Non-competitive inhibitor / binds to site other than active site;		
			Accept allosteric site		
			Do not accept 'changes shape' unqualified		
		2.	Causes TK to be in non-functional form / active site not formed / wrong shape / E-S complex not formed;		
		3.	So, (uncontrolled) cell division stopped / slowed / controlled;	2 max	
					[6]
	(a)	1.	Cell wall not formed / production inhibited;		
7	(0)		1. Q Accept: weakened cell wall, but do not accept 'cell wall is		
			broken down'		
		2.	Lower water potential in bacterium;		
		2.	2. Accept: converse		
			2. Must be clear that the lower water potential is in the bacterium		
			·		
		3.	Water enters and causes lysis / expansion / pressure;	2	
				2 max	
	(b)		nan cells lack enzyme (B) / have a different enzyme / produce different fatty acids e different substrates;		
			Neutral: 'human cells do not have cell walls' as out of context		
				1	

For more help visit our website https://www.exampaperspractice.co.uk/



	(c)	1.	Change in base sequence (of DNA / gene) leading to change in amino acid sequence / primary structure (of enzyme);		
			1. Accept: different amino acids coded for		
			1. Reject: different amino acids produced		
		2.	Change in hydrogen / ionic / disulphide bonds leading to change in the tertiary structure / active site (of enzyme);		
			2. Neutral: alters 3D structure / 3D shape		
		3.	Substrate not complementary / cannot bind (to enzyme / active site) / no enzyme-substrate complexes form;		
				3	[6]
					[6]
8	(a)	(i)	(Human cells) don't have a cell wall;		
			Accept "they" refers to human cells.	1	
				I	
		(ii)	(Affects) protein synthesis;		
			Allow description e.g. 'amino acids not joined together / translation.		
			Reject: affects transcription.	1	
	4.5			•	
	(b)	1.	Mutation present / occurs;		
			Ignore antibiotic causes mutation.		
		2.	Resistance gene / allele;		
			1. or 2.		
			Reference to immunity disqualifies first credited marking point.		
		3.	Resistant bacteria (survive and) reproduce;		
			Reference to mitosis negates marking point 3.		
				2	
					[4]
9	(a)	(i)	Hydrolysis;		
9			Accept phonetic spelling.		
			Ignore reaction.		
				1	
		(ii)	(Alpha) glucose;		
			Accept a glucose.		
			Reject β glucose / beta glucose		
				1	

For more help visit our website https://www.exampaperspractice.co.uk/



	(b)	(i)	Add Benedict's (reagent) and heat / warm;		
			Red / orange / yellow / green (colour);		
			Reject Add HCI		
			Accept brown, reject other colours	2	
		(ii)	2 products / 2 sugars produced;	-	
		()	Look for idea of two		
			Accept named monosaccharides produced.		
			"More" insufficient for mark		
			Neutral if incorrect products named		
			Neutral "lactose is a polysaccharide"		
			Neutral "lactose is not a reducing sugar"		
			Neutral: Reference to surface area.	1	
	(\mathbf{a})	1	Coloctopo io o cimilar chono (structure to loctopo (hoth complementary)	1	
	(c)	1.	Galactose is a similar shape / structure to lactose / both complementary; Q Reject: <u>Same</u> shape / structure		
			a nojeci. <u>Jame</u> shape / shakare		
		2.	(Inhibitor / Galactose) fits into / enters / binds with <u>active site</u> (of enzyme);		
			Accept blocks active site		
		3.	Prevents / less substrate fitting into / binding with (active site) / fewer or no E-S complexes;		
			Look for principles:		
			1. Shape		
			2. Binding to active site		
			3. Consequence	2 max	
				2 max	[7]
	(a)	2 ma	arks for Chymosin / enzyme is a protein; ;		
10	(u)				
		Prote	ein / peptide bonds present;	2 max	
	4.5			2 mux	
	(b)	wate	er removed;	1	
				_	[3]
11	Simi	lar str	ucture / shape (to PABA) / both complementary;		
	Com	petes	for / binds to active site / competitive inhibitor;		
		Less	PABA binds / less E-S complexes:		



OR

12

Specific reference to different structure / shape (to PABA) using the diagram;

Binds to position other than active site / binds to allosteric site / binds to inhibitor site / non-competitive inhibitor;

Changes the active site so substrate cannot bind / less PABA binds / less E-S complexes;

Q Reject: same structure / shape Note: competitive inhibitor binds to active site = 1 mark (same mark point) Assume that 'it' refers to sulfanilamide Accept: PABA / substrate cannot bind Neutral: less product produced as in question stem Neutral: different structure / shape to PABA Reject: active site on substrate for second marking point only

(a)				
	*	*	*	
				*
			×	×

One mark for each correct column Mark ticks only and ignore crosses

- (b) 1. Two marks for box round two hydrogens and one of the oxygens from OH groups on carbons 1 and 4;;
 - 2. One mark from incorrect answer involving any two hydrogens and an oxygen from carbons 1 and 4;

Do not award marks if all atoms concerned are on same carbon atom or are on carbon atoms other than 1 and 4 or where the answer does not have two hydrogen and one oxygen

- (c) (i) 1. Holds chains / cellulose molecules together / forms cross links between chains / cellulose molecules / forms microfibrils, providing strength / rigidity (to cellulose / cell wall);
 - Hydrogen bonds strong in large numbers;x
 Principles here are first mark for where hydrogen bonds are formed and second for a consequence of this.
 Accept microfibres

[3]

4



(ii) Compact / occupies small space / tightly packed;

Answer indicates depth required. Answers such as "good for storage", "easily stored" or "small" are insufficient.

1

13	(a)	Active site (complementary / specific) structure / shape; (Only) fits / binds to gangliosides;
		Forms enzyme-substrate complexes;

OR

	Active site (complementary / specific) structure / shape; (Does not) fit / bind with other lipids; Does not form enzyme-substrate complexes; <i>Note: 'active site has a specific shape' = 2 marks;</i>		
	<i>Reject: same shape Second mark for either route can refer to the enzyme or the substrate</i>		
	Accept: converse of second mark point and (different) structure / shape if referring to other lipids	3 max	
(b)	(i) No change / substrate remains high / horizontal line; <i>Curve should be labelled</i>	5 1114X	
	If curve H correctly labelled then assume other is curve T		
	Reject: obvious rise or fall / rise then plateau	1	
	 (ii) Curve decreases rapidly at first then more slowly; Curve should be labelled If curve <i>T</i> correctly labelled then assume other is curve <i>H</i> 		
	Reject: falling at a slower rate initially	1	
(c)	(Enzymes are) proteins; Digested / broken down / destroyed (by enzymes / acid);		
	OR		
	(Enzymes are) too large; To cross cell membranes / be absorbed / enter the bloodstream; Accept: denatured (by acid) Neutral: digested by saliva Reject: digested by amylase Neutral: will not reach the bloodstream		
		2	



14	(a)	Amr	nonia / ammonium / NH ₃ / NH ₄ +;	1	
	(b)		have similar shape / tertiary structure (as substrate) / complementary shape (to /e site);		
			Neutral: same shape as substrate		
		Fit /	bind with active site / forms enzyme-substrate complex;		
			Reject: same shape as active site	2	
	(c)	(i)	Provides ATP for the reaction / nitrogen fixation / reduction of nitrogen / formation of ammonia;		
			Accept: ATP or energy		
			Enzyme / nitrogenase produced quicker / <u>more</u> enzyme produced; Ignore references to temperature		
			Uses / removes oxygen (so nitrogenase works);		
			Use of oxygen must be in the correct context	2	
				2 max	
		(ii)	ATP used for / needed for nitrogen fixation / reduction of nitrogen / formation of ammonia / production of enzyme / nitrogenase;		
			Accept: ATP or energy		
			(So less ATP) available for growth / protein synthesis / production of new cells / production of biomass;		
			Accept: converse for those without fertiliser		
				2	[7]
	(a)	(i)	Faster / greater / more effective response in children;		
15	(u)	(1)	Do not accept children have more haemoglobin		
				1	
		(ii)	Use line of best fit;		
				1	
			Extrapolate / extend line (and read from graph);		
			Allow calculation using rate of increase per day = one mark.		
			However for both marks this must be linked to line of best fit.	1	
		(iii)	More than one polypeptide chain;		
		()	Allow many polypeptide chains.		
			'Haemoglobin has four polypeptide chains' must be in correct		
			context to gain mark.	1	
				1	



	(b)	(i)	Has same <u>water potential;</u> Allow converse for effect of using distilled water or a concentrated solution.	1	
			No (net) water movement / osmosis;	1	
			Cells will not swell / burst / change size; No osmotic lysis = two marks	1	
		(ii)	Pernicious anaemia (cells) greater range / spread / variation of diameters / widths;	1	
			Some pernicious anaemia (cells) wider than 9 (μ m) / some less than 5.5 (μ m) / without pernicious anaemia none more than 9 (μ m) / none less than 5.5 (μ m);		
			Pernicious anaemia (cells) peak / most frequent at 8.5 (μ m) / peak / most frequent at higher diameter / / without pernicious anaemia peak / most frequent at 7 (μ m) / peaks at lower diameter;		
			There are several alternatives for marking points 2 and 3	2 max	[9]
16	(a)	(i)	(Lactose +) <u>Water</u> ; \rightarrow (Glucose +) <u>Galactose</u> ; Accept: H ₂ O for water	2	
		(ii)	<u>Hydrolysis;</u> Accept: if phonetically correct	-	
	(b)	(i)	(Add Biuret reagent to both solutions) – no mark; Neutral: positive / negative result		
			Lactase / enzyme will give purple / lilac / mauve; Neutral: incorrect reference to the method		
			OR		
			Lactose / reducing sugar will not give purple / lilac / mauve / will remain blue;	1	
		(ii)	Lactase / enzyme is a protein;		
			Accept: lactase / enzyme contains peptide bonds	1	[5]



		EXAM PAPERS PRACTICE		
17	(a)	Enzyme / active site has a (specific) tertiary structure;		
		Only glucose has correct shape / is complementary / will bind / fit to active site;		
		 (Forming) enzyme-substrate <u>complex</u>; <i>Q</i> Allow second mark if candidate refers to correct shape or complementary in terms of the enzyme. Do not allow 'same' shape <i>Q</i> Do not allow third mark if active site is described as being on substrate. 	3	
	(b)	(Only detects glucose whereas) Benedict's detects (all) reducing sugars / named examples;		
		Provides a reading / is quantitative / Benedict's only provides a colour / doesn't measure concentration / is qualitative / semiquantitative;		
		Is more sensitive / detects low concentration;		
		Red colour / colour of blood masks result;		
		Can monitor blood glucose concentration continuously; Q Do not credit quicker / more accurate unless qualified.		
		Q Allow Benedict's detects monosaccharides for first mark point.	2 max	
	(c)	 Broken down by enzymes / digested / denatured (by pH) too large to be absorbed; 	1	
		 Study not carried out on humans / only carried out on rats; Long-term / side effects not known; Scientists have vested interest; Study should be repeated / further studies / sample size not known; 	2 max	
				[8]
18	(a)	(i) <u>Deoxyribose;</u> pentose / 5C sugar = neutral	1	
		 (ii) Phosphate / Phosphoric acid; phosphorus / P = neutral 		
	(b)	Hydrogen (bonds);	1	
	(c)	381 / 384 / 387;	1	
	(d)	(GIn) Met Met Arg Arg Arg Asn;	1	



(e) Change in (sequence of) amino acids / primary structure;

Change in hydrogen / ionic / disulfide bonds leads to change in tertiary structure / active site (of enzyme);

Substrate cannot bind / no enzyme-substrate complexes form;

Q Reject = different amino acids are formed

19 (a) (i) Glucose;

Fructose; Any order.

(ii) Lactose has a different shape / structure;

Does not fit / bind to active site of enzyme / sucrase;

Only allow a second mark if reference is made to the active site. Max 1 mark if active site is described as being on the substrate.

OR

Active site of enzyme / sucrase has a specific shape / structure; Does not fit / bind to lactose;

Do not accept same shape.

(b) (i) Rose and fell;

Peak at 45 (minutes) / concentration of 6.6 (mmol dm⁻³);

(ii) Glucose (produced by digestion) is absorbed / enters blood;

Decrease as used up / stored;

50

3

2

2

2

2

[8]



20	(a)	(i)	150;	1
		(ii)	27;	1
	(b)		; ber of peptide bond hydrolysed = total number present / all peptide bonds have n hydrolysed; accept calculation showing same number top and bottom.	2
	(c)	curv	re rising to peak at pH 2 and falling to zero by pH 6;	
				1
	(d)		ange in pH) leads to breaking of bonds holding tertiary structure / changes charge amino acids;	
		enzy	yme / protein / active site loses shape / denatured; substrate will not bind with / fit ve site / fewer / no ES complexes formed;	
				3
	(e)		e resistant to changes in pH and washing conditions variable / works in alkaline and washing powders alkaline;	
			mark awarded for indicating aspect of effect of pH and advantage of this in terms of washing powder and conditions in wash.	1
	(f)		ximum of three marks for specificity, points 1 - 3. Can only be given credit in text of specificity	
		1	each enzyme / protein has specific primary structure / amino acid sequence;	
		2	folds in a particular way / has particular tertiary structure giving an active site with a unique structure;	
		3	shape of active site complementary to / will only fit that of substrate; maximum of three marks for inhibition, points $5 - 8$	
		4	inhibitor fits at site on the enzyme other than active site;	
		5	distorts active site;	
		6	so substrate will no longer fit / form enzyme-substrate complex	6



- (a) Accept three suitable suggestions:
 - 1. (Lactase / beads) can be reused / not washed away;
 - 1. Accept lactase / beads not wasted
 - 1. Less lactase used is insufficient
 - 2. No need to remove from milk;
 - 2. Accept lactase not present in milk.
 - 3. Allows continuous process;
 - 4. The enzyme is more stable;
 - 5. Avoid end-product inhibition.

Ignore ref to SA

3 max

- (b) 1. (Lactose hydrolysed to) galactose and glucose;
 - 2. (So) more sugar molecules;
 - 2. Idea of more sugars essential
 - 3. (So) more / different receptors stimulated / sugars produced are sweeter (than lactose).

2 max

[5]

(Maintaining constant pH to avoid)

22

 Named protein / enzyme (in blood) sensitive to / affected by change in pH;

Accept converse for MP2 and MP3.

Named example should be a protein that might be affected (by change in pH) eg haemoglobin, carrier protein in plasma membrane.

Accept 'change in H + concentration' for 'change in pH'.

2. (Resultant) change of charge / shape / tertiary structure;

The change in charge idea relates to the enzyme / protein and not the blood (plasma) or red blood cells. 'Denaturation' alone is insufficient.

Described effect on named protein or enzyme.
 e.g. less oxygen binds with haemoglobin / less transport across membranes / fewer substrates can fit active site / fewer enzyme-substrate complexes.

Idea of 'less' or 'fewer' required. Ignore suggestion of 'no' or 'none'.



- (a) 1. (Enzyme has) active site;
 - 1. Reject active site is same shape as substrate
 - 1. Reject active site is on the substrate
 - 1. Accept active site forms during induced fit
 - 2. Only substrate fits (the active site);
 - 2. Accept converse statement
 - (b) Assume "it" = allopurinol
 - 1. (Allopurinol) is a similar shape to xanthine;
 - 1. Reject same shape. Accept similar structure
 - 2. (Allopurinol) enters active site / is a competitive inhibitor;
 - 2. Ignore e-s complexes in relation to inhibitor
 - 2. Reject non-competitive inhibitor in the context of binding to the active site
 - 2. Ignore complementary / fits
 - Less xanthine binds / fewer e-s complexes / fewer uric acid crystals formed / less uric acid formed;

3. Reject <u>no</u> e-s complexes / xanthine <u>cannot</u> enter active site, <u>no</u> uric acid

3. Can award in context of non-competitive inhibition

3

2