

Further Pure 2

1	Inequalities	3
	Algebraic solutions	3
	Graphical solutions	4
2	Series – Method of Differences	5
3	Complex Numbers	7
	Modulus and Argument	7
	Properties.....	7
	Euler's Relation $e^{i\theta}$	7
	Multiplying and dividing in mod-arg form	7
	De Moivre's Theorem.....	8
	Applications of De Moivre's Theorem	8
	$z^n + \frac{1}{z^n} = 2 \cos n\theta$ and $z^n - \frac{1}{z^n} = 2i \sin n\theta$	8
	n^{th} roots of a complex number	9
	Roots of polynomial equations with real coefficients	10
	Loci on an Argand Diagram.....	10
	Transformations of the Complex Plane	12
	Loci and geometry	13
4	First Order Differential Equations	14
	Separating the variables, families of curves.....	14
	Exact Equations	14
	Integrating Factors.....	15
	Using substitutions	15
5	Second Order Differential Equations	17
	Linear with constant coefficients	17
	(1) when $f(x) = 0$	17
	(2) when $f(x) \neq 0$, Particular Integrals	18
	D.E.s of the form $a \frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = f(x)$	21
6	Maclaurin and Taylor Series	22
	1) Maclaurin series.....	22
	2) Taylor series	22
	3) Taylor series – as a power series in $(x - a)$	22

4) Solving differential equations using Taylor series	22
Standard series	23
Series expansions of compound functions	25
7 Polar Coordinates	26
Polar and Cartesian coordinates.....	26
Sketching curves	26
Some common curves	26
Areas using polar coordinates.....	29
Tangents parallel and perpendicular to the initial line	30
Appendix	32
n^{th} roots of 1	32
Short method.....	32
Sum of n^{th} roots of 1	33
1 st order differential equations	34
Justification of the Integrating Factor method.....	34
Linear 2 nd order differential equations	35
Justification of the A.E. – C.F. technique for unequal roots	35
Justification of the A.E. – C.F. technique for equal roots	36
Justification of the A.E. – C.F. technique for complex roots	37
Justification that G.S. = C.F. + P.I.	38
Maclaurin's Series.....	39
Proof of Maclaurin's series.....	39
Proof of Taylor's series	39

1 Inequalities

Algebraic solutions

Remember that if you multiply both sides of an inequality by a negative number, you must turn the inequality sign round: $2x > 3 \Rightarrow -2x < -3$.

A difficulty occurs when multiplying both sides by, for example, $(x - 2)$; this expression is sometimes positive ($x > 2$), sometimes negative ($x < 2$) and sometimes zero ($x = 2$). In this case we multiply both sides by $(x - 2)^2$, which is always positive (provided that $x \neq 2$).

Example 1: Solve the inequality $2x + 3 < \frac{x^2}{x-2}$, $x \neq 2$

Solution: Multiply both sides by $(x - 2)^2$

we can do this since $(x - 2) \neq 0$

$$\Rightarrow (2x + 3)(x - 2)^2 < x^2(x - 2)$$


DO NOT MULTIPLY OUT

$$\Rightarrow (2x + 3)(x - 2)^2 - x^2(x - 2) < 0$$

$$\Rightarrow (x - 2)(2x^2 - x - 6 - x^2) < 0$$

$$\Rightarrow (x - 2)(x - 3)(x + 2) < 0$$

$$\Rightarrow x < -2, \text{ or } 2 < x < 3, \text{ below } x\text{-axis}$$



Note – care is needed when the inequality is \leq or \geq .

Example 2: Solve the inequality $\frac{x}{x+1} \geq \frac{2}{x+3}$, $x \neq -1$, $x \neq -3$

Solution: Multiply both sides by $(x + 1)^2(x + 3)^2$

which cannot be zero

$$\Rightarrow x(x + 1)(x + 3)^2 \geq 2(x + 3)(x + 1)^2$$

DO NOT MULTIPLY OUT

$$\Rightarrow x(x + 1)(x + 3)^2 - 2(x + 3)(x + 1)^2 \geq 0$$

$$\Rightarrow (x + 1)(x + 3)(x^2 + 3x - 2x - 2) \geq 0$$


$$\Rightarrow (x + 1)(x + 3)(x + 2)(x - 1) \geq 0$$

from sketch it looks as though the solution is

$$x \leq -3 \text{ or } -2 \leq x \leq -1 \text{ or } x \geq 1$$

BUT since $x \neq -1$, $x \neq -3$,

the solution is $x < -3$ or $-2 \leq x < -1$ or $x \geq 1$, above the x -axis



Graphical solutions

Example 1: On the same diagram sketch the graphs of $y = \frac{2x}{x+3}$ and $y = x - 2$.

Use your sketch to solve the inequality $\frac{2x}{x+3} \geq x - 2$

Solution: First find the points of intersection of the two graphs

$$\Rightarrow \frac{2x}{x+3} = x - 2$$


$$\Rightarrow 2x = x^2 + x - 6$$

$$\Rightarrow 0 = (x - 3)(x + 2)$$

$$\Rightarrow x = -2 \text{ or } 3$$

From the sketch we see that

$$x < -3 \text{ or } -2 \leq x \leq 3. \quad \text{Note that } x \neq -3$$



For inequalities involving $|2x - 5|$ etc., it is often essential to sketch the graphs first.

Example 2: Solve the inequality $|x^2 - 19| < 5(x - 1)$.

Solution: It is essential to sketch the curves first in order to see which solutions are needed.

To find the point A, we need to solve

$$-(x^2 - 19) = 5x - 5 \Rightarrow x^2 + 5x - 24 = 0$$

$$\Rightarrow (x + 8)(x - 3) = 0 \Rightarrow x = -8 \text{ or } 3$$

$$\text{From the sketch } x \neq -8 \Rightarrow x = 3$$


To find the point B, we need to solve

$$+(x^2 - 19) = 5x - 5 \Rightarrow x^2 - 5x - 14 = 0$$

$$\Rightarrow (x - 7)(x + 2) = 0 \Rightarrow x = -2 \text{ or } 7$$

$$\text{From the sketch } x \neq -2 \Rightarrow x = 7$$

$$\Rightarrow \text{the solution of } |x^2 - 19| < 5(x - 1) \text{ is } 3 < x < 7$$



2 Series – Method of Differences

The trick here is to write each line out in full and see what cancels when you add.

Do not be tempted to work each term out – you will lose the pattern which lets you cancel when adding.

Example 1: Write $\frac{1}{r(r+1)}$ in partial fractions, and then use the method of differences to find

the sum $\sum_{r=1}^n \frac{1}{r(r+1)} = \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{n(n+1)}.$

Solution: $\frac{1}{r(r+1)} = \frac{1}{r} - \frac{1}{r+1}$

put $r = 1 \Rightarrow \frac{1}{1 \times 2} = \frac{1}{1} - \frac{1}{2}$

put $r = 2 \Rightarrow \frac{1}{2 \times 3} = \frac{1}{2} - \frac{1}{3}$

put $r = 3 \Rightarrow \frac{1}{3 \times 4} = \frac{1}{3} - \frac{1}{4}$

etc.

put $r = n \Rightarrow \frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$

adding $\Rightarrow \sum_{r=1}^n \frac{1}{r(r+1)} = 1 - \frac{1}{n+1} = \frac{n}{n+1}$

Example 2: Write $\frac{2}{r(r+1)(r+2)}$ in partial fractions, and then use the method of differences to find the sum $\sum_{r=1}^n \frac{1}{r(r+1)(r+2)} = \frac{1}{1 \times 2 \times 3} + \frac{1}{2 \times 3 \times 4} + \frac{1}{3 \times 4 \times 5} + \dots + \frac{1}{n(n+1)(n+2)}$.

Solution:
$$\frac{2}{r(r+1)(r+2)} = \frac{1}{r} - \frac{2}{r+1} + \frac{1}{r+2}$$

put $r = 1 \Rightarrow \frac{2}{1 \times 2 \times 3} = \frac{1}{1} - \frac{2}{2} + \frac{1}{3}$

put $r = 2 \Rightarrow \frac{2}{2 \times 3 \times 4} = \frac{1}{2} - \frac{2}{3} + \frac{1}{4}$

put $r = 3 \Rightarrow \frac{2}{3 \times 4 \times 5} = \frac{1}{3} - \frac{2}{4} + \frac{1}{5}$

put $r = 4 \Rightarrow \frac{2}{4 \times 5 \times 6} = \frac{1}{4} - \frac{2}{5} + \frac{1}{6}$

...

etc.

...

put $r = n-1 \Rightarrow \frac{2}{(n-1)n(n+1)} = \frac{1}{n-1} - \frac{2}{n} + \frac{1}{n+1}$

put $r = n \Rightarrow \frac{2}{n(n+1)(n+2)} = \frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2}$

adding $\Rightarrow \sum_{r=1}^n \frac{2}{r(r+1)(r+2)} = \frac{1}{1} - \frac{2}{2} + \frac{1}{2} + \frac{1}{n+1} - \frac{2}{n+1} + \frac{1}{n+2}$

$$= \frac{1}{2} - \frac{1}{n+1} + \frac{1}{n+2}$$

$$= \frac{n^2 + 3n + 2 - 2n - 4 + 2n + 2}{2(n+1)(n+2)}$$

$$\Rightarrow \sum_{r=1}^n \frac{2}{r(r+1)(r+2)} = \frac{n^2 + 3n}{2(n+1)(n+2)}$$

$$\Rightarrow \sum_{r=1}^n \frac{1}{r(r+1)(r+2)} = \frac{n^2 + 3n}{4(n+1)(n+2)}$$

3 Complex Numbers


Modulus and Argument

The modulus of $z = x + iy$ is the length of z

$$\Rightarrow r = |z| = \sqrt{x^2 + y^2}$$

and the argument of z is the angle made by z with the positive x -axis, $-\pi < \arg z \leq \pi$.

N.B. $\arg z$ is **not always** equal to $\tan^{-1}\left(\frac{y}{x}\right)$



Properties

$$z = r \cos \theta + i r \sin \theta$$

$$|zw| = |z| |w|, \quad \text{and} \quad \left| \frac{z}{w} \right| = \frac{|z|}{|w|}$$

$$\arg(zw) = \arg z + \arg w, \quad \text{and} \quad \arg\left(\frac{z}{w}\right) = \arg z - \arg w$$

Euler's Relation $e^{i\theta}$

$$z = e^{i\theta} = \cos \theta + i \sin \theta$$

$$\frac{1}{z} = e^{-i\theta} = \cos \theta - i \sin \theta$$

Example: Express $5e^{\left(\frac{i3\pi}{4}\right)}$ in the form $x + iy$.

Solution: $5e^{\left(\frac{i3\pi}{4}\right)} = 5\left(\cos\left(\frac{3\pi}{4}\right) + i \sin\left(\frac{3\pi}{4}\right)\right)$

$$= \frac{-5\sqrt{2}}{2} + i \frac{5\sqrt{2}}{2}$$

Multiplying and dividing in mod-arg form

$$re^{i\theta} \times se^{i\phi} = rs e^{i(\theta+\phi)}$$

$$\equiv (r \cos \theta + i r \sin \theta) \times (s \cos \phi + i s \sin \phi) = rs \cos(\theta + \phi) + i rs \sin(\theta + \phi)$$

and

$$re^{i\theta} \div se^{i\phi} = \frac{r}{s} e^{i(\theta-\phi)}$$

$$\equiv (r \cos \theta + i r \sin \theta) \div (s \cos \phi + i s \sin \phi) = \frac{r}{s} \cos(\theta - \phi) + i \frac{r}{s} \sin(\theta - \phi)$$

De Moivre's Theorem

$$(re^{i\theta})^n = r^n e^{in\theta} \equiv (r \cos \theta + i r \sin \theta)^n = (r^n \cos n\theta + i r^n \sin n\theta)$$

Applications of De Moivre's Theorem

Example: Express $\sin 5\theta$ in terms of $\sin \theta$ only.

Solution: From De Moivre's Theorem we know that

$$\begin{aligned} \cos 5\theta + i \sin 5\theta &= (\cos \theta + i \sin \theta)^5 \\ &= \cos^5 \theta + 5i \cos^4 \theta \sin \theta + 10i^2 \cos^3 \theta \sin^2 \theta + 10i^3 \cos^2 \theta \sin^3 \theta + 5i^4 \cos \theta \sin^4 \theta + i^5 \sin^5 \theta \end{aligned}$$

Equating imaginary parts

$$\begin{aligned} \Rightarrow \sin 5\theta &= 5\cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin^5 \theta \\ &= 5(1 - \sin^2 \theta)^2 \sin \theta - 10(1 - \sin^2 \theta) \sin^3 \theta + \sin^5 \theta \\ &= 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta \end{aligned}$$

$$z^n + \frac{1}{z^n} = 2 \cos n\theta \quad \text{and} \quad z^n - \frac{1}{z^n} = 2i \sin n\theta$$

$$z = \cos \theta + i \sin \theta$$

$$\Rightarrow z^n = (\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$$

$$\text{and} \quad \frac{1}{z^n} = z^{-n} = (\cos \theta + i \sin \theta)^{-n} = (\cos n\theta - i \sin n\theta)$$

from which we can show that

$$\left(z + \frac{1}{z}\right) = 2 \cos \theta \quad \text{and} \quad \left(z - \frac{1}{z}\right) = 2i \sin \theta$$

$$z^n + \frac{1}{z^n} = 2 \cos n\theta \quad \text{and} \quad z^n - \frac{1}{z^n} = 2i \sin n\theta$$

Example: Express $\sin^5 \theta$ in terms of $\sin 5\theta$, $\sin 3\theta$ and $\sin \theta$.

Solution: Here we are dealing with $\sin \theta$, so we use

$$\begin{aligned} (2i \sin \theta)^5 &= \left(z - \frac{1}{z}\right)^5 \\ \Rightarrow 32i^5 \sin^5 \theta &= z^5 - 5z^4 \left(\frac{1}{z}\right) + 10z^3 \left(\frac{1}{z^2}\right) - 10z^2 \left(\frac{1}{z^3}\right) + 5z \left(\frac{1}{z^4}\right) - \left(\frac{1}{z^5}\right) \\ \Rightarrow 32i \sin^5 \theta &= \left(z^5 - \frac{1}{z^5}\right) - 5 \left(z^3 - \frac{1}{z^3}\right) + 10 \left(z - \frac{1}{z}\right) \\ \Rightarrow 32i \sin^5 \theta &= 2i \sin 5\theta - 5 \times 2i \sin 3\theta + 10 \times 2i \sin \theta \\ \Rightarrow \sin^5 \theta &= \frac{1}{16} (\sin 5\theta - 5 \sin 3\theta + 10 \sin \theta) \end{aligned}$$

n^{th} roots of a complex number

The technique is the same for finding n^{th} roots of any complex number.

Example: Find the 4th roots of $8\sqrt{2} + 8\sqrt{2}i$, and show the roots on an Argand Diagram.

Solution: We need to solve the equation $z^4 = 8\sqrt{2} + 8\sqrt{2}i$

1. Let $z = r \cos \theta + i r \sin \theta$

$$\Rightarrow z^4 = r^4 (\cos 4\theta + i \sin 4\theta)$$

2. $|8\sqrt{2} + 8\sqrt{2}i| = 8\sqrt{2+2} = 16$ and $\arg(8\sqrt{2} + 8\sqrt{2}i) = \frac{\pi}{4}$

$$\Rightarrow 8\sqrt{2} + 8\sqrt{2}i = 16 \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$$

3. Then $z^4 = 8\sqrt{2} + 8\sqrt{2}i$


$$\begin{aligned}
 \text{becomes } r^4 (\cos 4\theta + i \sin 4\theta) &= 16 \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) \\
 &= 16 \left(\cos \frac{9\pi}{4} + i \sin \frac{9\pi}{4} \right) && \text{adding } 2\pi \\
 &= 16 \left(\cos \frac{17\pi}{4} + i \sin \frac{17\pi}{4} \right) && \text{adding } 2\pi \\
 &= 16 \left(\cos \frac{25\pi}{4} + i \sin \frac{25\pi}{4} \right) && \text{adding } 2\pi
 \end{aligned}$$

4. $\Rightarrow r^4 = 16$ and $4\theta = \frac{\pi}{4}, \frac{9\pi}{4}, \frac{17\pi}{4}, \frac{25\pi}{4}$
 $\Rightarrow r = 2$ and $\theta = \frac{\pi}{16}, \frac{9\pi}{16}, \frac{17\pi}{16}, \frac{25\pi}{16} = \frac{-15\pi}{16}, \frac{25\pi}{16} = \frac{-7\pi}{16}; \quad -\pi < \arg z \leq \pi$

5. \Rightarrow roots are

$$\begin{aligned}
 z_1 &= 2 \left(\cos \frac{\pi}{16} + i \sin \frac{\pi}{16} \right) = 1.962 + 0.390i \\
 z_2 &= 2 \left(\cos \frac{9\pi}{16} + i \sin \frac{9\pi}{16} \right) = -0.390 + 1.962i \\
 z_3 &= 2 \left(\cos \frac{-15\pi}{16} + i \sin \frac{-15\pi}{16} \right) = -1.962 - 0.390i \\
 z_4 &= 2 \left(\cos \frac{-7\pi}{16} + i \sin \frac{-7\pi}{16} \right) = 0.390 - 1.962i
 \end{aligned}$$

Notice that the roots are symmetrically placed around the origin, and the angle between roots is $\frac{2\pi}{4} = \frac{\pi}{2}$.
 The angle between the n^{th} roots will always be $\frac{2\pi}{n}$.



For sixth roots the angle between roots will be $\frac{2\pi}{6} = \frac{\pi}{3}$, and so on.

Roots of polynomial equations with real coefficients

1. **Any** polynomial equation with real coefficients,
 $a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_2 x^2 + a_1 x + a_0 = 0$, (I)
 where all a_i are real, has a complex solution
2. \Rightarrow **any** complex n^{th} degree polynomial can be factorised into n linear factors over the complex numbers
3. If $z = a + ib$ is a root of (I), then its conjugate, $a - ib$ is also a root – see FP1.
4. By pairing factors with conjugate pairs we can say that any polynomial with real coefficients can be factorised into a combination of linear and quadratic factors over the real numbers.

Example: Given that $3 - 2i$ is a root of $z^3 - 5z^2 + 7z + 13 = 0$

- (a) Factorise over the real numbers
- (b) Find all three real roots

Solution:

- (a) $3 - 2i$ is a root $\Rightarrow 3 + 2i$ is also a root
 $\Rightarrow (z - (3 - 2i))(z - (3 + 2i)) = (z^2 - 6z + 13)$ is a factor
 $\Rightarrow z^3 - 5z^2 + 7z + 13 = (z^2 - 6z + 13)(z + 1)$ by inspection
- (b) \Rightarrow roots are $z = 3 - 2i, 3 + 2i$ and -1

Loci on an Argand Diagram

Two basic ideas

1. $|z - w|$ is the distance from w to z .
2. $\arg(z - (1 + i))$ is the angle made by the *half* line joining $(1 + i)$ to z , with the x -axis.

Example 1:

$|z - 2 - i| = 3$ is a circle with centre $(2 + i)$ and radius 3


Example 2:

$$|z + 3 - i| = |z - 2 + i|$$

$$\Leftrightarrow |z - (-3 + i)| = |z - (2 - i)|$$


is the locus of all points which are equidistant from the points

$A(-3, 1)$ and $B(2, -1)$, and so is the perpendicular bisector of AB .



Example 3:

$\arg(z - 4) = \frac{5\pi}{6}$ is a half line, from $(4, 0)$,
making an angle of $\frac{5\pi}{6}$ with the x -axis.



Example 4:

$|z - 3| = 2|z + 2i|$ is a circle
(Apollonius's circle).

To find its equation, put $z = x + iy$

$$\Rightarrow |(x - 3) + iy| = 2|x + i(y + 2)|$$

square both sides

$$\Rightarrow (x - 3)^2 + y^2 = 4(x^2 + (y + 2)^2)$$

leading to

$$\Rightarrow 3x^2 + 6x + 3y^2 + 16y + 7 = 0$$

$$\Rightarrow (x + 1)^2 + \left(y + \frac{8}{3}\right)^2 = \frac{52}{9}$$

which is a circle with centre $(-1, -\frac{8}{3})$, and radius $\frac{2\sqrt{13}}{3}$.


Example 5:

$$\arg\left(\frac{z-2}{z+5}\right) = \frac{\pi}{6}$$

$$\Rightarrow \arg(z - 2) - \arg(z + 5) = \frac{\pi}{6}$$

$$\Rightarrow \theta - \phi = \frac{\pi}{6}$$

which gives the arc of the circle as shown.




N.B.

The corresponding arc below the x -axis
would have equation

$$\arg\left(\frac{z-2}{z+5}\right) = -\frac{\pi}{6}$$

as $\theta - \phi$ would be negative in this picture.

(θ is a 'larger negative number' than ϕ .)



Transformations of the Complex Plane

Always start from the z -plane and transform to the w -plane, $z = x + iy$ and $w = u + iv$.

Example 1: Find the image of the circle $|z - 5| = 3$
under the transformation $w = \frac{1}{z-2}$.

Solution: First rearrange to find z

$$w = \frac{1}{z-2} \Rightarrow z - 2 = \frac{1}{w} \Rightarrow z = \frac{1}{w} + 2$$

Second substitute in equation of circle

$$\Rightarrow \left| \frac{1}{w} + 2 - 5 \right| = 3 \Rightarrow \left| \frac{1-3w}{w} \right| = 3$$

$$\Rightarrow |1 - 3w| = 3|w| \Rightarrow 3 \left| \frac{1}{3} - w \right| = 3|w|$$

$$\Rightarrow \left| w - \frac{1}{3} \right| = |w|$$

which is the equation of the perpendicular bisector of the line joining 0 to $\frac{1}{3}$,

$$\Rightarrow \text{the image is the line } u = \frac{1}{6}$$

Always consider the ‘modulus technique’ (above) first;

if this does not work then use the $u + iv$ method shown below.

Example 2: Show that the image of the line $x + 4y = 4$ under the transformation
 $w = \frac{1}{z-3}$ is a circle, and find its centre and radius.

Solution: First rearrange to find $z \Rightarrow z = \frac{1}{w} + 3$

The ‘modulus technique’ is not suitable here.

$$z = x + iy \quad \text{and} \quad w = u + iv$$

$$\Rightarrow z = \frac{1}{w} + 3 = \frac{1}{u+iv} + 3 = \frac{1}{u+iv} \times \frac{u-iv}{u-iv} + 3$$

$$\Rightarrow x + iy = \frac{u-iv}{u^2+v^2} + 3$$

$$\text{Equating real and imaginary parts } x = \frac{u}{u^2+v^2} + 3 \text{ and } y = \frac{-v}{u^2+v^2}$$

$$\Rightarrow x + 4y = 4 \text{ becomes } \frac{u}{u^2+v^2} + 3 - \frac{4v}{u^2+v^2} = 4$$

$$\Rightarrow u^2 - u + v^2 + 4v = 0$$

$$\Rightarrow \left(u - \frac{1}{2}\right)^2 + (v + 2)^2 = \frac{17}{4}$$

which is a circle with centre $\left(\frac{1}{2}, -2\right)$ and radius $\frac{\sqrt{17}}{2}$.

There are many more examples in the book, but these are the two important techniques.

Loci and geometry

It is always important to think of diagrams.

Example: z lies on the circle $|z - 2i| = 1$.
Find the greatest and least values of $\arg z$.

Solution: Draw a picture!


The greatest and least values of $\arg z$ will occur at B and A .

Trigonometry tells us that

$$\theta = \frac{\pi}{6}$$

and so greatest and least values of

$$\arg z \text{ are } \frac{2\pi}{3} \text{ and } \frac{\pi}{3}$$



4 First Order Differential Equations

Separating the variables, families of curves

Example: Find the general solution of

$$\frac{dy}{dx} = \frac{y}{2x(x+1)}, \quad \text{for } x > 0,$$


and sketch some of the family of solution curves.

Solution:
$$\frac{dy}{dx} = \frac{y}{2x(x+1)} \Rightarrow \int \frac{2}{y} dy = \int \frac{1}{x(x+1)} dx = \int \frac{1}{x} - \frac{1}{x+1} dx$$

$$\Rightarrow 2 \ln y = \ln x - \ln(x+1) + \ln A$$

$$\Rightarrow y^2 = \frac{Ax}{x+1}$$

Thus for varying values of A and for $x > 0$, we have



Exact Equations

In an exact equation the L.H.S. is an exact derivative (really a preparation for Integrating Factors).

Example: Solve $\sin x \frac{dy}{dx} + y \cos x = 3x^2$

Solution: Notice that the L.H.S. is an exact derivative

$$\sin x \frac{dy}{dx} + y \cos x = \frac{d}{dx}(y \sin x)$$

$$\Rightarrow \frac{d}{dx}(y \sin x) = 3x^2$$

$$\Rightarrow y \sin x = \int 3x^2 dx = x^3 + c$$

$$\Rightarrow y = \frac{x^3 + c}{\sin x}$$

Integrating Factors

$$\frac{dy}{dx} + Py = Q \quad \text{where } P \text{ and } Q \text{ are functions of } x \text{ only.}$$

In this case, multiply both sides by an Integrating Factor, $R = e^{\int P dx}$.

The L.H.S. will now be an exact derivative, $\frac{d}{dx}(Ry)$.

Proceed as in the above example.

Example: Solve $x \frac{dy}{dx} + 2y = 1$

Solution: First divide through by x

$$\Rightarrow \frac{dy}{dx} + \frac{2}{x}y = \frac{1}{x} \quad \text{now in the correct form}$$

Integrating Factor, I.F., is $R = e^{\int P dx} = e^{\int \frac{2}{x} dx} = e^{2 \ln x} = x^2$

$$\Rightarrow x^2 \frac{dy}{dx} + 2xy = x \quad \text{multiplying by } x^2$$

$$\Rightarrow \frac{d}{dx}(x^2 y) = x, \quad \text{check that it is an exact derivative}$$

$$\Rightarrow x^2 y = \int x dx = \frac{x^2}{2} + c$$

$$\Rightarrow y = \frac{1}{2} + \frac{c}{x^2}$$

Using substitutions

Example 1: Use the substitution $y = vx$ (where v is a function of x) to solve the equation

$$\frac{dy}{dx} = \frac{3yx^2 + y^3}{x^3 + xy^2}$$

Solution: $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

$$\Rightarrow \frac{dy}{dx} = \frac{3yx^2 + y^3}{x^3 + xy^2} \Rightarrow v + x \frac{dv}{dx} = \frac{3(vx)x^2 + (vx)^3}{x^3 + x(vx)^2} = \frac{3v + v^3}{1 + v^2}$$

and we can now separate the variables

$$\Rightarrow x \frac{dv}{dx} = \frac{3v + v^3}{1 + v^2} - v = \frac{3v + v^3 - v - v^3}{1 + v^2} = \frac{2v}{1 + v^2}$$

$$\Rightarrow \frac{1 + v^2}{2v} \frac{dv}{dx} = \frac{1}{x}$$

$$\Rightarrow \int \frac{1}{2v} + \frac{v}{2} dv = \int \frac{1}{x} dx$$

$$\Rightarrow \frac{1}{2} \ln v + \frac{v^2}{4} = \ln x + c$$

$$\text{But } v = \frac{y}{x}, \Rightarrow \frac{1}{2} \ln \frac{y}{x} + \frac{y^2}{4x^2} = \ln x + c$$

$$\Rightarrow 2x^2 \ln y + y^2 = 6x^2 \ln x + c' x^2 \quad c' \text{ is new arbitrary constant}$$

and I would not like to find $y!!!$

If told to use the substitution $v = \frac{y}{x}$, rewrite as $y = vx$ and proceed as in the above example.

Example 2: Use the substitution $y = \frac{1}{z}$ to solve the differential equation

$$\frac{dy}{dx} = y^2 + y \cot x.$$

Solution: $y = \frac{1}{z} \Rightarrow \frac{dy}{dx} = \frac{-1}{z^2} \frac{dz}{dx}$

$$\Rightarrow \frac{-1}{z^2} \frac{dz}{dx} = \frac{1}{z^2} + \frac{1}{z} \cot x$$

$$\Rightarrow \frac{dz}{dx} + z \cot x = -1$$

Integrating factor is $R = e^{\int \cot x \, dx} = e^{\ln(\sin x)} = \sin x$

$$\Rightarrow \sin x \frac{dz}{dx} + z \cos x = -\sin x$$

$$\Rightarrow \frac{d}{dx}(z \sin x) = -\sin x \quad \text{check that it is an exact derivative}$$

$$\Rightarrow z \sin x = \cos x + c$$

$$\Rightarrow z = \frac{\cos x + c}{\sin x} \quad \text{but } z = \frac{1}{y}$$

$$\Rightarrow y = \frac{\sin x}{\cos x + c}$$

Example 3: Use the substitution $z = x + y$ to solve the differential equation

$$\frac{dy}{dx} = \cos(x + y)$$

Solution: $z = x + y \Rightarrow \frac{dz}{dx} = 1 + \frac{dy}{dx}$

$$\Rightarrow \frac{dz}{dx} = 1 + \cos z$$

$$\Rightarrow \int \frac{1}{1 + \cos z} \, dz = \int dx \quad \text{separating the variables}$$

$$\Rightarrow \int \frac{1}{2} \sec^2\left(\frac{z}{2}\right) \, dz = x + c \quad 1 + \cos z = 1 + 2 \cos^2\left(\frac{z}{2}\right) - 1 = 2 \cos^2\left(\frac{z}{2}\right)$$

$$\Rightarrow \tan\left(\frac{z}{2}\right) = x + c$$

But $z = x + y \Rightarrow \tan\left(\frac{x+y}{2}\right) = x + c$

5 Second Order Differential Equations

Linear with constant coefficients

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = f(x) \quad \text{where } a, b \text{ and } c \text{ are constants.}$$

(1) when $f(x) = 0$

First write down the Auxiliary Equation, A.E

$$\text{A.E. } am^2 + bm + c = 0$$

and solve to find the roots $m = \alpha$ or β

- (i) If α and β are both real numbers, and if $\alpha \neq \beta$
then the Complimentary Function, C.F., is
 $y = A e^{\alpha x} + B e^{\beta x}$, where A and B are arbitrary constants of integration
- (ii) If α and β are both real numbers, and if $\alpha = \beta$
then the Complimentary Function, C.F., is
 $y = (A + Bx) e^{\alpha x}$, where A and B are arbitrary constants of integration
- (iii) If α and β are both complex numbers, and if $\alpha = a + ib$, $\beta = a - ib$
then the Complimentary Function, C.F.,
 $y = e^{ax}(A \sin bx + B \cos bx)$,
where A and B are arbitrary constants of integration

Example 1: Solve $\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} - 3y = 0$

Solution: A.E. is $m^2 + 2m - 3 = 0$

$$\Rightarrow (m - 1)(m + 3) = 0$$

$$\Rightarrow m = 1 \text{ or } -3$$

$$\Rightarrow y = A e^x + B e^{-3x}$$

when $f(x) = 0$, the C.F. is the solution

Example 2: Solve $\frac{d^2 y}{dx^2} + 6 \frac{dy}{dx} + 9y = 0$

Solution: A.E. is $m^2 + 6m + 9 = 0$

$$\Rightarrow (m + 3)^2 = 0$$

$$\Rightarrow m = -3 \text{ (and } -3)$$

$$\Rightarrow y = (A + Bx) e^{-3x}$$

repeated root

when $f(x) = 0$, the C.F. is the solution

Example 3: Solve $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 13y = 0$

Solution: A.E. is $m^2 + 4m + 13 = 0$

$$\Rightarrow (m + 2)^2 = -9 = (3i)^2$$

$$\Rightarrow (m + 2) = \pm 3i$$

$$\Rightarrow m = -2 - 3i \text{ or } -2 + 3i$$

$$\Rightarrow y = e^{-2x}(A \sin 3x + B \cos 3x)$$

when $f(x) = 0$, the C.F. is the solution

(2) when $f(x) \neq 0$, Particular Integrals

First proceed as in (1) to find the Complimentary Function, then use the rules below to find a Particular Integral, P.I.

Second the General Solution, G.S., is found by adding the C.F. and the P.I.

$$\Rightarrow \text{G.S.} = \text{C.F.} + \text{P.I.}$$

Note that it does not matter what P.I. you use, so you might as well find the easiest, which is what these rules do.

(1) $f(x) = e^{kx}$.

Try $y = Ae^{kx}$

unless e^{kx} appears, on its own, in the C.F., in which case try $y = Cxe^{kx}$

unless xe^{kx} appears, on its own, in the C.F., in which case try $y = Cx^2e^{kx}$.

(2) $f(x) = \sin kx$ or $f(x) = \cos kx$

Try $y = C \sin kx + D \cos kx$

unless $\sin kx$ or $\cos kx$ appear in the C.F., on their own, in which case

try $y = x(C \sin kx + D \cos kx)$

(3) $f(x) = \text{a polynomial of degree } n$.

Try $f(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_1 x + a_0$

unless a number, on its own, appears in the C.F., in which case

try $f(x) = x(a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_1 x + a_0)$

i.e. try $f(x) = \text{a polynomial of degree } n$.

(4) **In general**

to find a P.I., try something like $f(x)$, unless this appears in the C.F. (or if there is a problem), then try something like $xf(x)$.

Example 1: Solve $\frac{d^2y}{dx^2} + 6\frac{dy}{dx} + 5y = 2x$

Solution: A.E. is $m^2 + 6m + 5 = 0$

$$\Rightarrow (m+5)(m+1) = 0 \Rightarrow m = -5 \text{ or } -1$$

$$\Rightarrow \text{C.F. is } y = Ae^{-5x} + Be^{-x}$$

For the P.I., try $y = Cx + D$

$$\Rightarrow \frac{dy}{dx} = C \text{ and } \frac{d^2y}{dx^2} = 0$$

Substituting in the differential equation gives

$$0 + 6C + 5(Cx + D) = 2x$$

$$\Rightarrow 5C = 2$$

comparing coefficients of x

$$\Rightarrow C = \frac{2}{5}$$

$$\text{and } 6C + 5D = 0$$

comparing constant terms

$$\Rightarrow D = \frac{-12}{25}$$

$$\Rightarrow \text{P.I. is } y = \frac{2}{5}x - \frac{12}{25}$$

$$\Rightarrow \text{G.S. is } y = Ae^{-5x} + Be^{-x} + \frac{2}{5}x - \frac{12}{25}$$

Example 2: Solve $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 9y = e^{3x}$

Solution: A.E. is $m^2 - 6m + 9 = 0$

$$\Rightarrow (m-3)^2 = 0$$

$$\Rightarrow m = 3$$

repeated root

$$\Rightarrow \text{C.F. is } y = (Ax + B)e^{3x}$$

In this case, both e^{3x} and xe^{3x} appear, on their own, in the C.F., so for a P.I. we try $y = Cx^2e^{3x}$

$$\Rightarrow \frac{dy}{dx} = 2Cxe^{3x} + 3Cx^2e^{3x}$$

$$\text{and } \frac{d^2y}{dx^2} = 2Ce^{3x} + 6Cxe^{3x} + 6Cxe^{3x} + 9Cx^2e^{3x}$$

Substituting in the differential equation gives

$$2Ce^{3x} + 12Cxe^{3x} + 9Cx^2e^{3x} - 6(2Cxe^{3x} + 3Cx^2e^{3x}) + 9Cx^2e^{3x} = e^{3x}$$

$$\Rightarrow 2Ce^{3x} = e^{3x}$$

$$\Rightarrow C = \frac{1}{2}$$

$$\Rightarrow \text{P.I. is } y = \frac{1}{2}x^2e^{3x}$$

$$\Rightarrow \text{G.S. is } y = (Ax + B)e^{3x} + \frac{1}{2}x^2e^{3x}$$

Example 3: Solve $\frac{d^2x}{dt^2} - 3\frac{dx}{dt} + 2x = 4 \cos 2t$
 given that $x = 0$ and $\dot{x} = 1$ when $t = 0$.

Solution: A.E. is $m^2 - 3m + 2 = 0$

$$\Rightarrow m = 1 \text{ or } 2$$

$$\Rightarrow \text{C.F. is } x = Ae^t + Be^{2t}$$

For the P.I. try $x = C \sin 2t + D \cos 2t$

BOTH $\sin 2t$ AND $\cos 2t$ are needed

$$\Rightarrow \dot{x} = 2C \cos 2t - 2D \sin 2t$$

$$\text{and } \ddot{x} = -4C \sin 2t - 4D \cos 2t$$

Substituting in the differential equation gives

$$(-4C \sin 2t - 4D \cos 2t) - 3(2C \cos 2t - 2D \sin 2t) + 2(C \sin 2t + D \cos 2t) = 4 \cos 2t$$

$$\Rightarrow -2C + 6D = 0 \quad \Rightarrow -C + 3D = 0 \quad \text{comparing coefficients of } \sin 2t$$

$$\text{and } -6C - 2D = 4 \quad \Rightarrow 3C + D = -2 \quad \text{comparing coefficients of } \cos 2t$$

$$\Rightarrow C = \frac{-3}{5} \text{ and } D = \frac{-1}{5}$$

$$\Rightarrow \text{P.I. is } x = -\frac{3}{5} \sin 2t - \frac{1}{5} \cos 2t$$

$$\Rightarrow \text{G.S. is } x = Ae^t + Be^{2t} - \frac{3}{5} \sin 2t - \frac{1}{5} \cos 2t$$

$$\Rightarrow \dot{x} = Ae^t + 2Be^{2t} - \frac{6}{5} \cos 2t + \frac{2}{5} \sin 2t$$

$$x = 0 \text{ and when } t = 0 \quad \Rightarrow 0 = A + B - \frac{1}{5}$$

$$\text{and } \dot{x} = 1 \text{ when } t = 0 \quad \Rightarrow 1 = A + 2B - \frac{6}{5}$$

$$\Rightarrow A = \frac{-9}{5} \text{ and } B = 2$$

$$\Rightarrow \text{solution is } x = \frac{-9}{5} e^t + 2e^{2t} - \frac{6}{5} \sin 2t - \frac{2}{5} \cos 2t$$

D.E.s of the form $ax^2 \frac{d^2y}{dx^2} + bx \frac{dy}{dx} + cy = f(x)$

Substitute $x = e^u$

$$\Rightarrow \frac{dx}{du} = e^u = x \Rightarrow \frac{du}{dx} = \frac{1}{x}$$

and $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{x} \frac{dy}{du} \Leftrightarrow \quad \mathbf{x \frac{dy}{dx} = \frac{dy}{du}} \quad \mathbf{I}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{1}{x^2} \frac{dy}{du} + \frac{1}{x} \frac{d\left(\frac{dy}{du}\right)}{dx}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{1}{x^2} \frac{dy}{du} + \frac{1}{x} \frac{d\left(\frac{dy}{du}\right)}{du} \frac{du}{dx} \quad \text{chain rule}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{1}{x^2} \frac{dy}{du} + \frac{1}{x^2} \frac{d^2y}{du^2}$$

$$\Rightarrow \quad \mathbf{x^2 \frac{d^2y}{dx^2} = \frac{d^2y}{du^2} - \frac{dy}{du}} \quad \mathbf{II}$$

Thus we have $x^2 \frac{d^2y}{dx^2} = \frac{d^2y}{du^2} - \frac{dy}{du}$ and $x \frac{dy}{dx} = \frac{dy}{du}$ from **I** and **II**

substituting these in the original equation leads to a second order D.E. with constant coefficients.

Example: Solve the differential equation $x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 3y = -2x^2$.

Solution: Using the substitution $x = e^u$, and proceeding as above

$$x^2 \frac{d^2y}{dx^2} = \frac{d^2y}{du^2} - \frac{dy}{du} \quad \text{and} \quad x \frac{dy}{dx} = \frac{dy}{du}$$

$$\Rightarrow \frac{d^2y}{du^2} - \frac{dy}{du} - 3 \frac{dy}{du} + 3y = -2e^{2u}$$

$$\Rightarrow \frac{d^2y}{du^2} - 4 \frac{dy}{du} + 3y = -2e^{2u}$$

$$\Rightarrow \text{A.E. is } m^2 - 4m + 3 = 0$$

$$\Rightarrow (m-3)(m-1) = 0 \Rightarrow m = 3 \text{ or } 1$$

$$\Rightarrow \text{C.F. is } y = Ae^{3u} + Be^u$$

For the P.I. try $y = Ce^{2u}$

$$\Rightarrow \frac{dy}{du} = 2Ce^{2u} \quad \text{and} \quad \frac{d^2y}{du^2} = 4Ce^{2u}$$

$$\Rightarrow 4Ce^{2u} - 8Ce^{2u} + 3Ce^{2u} = -2e^{2u}$$

$$\Rightarrow C = 2$$

$$\Rightarrow \text{G.S. is } y = Ae^{3u} + Be^u + 2e^{2u}$$

But $x = e^u \Rightarrow \text{G.S. is } y = Ax^3 + Bx + 2x^2$

6 Maclaurin and Taylor Series

1) Maclaurin series

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \frac{x^3}{3!}f'''(0) + \dots + \frac{x^n}{n!}f^n(0) + \dots$$

2) Taylor series

$$f(x+a) = f(a) + xf'(a) + \frac{x^2}{2!}f''(a) + \frac{x^3}{3!}f'''(a) + \dots + \frac{x^n}{n!}f^n(a) + \dots$$

3) Taylor series – as a power series in $(x-a)$

replacing x by $(x-a)$ in 2) we get

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!}f''(a) + \frac{(x-a)^3}{3!}f'''(a) + \dots + \frac{(x-a)^n}{n!}f^n(a) + \dots$$

4) Solving differential equations using Taylor series

(a) If we are given the value of y when $x=0$, then we use the Maclaurin series with

$$f(0) = y_0 \quad \text{the value of } y \text{ when } x=0$$

$$f'(0) = \left(\frac{dy}{dx}\right)_0 \quad \text{the value of } \frac{dy}{dx} \text{ when } x=0$$

etc. to give

$$f(x) = y = y_0 + x\left(\frac{dy}{dx}\right)_0 + \frac{x^2}{2!}\left(\frac{d^2y}{dx^2}\right)_0 + \frac{x^3}{3!}\left(\frac{d^3y}{dx^3}\right)_0 + \dots + \frac{x^n}{n!}\left(\frac{d^ny}{dx^n}\right)_0 + \dots$$

(b) If we are given the value of y when $x=a$, then we use the Taylor power series with

$$f(a) = y_a \quad \text{the value of } y \text{ when } x=a$$

$$f'(a) = \left(\frac{dy}{dx}\right)_a \quad \text{the value of } \frac{dy}{dx} \text{ when } x=a$$

etc. to give

$$y = y_a + (x-a)\left(\frac{dy}{dx}\right)_a + \frac{(x-a)^2}{2!}\left(\frac{d^2y}{dx^2}\right)_a + \frac{(x-a)^3}{3!}\left(\frac{d^3y}{dx^3}\right)_a + \dots$$

NOTE THAT 4 (a) and 4 (b) are not in the formula book, but can easily be found using the results in 1) and 3).

Standard series

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots \quad \text{converges for all real } x$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} + \dots \quad \text{converges for all real } x$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^{n-1} \frac{x^{2n-2}}{(2n-2)!} + \dots \quad \text{converges for all real } x$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^{n-1} \frac{x^n}{n} + \dots \quad \text{converges for } -1 < x \leq 1$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \quad \text{converges for } -1 < x < 1$$

Example 1: Find the Maclaurin series for $f(x) = \tan x$, up to and including the term in x^3

Solution: $f(x) = \tan x \quad \Rightarrow \quad f(0) = 0$

$$\Rightarrow f'(x) = \sec^2 x \quad \Rightarrow \quad f'(0) = 1$$

$$\Rightarrow f''(x) = 2 \sec^2 x \tan x \quad \Rightarrow \quad f''(0) = 0$$

$$\Rightarrow f'''(x) = 4 \sec^2 x \tan^2 x + 2 \sec^4 x \quad \Rightarrow \quad f'''(0) = 2$$

and $f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots + \frac{x^n}{n!} f^n(0) + \dots$

$$\Rightarrow \tan x \cong 0 + x \times 1 + \frac{x^2}{2!} \times 0 + \frac{x^3}{3!} \times 2 \quad \text{up to the term in } x^3$$

$$\Rightarrow \tan x \cong x + \frac{x^3}{3}$$

Example 2: Using the Maclaurin series for e^x to find an expansion of e^{x+x^2} , up to and including the term in x^3 .

Solution: $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

$$\Rightarrow e^{x+x^2} \cong 1 + (x+x^2) + \frac{(x+x^2)^2}{2!} + \frac{(x+x^2)^3}{3!} \quad \text{up to the term in } x^3$$

$$\cong 1 + x + x^2 + \frac{x^2+2x^3+\dots}{2!} + \frac{x^3+\dots}{3!} \quad \text{up to the term in } x^3$$

$$\Rightarrow e^{x+x^2} \cong 1 + x + \frac{3}{2}x^2 + \frac{7}{6}x^3 \quad \text{up to the term in } x^3$$

Example 3: Find a Taylor series for $\cot\left(x + \frac{\pi}{4}\right)$, up to and including the term in x^2 .

Solution: $f(x) = \cot x$ and we are looking for

$$f\left(x + \frac{\pi}{4}\right) = f\left(\frac{\pi}{4}\right) + xf'\left(\frac{\pi}{4}\right) + \frac{x^2}{2!}f''\left(\frac{\pi}{4}\right)$$

$$f(x) = \cot x \quad \Rightarrow \quad f\left(\frac{\pi}{4}\right) = 1$$

$$\Rightarrow f'(x) = -\operatorname{cosec}^2 x \quad \Rightarrow \quad f'\left(\frac{\pi}{4}\right) = -2$$

$$\Rightarrow f''(x) = 2\operatorname{cosec}^2 x \cot x \quad \Rightarrow \quad f''\left(\frac{\pi}{4}\right) = 4$$

$$\Rightarrow \cot\left(x + \frac{\pi}{4}\right) \cong 1 - 2x + \frac{x^2}{2!} \times 4 \quad \text{up to the term in } x^2$$

$$\Rightarrow \cot\left(x + \frac{\pi}{4}\right) \cong 1 - 2x + 2x^2 \quad \text{up to the term in } x^2$$

Example 4: Use a Taylor series to solve the differential equation,

$$y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx}\right)^2 + y = 0 \quad \text{equation I}$$

up to and including the term in x^3 , given that $y = 1$ and $\frac{dy}{dx} = 2$ when $x = 0$.

In this case the initial value of x is 0, so we shall use

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \frac{x^3}{3!}f'''(0) + \dots + \frac{x^n}{n!}f^n(0) + \dots$$

$$\Leftrightarrow y = y_0 + x\left(\frac{dy}{dx}\right)_0 + \frac{x^2}{2!}\left(\frac{d^2 y}{dx^2}\right)_0 + \frac{x^3}{3!}\left(\frac{d^3 y}{dx^3}\right)_0.$$

We already know that $y_0 = 1$ and $\left(\frac{dy}{dx}\right)_0 = 2$ values when $x = 0$

$$\Rightarrow \left(\frac{d^2 y}{dx^2}\right)_0 = \left(-\frac{1}{y}\left(\frac{dy}{dx}\right)^2 - 1\right)_0 = -5 \quad \text{values when } x = 0$$

$$\text{equation I} \quad y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx}\right)^2 + y = 0$$

$$\text{Differentiating} \quad \Rightarrow \quad y \frac{d^3 y}{dx^3} + \frac{dy}{dx} \times \frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} \times \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 0$$

Substituting $y_0 = 1$, $\left(\frac{dy}{dx}\right)_0 = 2$ and $\left(\frac{d^2 y}{dx^2}\right)_0 = -5$ values when $x = 0$

$$\Rightarrow \left(\frac{d^3 y}{dx^3}\right)_0 + 2 \times (-5) + 2 \times 2 \times (-5) + 2 = 0$$

$$\Rightarrow \left(\frac{d^3 y}{dx^3}\right)_0 = 28$$

$$\Rightarrow \text{solution is } y \cong 1 + 2x + \frac{x^2}{2!} \times (-5) + \frac{x^3}{3!} \times 28$$

$$\Rightarrow y \cong 1 + 2x - \frac{5}{2}x^2 + \frac{14}{3}x^3$$

Series expansions of compound functions

Example: Find a polynomial expansion for

$$\frac{\cos 2x}{1-3x}, \quad \text{up to and including the term in } x^3.$$

Solution: Using the standard series

$$\cos 2x = 1 - \frac{(2x)^2}{2!} + \dots \quad \text{up to and including the term in } x^3$$

$$\begin{aligned} \text{and } (1-3x)^{-1} &= 1 + 3x + \frac{-1 \times -2}{2!}(-3x)^2 + \frac{-1 \times -2 \times -3}{3!}(-3x)^3 \\ &= 1 + 3x + 9x^2 + 27x^3 \quad \text{up to and including the term in } x^3 \end{aligned}$$

$$\begin{aligned} \Rightarrow \frac{\cos 2x}{1-3x} &= \left(1 - \frac{(2x)^2}{2!}\right)(1 + 3x + 9x^2 + 27x^3) \\ &= 1 + 3x + 9x^2 + 27x^3 - 2x^2 - 6x^3 \quad \text{up to and including the term in } x^3 \end{aligned}$$


$$\Rightarrow \frac{\cos 2x}{1-3x} = 1 + 3x + 7x^2 + 21x^3 \quad \text{up to and including the term in } x^3$$

7 Polar Coordinates

The polar coordinates of P are (r, θ)

$r = OP$, the distance from the origin or *pole*,

and θ is the angle made anti-clockwise with the initial line.



In the Edexcel syllabus r is always taken as positive or 0, and $0 \leq \theta < 2\pi$

(But in most books r can be negative, thus $(-4, \frac{\pi}{2})$ is the same point as $(4, \frac{3\pi}{2})$)


Polar and Cartesian coordinates

From the diagram

$$r = \sqrt{x^2 + y^2}$$

and $\tan \theta = \frac{y}{x}$ (use sketch to find θ).

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta.$$



Sketching curves

In practice, if you are asked to sketch a curve, it will probably be best to plot a few points. The important values of θ are those for which $r = 0$.


The sketches in these notes will show when r is negative by plotting a dotted line; these sections should be ignored as far as Edexcel A-level is concerned.

Some common curves

$$r = a + b \cos \theta$$


Cardioid

$$a = b$$




Limacon without dimple

$$a \geq 2b,$$




Limacon with a dimple

$$b \leq a < 2b$$




Limacon with a loop


$a < b$
 r negative in the loop




Circle




Half line




Line ($x = 3$)



Line ($y = 3$)




Circle




With Cartesian coordinates the graph of $y = f(x - a)$ is the graph of $y = f(x)$ translated through a in the x -direction.

In a similar way the graph of $r = 3 \sec(\theta - \alpha)$, or $r = 3 \sec(\alpha - \theta)$, is a rotation of the graph of $r = \sec \theta$ through α , anti-clockwise.

Line ($x = 3$ rotated through $\frac{\pi}{6}$)




Line ($y = 3$ rotated through $\frac{\pi}{6}$)



Rose Curves

$$r = 4 \cos 3\theta$$


$$0 \leq \theta < \pi$$



below x-axis, r negative

$$r = 4 \cos 3\theta$$


$$\pi \leq \theta < 2\pi$$



above x-axis, r negative

$$r = 4 \cos 3\theta$$

$$0 \leq \theta < 2\pi$$




whole curve for $r \geq 0$

The rose curve will always have n petals when n is odd, for $0 \leq \theta < 2\pi$.

$$r = 3 \cos 4\theta$$


When n is even there will be n petals for $r \geq 0$ and $0 \leq \theta < 2\pi$.




Thus, whether n is odd or even, the rose curve $r = a \cos \theta$ always has n petals, when only the positive (or 0) values of r are taken.

Edexcel only allow positive or 0 values of r .


Lemniscate of Bernoulli



Spiral $r = 2\theta$




Spiral $r = e^\theta$




Circle $r = 10 \cos \theta$

Notice that in the circle on OA as diameter, the angle P is 90° (angle in a semi-circle) and trigonometry gives us that $r = 10 \cos \theta$.



Circle $r = 10 \sin \theta$

In the same way $r = 10 \sin \theta$ gives a circle on the y -axis.



Areas using polar coordinates


Remember: area of a sector is $\frac{1}{2}r^2\theta$

$$\text{Area of } OPQ = \delta A \approx \frac{1}{2}r^2\delta\theta$$

$$\Rightarrow \text{Area } OAB \approx \sum \left(\frac{1}{2}r^2\delta\theta \right)$$

as $\delta\theta \rightarrow 0$


$$\Rightarrow \text{Area } OAB = \int_{\theta_1}^{\theta_2} \frac{1}{2}r^2 d\theta$$



Example: Find the area between the curve $r = 1 + \tan \theta$ and the half lines $\theta = 0$ and $\theta = \frac{\pi}{3}$

Solution:

$$\begin{aligned}
 \text{Area} &= \int_0^{\pi/3} \frac{1}{2} r^2 d\theta \\
 &= \int_0^{\pi/3} \frac{1}{2} (1 + \tan \theta)^2 d\theta \\
 &= \int_0^{\pi/3} \frac{1}{2} (1 + 2 \tan \theta + \tan^2 \theta) d\theta \\
 &= \int_0^{\pi/3} \frac{1}{2} (2 \tan \theta + \sec^2 \theta) d\theta \\
 &= \frac{1}{2} [2 \ln(\sec \theta) + \tan \theta]_0^{\pi/3} \\
 &= \ln 2 + \frac{\sqrt{3}}{2}
 \end{aligned}$$



Tangents parallel and perpendicular to the initial line

$$y = r \sin \theta \quad \text{and} \quad x = r \cos \theta$$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta}$$

- 1) Tangents will be parallel to the initial line ($\theta = 0$), or horizontal, when $\frac{dy}{dx} = 0$

$$\Rightarrow \frac{dy}{d\theta} = 0$$

$$\Rightarrow \frac{d}{d\theta} (r \sin \theta) = 0$$

- 2) Tangents will be perpendicular to the initial line ($\theta = 0$), or vertical, when $\frac{dy}{dx}$ is infinite

$$\Rightarrow \frac{dx}{d\theta} = 0$$

$$\Rightarrow \frac{d}{d\theta} (r \cos \theta) = 0$$

Note that if both $\frac{dy}{d\theta} = 0$ and $\frac{dx}{d\theta} = 0$, then $\frac{dy}{dx}$ is not defined, and you should look at a sketch to help (or use l'Hôpital's rule).

Example: Find the coordinates of the points on $r = 1 + \cos \theta$ where the tangents are
 (a) parallel to the initial line,
 (b) perpendicular to the initial line.

Solution: $r = 1 + \cos \theta$ is shown in the diagram.

(a) Tangents parallel to $\theta = 0$ (horizontal)

$$\Rightarrow \frac{dy}{d\theta} = 0 \Rightarrow \frac{d}{d\theta}(r \sin \theta) = 0$$

$$\Rightarrow \frac{d}{d\theta}((1 + \cos \theta) \sin \theta) = 0 \Rightarrow \frac{d}{d\theta}(\sin \theta + \sin \theta \cos \theta) = 0$$

$$\Rightarrow \cos \theta - \sin^2 \theta + \cos^2 \theta = 0 \Rightarrow 2 \cos^2 \theta + \cos \theta - 1 = 0$$

$$\Rightarrow (2 \cos \theta - 1)(\cos \theta + 1) = 0 \Rightarrow \cos \theta = \frac{1}{2} \text{ or } -1$$

$$\Rightarrow \theta = \pm \frac{\pi}{3} \text{ or } \pi$$

(b) Tangents perpendicular to $\theta = 0$ (vertical)

$$\Rightarrow \frac{dx}{d\theta} = 0 \Rightarrow \frac{d}{d\theta}(r \cos \theta) = 0$$

$$\Rightarrow \frac{d}{d\theta}((1 + \cos \theta) \cos \theta) = 0 \Rightarrow \frac{d}{d\theta}(\cos \theta + \cos^2 \theta) = 0$$

$$\Rightarrow -\sin \theta - 2 \cos \theta \sin \theta = 0 \Rightarrow \sin \theta (1 + 2 \cos \theta) = 0$$

$$\Rightarrow \cos \theta = -\frac{1}{2} \text{ or } \sin \theta = 0$$

$$\Rightarrow \theta = \pm \frac{2\pi}{3} \text{ or } 0, \pi$$


From the above we can see that

- (a) the tangent is parallel to $\theta = 0$
 at $B \left(\theta = \frac{\pi}{3} \right)$, and $E \left(\theta = -\frac{\pi}{3} \right)$,
 also at $\theta = \pi$, the origin – see below (c)

- (b) the tangent is perpendicular to $\theta = 0$
 at $A \left(\theta = 0 \right)$, $C \left(\theta = \frac{2\pi}{3} \right)$ and $D \left(\theta = \frac{-2\pi}{3} \right)$

- (c) we also have both $\frac{dx}{d\theta} = 0$ and $\frac{dy}{d\theta} = 0$ when $\theta = \pi$!!!

From the graph it looks as if the tangent is parallel to $\theta = 0$ at the origin, when $\theta = \pi$,
 and from l'Hôpital's rule it can be shown that this is true.



Appendix

n^{th} roots of 1

Short method

Example: Find the 5th roots of $-4 + 4i = 4\sqrt{2} e^{3\pi i/4}$

Solution: First find the root with the smallest argument

$$\left(4\sqrt{2} e^{3\pi i/4}\right)^{1/5} = \sqrt{2} e^{3\pi i/20}$$


Then sketch the symmetrical ‘spider’ diagram where the angle between successive roots is $2\pi/5 = 8\pi/20$

then find all five roots by successively adding $8\pi/20$ to the argument of each root

to give

$$\sqrt{2} e^{3\pi i/20}, \sqrt{2} e^{11\pi i/20}, \sqrt{2} e^{19\pi i/20},$$

$$\sqrt{2} e^{27\pi i/20} = \sqrt{2} e^{-13\pi i/20}, \text{ and } \sqrt{2} e^{35\pi i/20} = \sqrt{2} e^{-5\pi i/20}.$$




This can be generalized to find the n^{th} roots of any complex number, adding $2\pi/n$ successively to the argument of each root.

Warning: You must make sure that your method is very clear in an examination.

Sum of n^{th} roots of 1

Consider the solutions of $z^{10} = 1$, the complex 10^{th} roots of 1.

Suppose that ω is the complex 10^{th} root of 1 with the smallest argument. The 'spider' diagram shows that the roots are $\omega, \omega^2, \omega^3, \omega^4, \dots, \omega^9$ and 1.



Symmetry indicates that the sum of all these roots is a real number, but to prove that this sum is 0 requires algebra.

$$\omega \neq 1, \text{ and } \omega^{10} = 1$$

$$\Rightarrow 1 - \omega^{10} = 0$$

$$\Rightarrow (1 - \omega)(1 + \omega + \omega^2 + \omega^3 + \omega^4 + \dots + \omega^9) = 0 \quad \text{factorising}$$

$$\Rightarrow 1 + \omega + \omega^2 + \omega^3 + \omega^4 + \dots + \omega^9 = 0, \quad \text{since } 1 - \omega \neq 0$$

$$\Leftrightarrow \text{the sum of the complex } 10^{\text{th}} \text{ roots of 1 is 0.}$$

This can be generalized to show that the sum of the n^{th} roots of 1 is 0, for any n .

1st order differential equations

Justification of the Integrating Factor method.

$$\frac{dy}{dx} + Py = Q \quad \text{where } P \text{ and } Q \text{ are functions of } x \text{ only.}$$

We are looking for an Integrating Factor, R (a function of x), so that multiplication by R of the L.H.S. of the differential equation gives an exact derivative.

Multiplying the L.H.S. by R gives

$$R \frac{dy}{dx} + RPy$$

If this is to be an **exact** derivative we can see, by looking at the first term, that we should try

$$\frac{d}{dx}(Ry) = R \frac{dy}{dx} + y \frac{dR}{dx} = R \frac{dy}{dx} + RPy$$

$$\Rightarrow y \frac{dR}{dx} = RPy$$

$$\Rightarrow \int \frac{1}{R} dR = \int P dx$$

$$\Rightarrow \ln R = \int P dx$$

$$\Rightarrow R = e^{\int P dx}$$

Thus $e^{\int P dx}$ is the required I.F., Integrating Factor.

Linear 2nd order differential equations

Justification of the A.E. – C.F. technique for unequal roots

$$\frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = 0$$

without loss of generality we can take the coefficient of $\frac{d^2y}{dx^2}$ as 1.

Let the roots of the A.E. be α and β ($\alpha \neq \beta$), then the A.E. can be written as

$$(m - \alpha)(m - \beta) = 0 \Leftrightarrow m^2 - (\alpha + \beta)m + \alpha\beta = 0$$

So the differential equation can be written

$$\frac{d^2y}{dx^2} - (\alpha + \beta) \frac{dy}{dx} + \alpha\beta y = 0 \quad \text{I}$$

We can 'sort of factorise' this to give

$$\left(\frac{d}{dx} - \alpha\right)\left(\frac{dy}{dx} - \beta y\right) = 0 \quad \text{II} \quad \text{'multiply' out to check}$$

Now put $\left(\frac{dy}{dx} - \beta y\right) = z$, in II, and we get $\frac{dz}{dx} - \alpha z = 0$

$$\Rightarrow \int \frac{1}{z} dz = \int \alpha dx \Rightarrow z = A e^{\alpha x}$$

$$\text{But } \left(\frac{dy}{dx} - \beta y\right) = z \Rightarrow \frac{dy}{dx} - \beta y = A e^{\alpha x}$$

The Integrating Factor is $e^{-\beta x}$

$$\Rightarrow e^{-\beta x} \frac{dy}{dx} - \beta e^{-\beta x} y = A e^{\alpha x} e^{-\beta x} \Rightarrow \frac{d(e^{-\beta x} y)}{dx} = A e^{(\alpha - \beta)x}$$

$$\Rightarrow e^{-\beta x} y = \frac{A}{(\alpha - \beta)} e^{(\alpha - \beta)x} + B$$

$$\Rightarrow y = A' e^{\alpha x} + B e^{\beta x}$$

which is the C.F., for **unequal** roots of the A.E.

Justification of the A.E. – C.F. technique for equal roots

$$\frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = 0$$

without loss of generality we can take the coefficient of $\frac{d^2y}{dx^2}$ as 1.

Let the roots of the A.E. be α and α , (**equal** roots) then the A.E. can be written as

$$(m - \alpha)(m - \alpha) = 0 \Leftrightarrow m^2 - 2\alpha m + \alpha^2 = 0$$

So the differential equation can be written

$$\frac{d^2y}{dx^2} - 2\alpha \frac{dy}{dx} + \alpha^2 y = 0 \quad \text{I}$$

We can 'sort of factorise' this to give

$$\left(\frac{d}{dx} - \alpha\right)\left(\frac{dy}{dx} - \alpha y\right) = 0 \quad \text{II} \quad \text{'multiply' out to check}$$

Now put $\left(\frac{dy}{dx} - \alpha y\right) = z$, in II, and we get $\frac{dz}{dx} - \alpha z = 0$

$$\Rightarrow \int \frac{1}{z} dz = \int \alpha dx \Rightarrow z = A e^{\alpha x}$$

$$\text{But } \left(\frac{dy}{dx} - \alpha y\right) = z \Rightarrow \frac{dy}{dx} - \alpha y = A e^{\alpha x}$$

The Integrating Factor is $e^{-\alpha x}$

$$\Rightarrow e^{-\alpha x} \frac{dy}{dx} - \alpha e^{-\alpha x} y = A e^{\alpha x} e^{-\alpha x} \Rightarrow \frac{d(e^{-\alpha x} y)}{dx} = A$$

$$\Rightarrow e^{-\alpha x} y = Ax + B$$

$$\Rightarrow y = (Ax + B)e^{\alpha x}$$

which is the C.F., for **equal** roots of the A.E.

Justification of the A.E. – C.F. technique for complex roots

Suppose that α and β are complex roots of the A.E., then they must occur as a conjugate pair (see FP1),

$$\Rightarrow \alpha = a + ib \text{ and } \beta = a - ib$$

$$\Rightarrow \text{C.F. is } y = A e^{(a+ib)x} + B e^{(a-ib)x} \text{ assuming that calculus works for complex nos. which it does}$$

$$\Rightarrow y = e^{ax} (A e^{ibx} + B e^{-ibx}) = e^{ax} (A(\cos x + i \sin x) + B(\cos x - i \sin x))$$

$$\Rightarrow \text{C.F. is } y = e^{ax} (C \cos x + D \sin x), \quad \text{where } C \text{ and } D \text{ are arbitrary constants.}$$

We now have the rules for finding the C.F. as before

$$a \frac{d^2y}{dx^2} + b \frac{dy}{dx} + cy = 0 \quad \text{where } a, b \text{ and } c \text{ are constants.}$$

First write down the Auxiliary Equation, A.E

$$\text{A.E. } am^2 + bm + c = 0$$

and solve to find the roots $m = \alpha$ or β

- If α and β are both real numbers, and if $\alpha \neq \beta$ then the Complimentary Function, C.F., is
- $y = A e^{\alpha x} + B e^{\beta x}$, where A and B are arbitrary constants of integration
- If α and β are both real numbers, and if $\alpha = \beta$ then the Complimentary Function, C.F., is
- $y = (A + Bx) e^{\alpha x}$, where A and B are arbitrary constants of integration
- If α and β are both complex numbers, and if $\alpha = a + ib$, $\beta = a - ib$ then the Complimentary Function, C.F.,
- $y = e^{ax} (A \sin bx + B \cos bx)$, where A and B are arbitrary constants of integration

Justification that G.S. = C.F. + P.I.

Consider the differential equation $ay'' + by' + cy = f(x)$

Suppose that u (a function of x) is any member of the Complimentary Function, and that v (a function of x) is a Particular Integral of the above D.E.

$$\Rightarrow au'' + bu' + cu = 0$$

$$\text{and } av'' + bv' + cv = f(x)$$

Let $w = u + v$

$$\begin{aligned} \text{then } aw'' + bw' + cw &= a(u + v)'' + b(u + v)' + c(u + v) \\ &= (au'' + bu' + cu) + (av'' + bv' + cv) = 0 + f(x) = f(x) \end{aligned}$$

$$\Rightarrow w \text{ is a solution of } ay'' + by' + cy = f(x)$$

$$\Rightarrow \text{all possible solutions } y = u + v \text{ are part of the General Solution.} \quad \mathbf{I}$$

We now have to show that **any** member of the G.S. can be written in the form $u + v$, where u is some member of the C.F., and v is the P.I. used above.

Let z be **any** member of the G.S, then $az'' + bz' + cz = f(x)$.

Consider $z - v$

$$a(z - v)'' + b(z - v)' + c(z - v) = (az'' + bz' + cz) - (av'' + bv' + cv) = f(x) - f(x) = 0$$

$$\Rightarrow (z - v) \text{ is some member of the C.F. - call it } u$$

$$\Rightarrow z - v = u \Rightarrow z = u + v$$

thus **any** member, z , of the G.S. can be written in the form $u + v$, where u is some member of the C.F., and v is the P.I. used above. **II**

I and **II** \Rightarrow the Complementary Function + a Particular Integral forms the complete General Solution.

Maclaurin's Series

Proof of Maclaurin's series

To express any function as a power series in x

$$\text{Let } f(x) = a + bx + cx^2 + dx^3 + ex^4 + fx^5 + \dots \quad \mathbf{I}$$

$$\text{put } x = 0 \Rightarrow f(0) = a$$

$$\frac{d}{dx} \Rightarrow f'(x) = b + 2cx + 3dx^2 + 4ex^3 + 5fx^4 + \dots$$

$$\text{put } x = 0 \Rightarrow f'(0) = b$$

$$\frac{d}{dx} \Rightarrow f''(x) = 2 \times 1c + 3 \times 2dx + 4 \times 3ex^2 + 5 \times 4fx^3 + \dots$$

$$\text{put } x = 0 \Rightarrow f''(0) = 2 \times 1c \Rightarrow c = \frac{1}{2!} f''(0)$$

$$\frac{d}{dx} \Rightarrow f'''(x) = 3 \times 2 \times 1d + 4 \times 3 \times 2ex + 5 \times 4 \times 3fx^2 + \dots$$

$$\text{put } x = 0 \Rightarrow f'''(0) = 3 \times 2 \times 1d \Rightarrow d = \frac{1}{3!} f'''(0)$$

continuing in this way we see that the coefficient of x^n in **I** is $\frac{1}{n!} f^n(0)$

$$\Rightarrow f(x) = f(0) + xf'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots + \frac{x^n}{n!} f^n(0) + \dots$$

The range of x for which this series converges depends on $f(x)$, and is beyond the scope of this course.

Proof of Taylor's series

If we put $f(x) = g(x + a)$ then

$$f(0) = g(a), f'(0) = g'(a), f''(0) = g''(a), \dots, f^n(0) = g^n(a), \dots$$

and Maclaurin's series becomes

$$g(x + a) = g(a) + xg'(a) + \frac{x^2}{2!} g''(a) + \frac{x^3}{3!} g'''(a) + \dots + \frac{x^n}{n!} g^n(a) + \dots$$

which is Taylor's series for $g(x + a)$ as a power series in x

Replace x by $(x - a)$ and we get

$$g(x) = g(a) + (x - a)g'(a) + \frac{(x - a)^2}{2!} g''(a) + \frac{(x - a)^3}{3!} g'''(a) + \dots + \frac{(x - a)^n}{n!} g^n(a) + \dots$$

which is Taylor's series for $g(x)$ as a power series in $(x - a)$

Index

- complex numbers, 7
 - applications of De Moivre's theorem, 8
 - argument, 7
 - De Moivre's theorem, 8
 - Euler's relation, 7
 - loci, 10
 - loci and geometry, 13
 - modulus, 7
 - n th roots, 9
 - n^{th} roots of 1, 32
 - roots of polynomial equations, 10
 - transformations, 12
- first order differential equations, 14
 - exact equations, 14
 - families of curves, 14
 - integrating factors, 15
 - integrating factors – proof of method, 34
 - separating the variables, 14
 - using substitutions, 15
- inequalities, 3
 - algebraic solutions, 3
 - graphical solutions, 4
- Maclaurin and Taylor series, 22
 - expanding compound functions, 25
 - proofs, 39
 - standard series, 23
 - worked examples, 23
- method of differences, 5
- polar coordinates, 26
 - area, 29
 - cardioid, 26
 - circle, 29
 - lemniscate, 28
 - polar and cartesian, 26
 - $r = a \cos n\theta$, 28
 - spiral, 28
 - tangent, 30
- second order differential equations, 17
 - auxiliary equation, 17, 37
 - complimentary function, 17, 37
 - general solution, 18
 - justification of technique, 35
 - linear with constant coefficients, 17
 - particular integral, 18
 - using substitutions, 21