



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2002

XVIII

1583

Time allowed
182 Minutes

Score

/152

Percentage

%

CHEMISTRY

**OCR
AS & A LEVEL**

Mark Scheme

**Module 4: Core organic
chemistry**

www.exampaperspractice.co.uk



1. Propene ✓

ALLOW prop-1-ene ✓
DO NOT ALLOW prop-2-ene

[1]

2. (i) $\text{—CH}_2\text{CHCl—} + 2\frac{1}{2}\text{O}_2 \rightarrow 2\text{CO}_2 + \text{H}_2\text{O} + \text{HCl}$ ✓

1

(ii) Alkali **OR** base **OR** carbonate ✓

ALLOW correct formula of or named carbonate
OR alkali OR base
Correct name and wrong formula does not score

1

[2]

3. Any two marks from the following:

Develop photodegradable polymers ✓

Develop biodegradable polymers

OR develop compostable polymers ✓

Develop techniques for cracking polymers

OR develop use as a chemical feedstock ✓

Develop ways of making polymers from plant-based substances

OR reduce the need to use finite raw materials such as crude oil ✓


Designing processes with high atom economy

OR reduce waste products during manufacture ✓

Develop ways of sorting **AND** recycling polymers ✓

[2]

4.



one mark for each correct structure ✓ ✓ ✓ ✓

ALLOW skeletal formula **OR** displayed formulae

IGNORE molecular formulae

IF two answers given e.g. name and structure then both must be correct to be given a mark

ALLOW methylpropane **OR** $(\text{CH}_3)_3\text{CH}$ ✓

ALLOW 1, 2-dibromo-methylpropane **OR**

$\text{CH}_2\text{BrCBr}(\text{CH}_3)_2$ ✓


ALLOW 1-bromo-methylpropane **OR** $\text{CH}_2\text{BrCH}(\text{CH}_3)_2$ ✓

ALLOW 2-bromo-methylpropane **OR** $\text{CH}_3\text{CBr}(\text{CH}_3)_2$ ✓

ALLOW ecf if wrong carbon skeleton is used in all of the structures mark first structure wrong and then apply ecf for the rest

[4]

5. curly arrow from double bond to $\text{Br}^{\delta+}$ and curly arrow from $\text{Br}-\text{Br}$ bond pair to $\text{Br}^{\delta-}$ in 1st step ✓
 curly arrow in 2nd step from bromide ion ✓
 correct dipole shown on Br_2 ✓
 correct carbocation shown ✓



Curly arrow must start from the double bond and not a carbon atom, other curly arrow must start from $\text{Br}-\text{Br}$ bond

ALLOW curly arrow from any part of bromide ion

The bromide ion does not need to show a lone pair


Dipole must be partial charge and not full charge

Carbocation needs a full charge and not a partial charge (charges do not need to be surrounded by a circle)

ALLOW carbocation on carbon 1 where electrophile attacks carbon 2 i.e. $^+\text{CH}_2\text{CBr}(\text{CH}_3)_2$

[4]

6. (i)



Free bonds at bond ends must be present

ALLOW minor slip e.g. missing one hydrogen and left as a stick


ALLOW more than two repeat units but must be a whole number of repeat units

IGNORE brackets, use of numbers and n in the drawn structure

1



(ii)



✓
ALLOW skeletal formula
ALLOW CH₂CHF

1

[2]

7. Any two from:

separation into types and recycling **OR** sort plastics, melt and remould ✓

combustion for energy generation ✓

used for cracking **OR** feedstock for plastics or chemicals ✓

IGNORE biodegradable

used as a fuel is insufficient

releases energy is insufficient

ALLOW burning plastics to release energy

ALLOW organic feedstock / raw materials to make organic compounds

[2]

8. 1st bullet

product: CH₃CH₂CHBrCH₂Br (1)

equation: CH₃CH₂CH=CH₂ + Br₂ → CH₃CH₂CHBrCH₂Br (1)

products: CH₃CH₂CHBrCH₃ **and** CH₃CH₂CH₂CH₂Br (1)

(or statement that 2-bromo- is formed)

equation: CH₃CH=CHCH₃ + HBr → CH₃CH₂CHBrCH₃ (1)

(i.e. for one product)


products: CH₃CH₂CHOHCH₃ **and** CH₃CH₂CH₂CH₂OH (1)

(or statement that 2-ol is formed)

equation: CH₃CH=CHCH₃ + H₂O → CH₃CH₂CHOHCH₃ (1)

(i.e. for one product)

6

2nd bullet


1 mark for skeleton with two repeat units (1)

1 mark for correct groups on side chains (1)

2

3rd bullet

two (1) (1) from

energy from incineration

development of biodegradable polymers


cracking of waste polymers

2

[10]

9. (a) (i) phosphoric acid/H⁺/sulphuric acid 1
- (ii) lone/electron pair of electrons acceptor 1

(b) (i)


 Step 1 curly arrow from π -bond to H⁺ 1

 Step 2 curly arrow from lone pair on the O^{δ-} to C⁺ 1


 Step 3 curly arrow from O—H bond to O⁺ 1

- (ii) catalyst ... no marks because it is **not** consumed/used up in the reaction/owtte 1

[6]

10. (a) 3-chloro(-2-)methylprop-1-ene/1-chloro(-2-)methylprop-2-ene 1

(b)




Backbone of 4 carbons and a reasonable attempt gets 1 mark.

2

[3]

11. (a)



curly	1
dipoles shown correctly on the Br-Br and curly arrow from the Br-Br bond towards the Br ^δ	1
correct intermediate shown	1
curly arrow from the lone pair or the negative charge on the Br ⁻ to the C ⁺	1

- | | | | | |
|--|-----|------|--|---|
| | (b) | (i) | Hs are diagonal to each other in the <i>trans</i> / difference clearly shown in a diagram | 1 |
| | | (ii) | (the product is saturated hence) there is no restricted rotation/single bonds allow rotation/because C=C prevents rotation | 1 |

[6]

- | | | |
|-----|---------------------|---|
| 12. | H ₂ | 1 |
| | Ni/Pt/Pd (catalyst) | 1 |

[2]



- | | | | |
|-----|------|--|---|
| 13. | (i) | alkene | 1 |
| | | bromine | 1 |
| | | decolourises | 1 |
| | (ii) | 3-methylhex-2-en-1-ol/ 1-hydroxy-3-methylhex-2-ene | 1 |

[4]

14. **margarine**


Ni catalyst 1

hydrogen/ hydrogenated 1

unsaturated vegetable oil/fat 1


poly(propene)

equation



1

two repeat units



1

(Ziegler) catalyst / high temp/heat/use of an initiator

Problems with disposal

non-biodegradable/don't decompose/not broken down by bacteria etc 1

when burnt produces toxic fumes 1

Future methods of disposal

recycling (to produce new polymers) 1

incineration for energy (production) 1

cracking/owtte (to produce useful organic molecules)

use gas scrubbers to reduce toxic fumes

any two

max = 9

QWC

Answer is well organised/structure and using at least three of:

catalyst, hydrogenation, addition polymerisation, Ziegler, incineration, feedstock, recycling, non-biodegradable, initiator, monomer, unsaturated.

in the correct context. 1


[10]

15. (a) (i) C₅H₈ 1

(ii) C₅H₈ 1

(b) (i) Ni/Pt/Pd 1

- (ii) 1 mark for C_5H_{12} 1
 1 mark for correct balancing 1
- (iii)




1

[6]

16. (i) electron/lone pair acceptor 1

(ii)



curly arrow from π -bond to $Br^{\delta+}$

Dipoles on the $Br-Br$ bond
and
 curly arrow from $Br-Br$ bond to $Br^{\delta-}$

Curly arrow from Br^- to C^+


1

1

1

[4]

17. (i) correctly shows three repeat units with 'end bonds' 1
 correctly identifies the repeat unit 1



- (ii) harmful/toxic fumes are produced 1


- (iii) recycle/remove HCl by using gas scrubbers or wtte/crack polymers/used a feedstock/ source of fuel (in an incinerator)/developing biodegradable alternatives. 2

[5]



18. (a) (i) $24.7/12 : 2.1/1 : 73.2/35.5$
 $2.06 : 2.1 : 2.06$ 1
CHCl 1
(ii) $(CHCl = 12 + 1 + 35.5 =) 48.5$ 1
 $48.5 \times 3 = 145.5$ 1

- (b) (i) Any two from 2



- (ii) 1, 2,3-trichloropropene
(trichloropropene scores 1 mark ✓)


3 marking points:

- correct numbers 1, 2,3
- trichloro
- propene/prop-1-ene

any two gets 1 mark

2

- (c) (i) 2




1 mark if backbone contains 4 carbons with 'end-bonds' and a reasonable attempt has been made
e.g used the wrong isomer.... max = 1 mark

- (ii) non-biodegradable 1
toxic fumes evolved when burnt 1
HCl or Cl• or chlorinated organic compounds such as COCl₂ also evolved when burnt 1

[13]



19. (i) decolourises 1
(ii)




- curly arrow from C=C bond to bromine 1
dipoles on Br₂ or curly arrow to show movement of bonded pair of electrons 1
intermediate carbonium ion/carbocation 1
curly arrow from lone pair on the Br⁻ ion to carbonium ion (Br^{δ-} loses 1 mark) 1

[5]

20. (a) (i) reaction 1 1
(ii) reaction 4 1
(iii) reaction 3 1

- (b) (i) lone pair/electron pair donor 1




- Correct dipole 1
Curly arrow from the O in the OH⁻ to C in the CH₂ 1
Curly arrow to show movement of bonded pair in the C-Cl bond 1
Cl⁻ as a product 1



(c) (i) same molecular formula, different structure/arrangement of atoms. (same formula, different structure.) 2


(ii) 2



(d) (i) addition, (not additional) 1

(ii) poly(propene)/ polypropene/ polypro-1-ene, polypropylene 1


(iii) 1



[15]

21. (i) decolourises/not clear/not discolours 1

(ii)



curly arrow from C=C to Br^{δ+} 1

dipole on Br-Br **and** curly arrow showing movement of bonded pair of electrons 1

correct intermediate/carbonium ion/carbocation **and** curly arrow from Br⁻ to C⁺ 1

1, 2-dibromopropane as product 1

[5]

22. $\text{CH}_3\text{CBr}_2\text{CH}_3$ 1
 $\text{CH}_3\text{CHBrCH}_2\text{Br}$ 1
 $\text{CH}_3\text{CH}_2\text{CHBr}_2$ 1

($\text{CH}_3\text{CHBrCH}_2\text{Br}$ has a chiral centre, hence optical isomers of 1, 2-dibromopropane are acceptable but must be drawn with 'wedge-shape' bonds and be non-superimposable mirror images)

[3]

23. (i) *unsaturated* contains a double/multiple/ π bond ✓ 1
hydrocarbon contains hydrogen and carbon **only**. ✓ 1
 (ii) angle **a** $109 - 110^\circ$ ✓ 1
 angle **b** $117 - 120^\circ$ ✓ 1
 (iii)




Diagram to show a minimum of 2 carbons, each with a σ -bond and p-orbitals ✓


Overlap of adjacent p-orbitals (in words or in diagram) ✓ 2

[6]

24. (i) *electrophile*: lone pair (of electrons) acceptor. ✓

1

(ii)



essential mark intermediate carbocation/carbonium ion, accept primary /"triangular"/ ✓

essential mark product ✓

curly arrow from double bond to Br_2 ✓

curly arrow showing movement of electrons in the Br-Br bond **or** the dipole in the Br-Br ✓

curly arrow from lone pair of electrons in Br^- to intermediate ✓
mark any errors first

5 max


[5]

25. (i) Addition (not additional) ✓

1


(ii) ✓

1



(iii) ✓

1




(iv) Poly(but-1-ene) ✓

1


[4]

26. (a) (i) alkene ✓ 1
 alcohol/hydroxy/hydroxyl ✓ 1

- (b) (i) I = alkene & II = alcohol... both are needed ✓ 1
 (ii) decolourised / colourless ✓ 1
 (iii) ✓ 1




- (iv) X as shown below ✓ 1



- (c) (i) Ni/Pt/Rh/Pd ✓ 1
 (ii) compound B is C₁₀H₂₂O ✓ 1
 (iii) C₁₀H₂₀O + H₂ → C₁₀H₂₂O ✓ 1

[9]

27.



1 mark is available if the backbone consists of 4 C atoms and a reasonable attempt has been made ✓✓

[2]

28. (a) (i) Alkene/C=C ✓ 1
 Alcohol/ROH/hydroxy/hydroxyl/OH (not OH⁻ or hydroxide) ✓ 1
 (ii) One of the C in both C=C is joined to two atoms or groups that are the same ✓ 1
- (b) Observation decolourisation (of Br₂) ✓ 1
 Molecular formula C₁₀H₁₈OBr₄ ✓✓ 2
 C₁₀H₁₈OBr₂ gets 1 mark
- (c) reagent CH₃COOH ✓ 1
 catalyst H₂SO₄/H⁺/HCl (aq) or dilute loses the mark ✓ 1
- (d) (i) C₁₀H₁₈O + 2[O] → C₁₀H₁₆O₂ + H₂O ✓✓ 2
 1 mark for H₂O and 1 mark for 2[O]
 (ii) The infra-red spectrum was of compound Y
 because absorption between 1680 – 1750 cm⁻¹ indicates a C=O ✓ 1
 and the absence of a peak between 2500 – 3300 cm⁻¹ shows the absence of the OH hydrogen bonded in a carboxylic acid ✓ 1

[12]