



**EXAM PAPERS PRACTICE**

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2002

**XVIII**

1583

Time allowed  
**66 Minutes**

Score

**/55**

Percentage

**%**

**CHEMISTRY**

**Edexcel  
AS & A LEVEL**

**Mark Scheme**

**Paper 1: Advanced Inorganic  
and Physical Chemistry**

[www.exampaperspractice.co.uk](http://www.exampaperspractice.co.uk)



Question Number	Correct Answer	Reject	Mark
<b>1</b> (a)(i)	Mass of ethanoic acid = $0.04 \times 60.1$ = (2.404 g) (1)  Volume of ethanoic acid = $2.404 \div 1.049$ =  $2.2917 = 2.3 \text{ (cm}^3\text{)}$ (1)  Correct answer with no working (2)  Ignore SF except only one  ALLOW  60.0 for molar mass which gives mass 2.4 and volume 2.288 = $2.3 \text{ cm}^3$ (2)  OR  First step $1.049 \div 60/60.1$ to find number of moles in $1 \text{ cm}^3 = 0.017$ (1)  Then volume = $0.04 \div 0.017$ = $2.3529 \text{ (cm}^3\text{)}$  But note, if whole calculation done on calculator, 60 gives 2.2879 and 61 gives 2.2917. (1)  If units given, they must be correct, but penalise wrong units only once here.		2

Question Number	Correct Answer	Reject	Mark
<b>1</b> (a)(ii)	Syringe  ALLOW Burette  Graduated/adjustable pipette	Gas syringe  Biuret  Just 'pipette'	1



Question Number	Correct Answer	Reject	Mark
<b>1</b> (a) (iii)	To prevent... evaporation/vapour escaping water vapour entering  OR To maintain a closed system  OR To maintain a closed environment  ALLOW  To prevent: air oxidizing the alcohol reaction with air  OR Due to volatility (of chemicals)  IGNORE ...gas escaping ...HCl escaping		1



Question Number	Correct Answer	Reject	Mark
<b>1</b> (a) (iv)	<p>First and second mark</p> <p>Phenolphthalein (1)</p> <p>From colourless to (pale) pink/red (1)</p> <p>ALLOW Other indicators with <math>pK_{in}</math> in range 7.5 – 10</p> <p>Some examples are:</p> <p>Thymol blue ((base)) (yellow to blue)</p> <p>Phenol red (yellow to red)</p> <p>Thymolphthalein (colourless to blue)</p> <p>Second mark depends on correct indicator except bromothymol blue, which is incorrect but very close to range so allow colour yellow to blue.</p> <p>Third mark Sodium ethanoate is (slightly) alkaline</p> <p>OR Ethanoic acid is a weak acid</p> <p>OR Phenolphthalein pH range coincides with vertical section of the pH/titration curve</p> <p>OR Titration of weak acid with strong base</p> <p>OR Neutralisation/equivalence point is at 8-10/ any number between 8 and 10.</p> <p>OR <math>pK_{in} \pm 1</math> lies within vertical region (1)</p> <p>Third mark is independent</p>	<p>Litmus/universal indicator</p> <p>Pink to colourless</p> <p>Thymol blue (acid)</p> <p>Phenyl red Methyl red</p>	3



Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (i)	$\text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{OH} \rightleftharpoons \text{CH}_3\text{COOCH}_2\text{CH}_3 + \text{H}_2\text{O}$  ALLOW  Single arrow  -CO <sub>2</sub> H  -C <sub>2</sub> H <sub>5</sub>  Displayed formulae  IGNORE state symbols even if incorrect		1

Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (ii)	Volume of alkali reacting with ethanoic acid = $77.1 - 11.7 = 65.4 \text{ cm}^3$ (1)  Moles of ethanoic acid = $\frac{65.4 \times 0.200}{1000}$ = $0.01308 / 1.308 \times 10^{-2} \text{ (mol)}$ (1)  Correct answer no working (2)  Ignore SF except 1  Allow internal TE for use of  Moles of ethanoic acid = $\frac{77.1 \times 0.200}{1000}$ = $0.01542 / 1.542 \times 10^{-2} \text{ (mol)}$ max(1)		2

Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (iii)	Number of moles of ethanol =  $0.01308 / 1.308 \times 10^{-2} \text{ (mol)}$  TE same as (ii)		1



Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (iv)	Number of moles of ethyl ethanoate $= 0.0400 - 0.01308 = 0.02692$ (mol)  Allow TE from (ii)/(iii) for example  0.01542 gives 0.02458		1

Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (v)	$K_c = \frac{[\text{CH}_3\text{CO}_2\text{CH}_2\text{CH}_3][\text{H}_2\text{O}]}{[\text{CH}_3\text{CO}_2\text{H}][\text{CH}_3\text{CH}_2\text{OH}]}$ <p style="text-align: right;">(1)</p> $= \frac{0.02692 \times 0.02692}{0.01308 \times 0.01308}$ $= 4.23579 = 4.24$ <p style="text-align: right;">(1)</p> Ignore SF except one  Allow TE from (ii), (iii) and (iv) for example  0.01542 etc gives 2.54  No TE for incorrect expression of $K_c$		2

Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (vi)	The units cancel  OR  There are the same numbers of moles of reactants and products		1

Question Number	Correct Answer	Reject	Mark
<b>1</b> (b) (vii)	(Concentrated) hydrochloric acid contains water		1



Question Number	Correct Answer	Reject	Mark
<b>1</b> (c) (i)	First test tube esterification  OR  addition/elimination  ALLOW Condensation (1)  Second test tube (acid) hydrolysis (1)  Two fully correct answers in wrong order (1) ma	Alkaline hydrolysis followed by acidification	2

Question Number	Correct Answer	Reject	Mark
<b>1</b> (c) (ii)	The values are the same within experimental error  OR  The values are concordant  ALLOW  The values are similar (1)  The equilibrium can be approached from either direction  OR  The reaction is reversible  OR  Any comment relating equilibrium to reversibility  IGNORE Dynamic equilibrium  OR  Rate of reverse reaction = rate of forward reaction (1)	Just...the same	2

Question Number	Correct Answer	Reject	Mark
<b>1</b> (c) (iii)	(Acid) catalyst (makes it faster)  OR Provides H <sup>+</sup> (as a catalyst)  OR Protonates...  OR Protonating agent...  OR Donates protons  OR Increases H <sup>+</sup> concentration	Initiates  Reacts with...  Protates	1



Question Number	Acceptable Answers	Reject	Mark
* <b>2(a)</b>	(A green solution) forms a yellow / orange / brown (solution) ALLOW reddish-brown (1)  A grey / black precipitate ALLOW silver ppt ALLOW solid / crystals for precipitate (1)	Red 'Green(ish)' with any other colour  Silver mirror silver compound	2

Question Number	Acceptable Answers	Reject	Mark
<b>2(b) (i)</b>	0.05(00) (mol dm <sup>-3</sup> )		1

Question Number	Acceptable Answers	Reject	Mark
<b>2(b) (ii)</b>	Amount of silver ion in 10 cm <sup>3</sup> = amount of thiocyanate = $\frac{5.6 \times 0.0200}{1000} = 0.000112/1.12 \times 10^{-4}$ (mol) (1)  So concentration of silver ion = $0.000112 \times \frac{1000}{10} = 0.0112/1.12 \times 10^{-2}$ (mol dm <sup>-3</sup> ) (1)		2

Question Number	Acceptable Answers	Reject	Mark
<b>2(b) (iii)</b>	0.0112/1.12 x 10 <sup>-2</sup> (mol dm <sup>-3</sup> )  Accept TE = answer to (ii)		1

Question Number	Acceptable Answers	Reject	Mark
<b>2(b) (iv)</b>	0.0500 – 0.0112 = 0.0388/3.88 x 10 <sup>-2</sup> (mol dm <sup>-3</sup> )  Accept TE = 0.05 - answer to (iii)  Accept answer to (i) – answer to (iii)		1



Question Number	Acceptable Answers	Reject	Mark
<b>2(b)(v)</b>	$K_c = \frac{[\text{Fe}^{3+}(\text{aq})]}{[\text{Fe}^{2+}(\text{aq})][\text{Ag}^+(\text{aq})]}$ <p>ALLOW <math display="block">K_c = \frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}][\text{Ag}^+]} \quad (1)</math></p> $= \frac{0.0388}{0.0112^2}$ $= 309.311 = 309 \text{ dm}^3 \text{ mol}^{-1}$ <p>Value (1)</p> <p>Unit (any order) (1)</p> <p>Three SF (1)</p> <p>Accept TE from (iii) and (iv): ( use of 0.1 from (i) gives <math>708 \text{ dm}^3 \text{ mol}^{-1}</math>)</p> <p>If [Ag] is included in the numerator and taken as <math>=[\text{Fe}^{3+}(\text{aq})]</math>, then allow unit and SF marks ONLY, but must either state 'no units' or show working</p>	[Ag] in numerator	4

Question Number	Acceptable Answers	Reject	Mark
<b>2(c)(i)</b>	$\Delta S^\ominus_{\text{total}} = 8.31 \times \ln 309$ $= +47.6(4) / +47.6(5) \text{ J mol}^{-1} \text{ K}^{-1}$ <p>OR</p> $= 8.31 \times \ln 309.311 = +47.6(5) \text{ J mol}^{-1} \text{ K}^{-1}$ <p>Accept TE : <math>8.31 \times \ln(\text{answer from b(v)})</math></p> <p>Value (1)</p> <p>Sign <u>and</u> Unit (any order) (1)</p> <p>IGNORE sf except 1</p>		2

Question Number	Acceptable Answers	Reject	Mark
<b>2(c)(ii)</b>	<p>First Mark: One of the products is a solid</p> <p>OR</p> <p>Two moles going to two moles but one of them is a solid</p> <p>OR</p> <p>Two moles of solution react to form one mole of solution / liquid and one mole of solid (1)</p> <p>Second Mark (Hence) RHS more ordered / LHS less ordered (1)</p>		2

Question Number	Acceptable Answers	Reject	Mark
<b>2(c)(iii)</b>	$\Delta S^{\circ}_{\text{surroundings}} = \Delta S^{\circ}_{\text{total}} - \Delta S^{\circ}_{\text{system}}$ $= +47.6 - (-208.3) = (+)255.9 \text{ (J mol}^{-1} \text{K}^{-1}\text{)}$ <p>Accept TE on c(i)</p> <p>IGNORE sf except 1</p>		1

Question Number	Acceptable Answers	Reject	Mark
<b>2(c)(iv)</b>	<p>Because <math>\Delta S^{\circ}_{\text{surroundings}} = \frac{-\Delta H^{\circ}}{T}</math> (1)</p> <p><math>\Delta H = -298 \times 255.9 = -76258 \text{ (J mol}^{-1}\text{)}</math>  <math>= -76.258 \text{ (kJ mol}^{-1}\text{)}</math> (1)</p> <p>Units if given must be correct            Correct answer with or without working scores 2 marks</p> <p>IGNORE SF except 1</p> <p>As T increases <math>\Delta S^{\circ}_{\text{surroundings}}</math> becomes less positive / decreases            therefore  <math>\Delta S_{\text{total}}</math> becomes less positive / decreases            ALLOW more negative for less positive (1)</p>	$\Delta S^{\circ}_{\text{total}} = \frac{-\Delta H^{\circ}}{T}$	3



Question Number	Acceptable Answers	Reject	Mark
<b>2*</b> (d)	No change in the titre ALLOW No significant change Stand alone mark (1)  (though silver solid was removed the equilibrium constant remains the same so) the equilibrium concentration(s) would remain the same (1)  Second mark dependent on first IGNORE references to temperature		2



Question Number	Acceptable Answers	Reject	Mark
3 (a)(i)	$(K_p =) \frac{p_{\text{CH}_3\text{CO}_2\text{H}}}{p_{\text{CH}_3\text{OH}} (x) p_{\text{CO}}}$  Partial pressure symbol can be shown in various ways, eg pp, $p_{\text{CO}}$ , (CO)p, etc  <i>ALLOW</i> p in upper or lower case, round brackets <i>IGNORE</i> units	[ ] State symbols given as (l)  + in botto line	1

Question Number	Acceptable Answers	Reject	Mark
3 (a)(ii)	$P_{\text{CH}_3\text{OH}} = 4.9 \text{ (atm) (1)}$ $P_{\text{CO}} = 4.9 \text{ (atm) (1)}$  1 mark for recognition that pressures are equal  <i>IGNORE</i> units		2

Question Number	Acceptable Answers	Reject	Mark
3 (a)(iii)	$K_p = ((22.2)/(4.9)^2)$ $= 0.925 \text{ (1)}$  $\text{atm}^{-1} \text{ (1)}$ stand alone mark but must match expression used in (a)(iii)  OR  $9.25 \times 10^4 \text{ Pa}^{-1} / 92.5 \text{ kPa}^{-1} \text{ (2)}$  <i>ALLOW</i> TE from (a)(i) if inverted and/or (a)(ii)	Answers to other than 3 significant figures	2



Question Number	Acceptable Answers	Reject	Mark
3 (b)(i)	$\text{CH}_3\text{OH}$ : 3.2 $\text{CO}$ : 3.2 (1) for both values  $\text{CH}_3\text{CO}_2\text{H}$ : 46.8 (1)  <i>ALLOW</i> TE for moles of ethanoic acid based on numbers of methanol and carbon monoxide used, as long as moles of methanol and carbon monoxide are equal and moles ethanoic acid + moles methanol = 50		2

Question Number	Acceptable Answers	Reject	Mark
3 (b)(ii)	$\left(\frac{46.8 \times 32}{53.2}\right) = 28.2 / 28.1504$ (atm)  <i>IGNORE</i> sf except 1  Value = 28.16 if mol fraction rounded  <i>ALLOW</i> TE from (b)(i)	28.1  $\frac{46.8 \times 32}{50} =$ 29.95 (atm)	1

Question Number	Acceptable Answers	Reject	Mark
3 (b)(iii)	exothermic as yield / pp of ethanoic acid / conversion of reactants/ $K_p$ is higher at lower temperature / as equilibrium moves (right) at lower temperature  <i>ALLOW</i> if partial pressure of ethanoic acid < 22.2 atm in (b)(ii), endothermic as yield / pp of ethanoic acid / conversion of reactants/ $K_p$ is lower at lower temperature		1

Question Number	Acceptable Answers	Reject	Mark
3 (c)(i)	No effect and other concentrations change to keep $K_p$ constant / $K_p$ is only affected by temperature/ as equilibrium moves (right) to keep $K_p$ constant / change in pressure does not change $K_p$	As $K_p$ is a constant	1

Question Number	Acceptable Answers	Reject	Mark
3 (c)(ii)	Yield increased to restore fraction / quotient / partial pressure ratio back to $K_p$  <i>ALLOW</i> (equilibrium moves) to use up the methanol / answers based on entropy or Le Chatelier  Correct prediction in (c)(i) and (c)(ii) with inadequate explanations scores 1 mark in (c)(ii)	Just 'equilibrium moves to the right'	1

Question Number	Acceptable Answers	Reject	Mark
3 (d)	Mark independently  Reaction can occur at lower temperature / has lower activation energy / requires less energy (1) less fuel needed / fewer emissions (from fuels) / fewer raw materials needed / less natural resources used (1)  OR  Enables use of an alternative process with higher atom economy (1) fewer raw materials needed / less natural resources used (1)	Answer based on car exhaust emissions	2