



**EXAM PAPERS PRACTICE**

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2002

**XVIII**

1583

Time allowed

Score

Percentage

/

%

**Maths**

**AQA  
AS & A LEVEL**

**Mark Scheme**

**3.9 H: Integration**

[www.exampaperspractice.co.uk](http://www.exampaperspractice.co.uk)



3(a)(i)	$f' = \frac{dy}{dx} = 4x^3 + 2$	B1	1	
(ii)	$\int \frac{2x^3 + 1}{x^4 + 2x} dx$ $= \frac{1}{2} \ln(x^4 + 2x) (+c)$	M1 A1	2	For $k \ln(x^4 + 2x)$ By substitution $k \ln u$ M1 correct A1
(b)(i)	$u = 2x + 1$ $du = 2 dx$ $\int x\sqrt{2x+1} dx =$ $\int \left(\frac{u-1}{2}\right)\sqrt{u} \frac{du}{2}$ $= \frac{1}{4} \int \left(u^{\frac{3}{2}} - u^{\frac{1}{2}}\right) du$	B1  M1  A1	   3	  Must be in terms of $u$ only incl. $du$  AG
(ii)	$\int_0^4 dx = \int_1^9 du$ $\frac{1}{4} \int u^{\frac{3}{2}} - u^{\frac{1}{2}} = \frac{1}{4} \left[ \frac{u^{\frac{5}{2}}}{\frac{5}{2}} - \frac{u^{\frac{3}{2}}}{\frac{3}{2}} \right]$ $= \frac{1}{4} \left[ \left( \frac{2}{5} (9)^{\frac{5}{2}} - \frac{2}{3} (9)^{\frac{3}{2}} \right) - \left( \frac{2}{5} - \frac{2}{3} \right) \right]$ $= \frac{1}{4} [79.2 + 0.2\bar{6}]$ $= 19.86$ $= 19.9$	B1  M1 A1    A1	     4	Or changing $u$ 's to $x$ 's at end    Sight of any of these 3 lines  AG
<b>Total</b>			<b>10</b>	



(b)	$\int x^{-2} \ln x \, dx$	$u = \ln x$	$dv = x^{-2}$	M1	4	Attempt at integration by parts
		$du = \frac{1}{x}$	$v = -x^{-1}$	A1		
		$\int = -\frac{1}{x} \ln x + \int x^{-2} \, dx$		A1		
		$= -\frac{1}{x} \ln x - \frac{1}{x} (+c)$		A1		

(ii)	$R = \left[ -\frac{1}{x} (\ln x + 1) \right]_1^5$	M1		$R = \left[ \text{Their (b)} \right]_1^5$
	$= -\frac{1}{5} (\ln 5 + 1) + (\ln 1 + 1)$	A1		OE
	$= \frac{1}{5} (4 - \ln 5)$	A1	3	convincing argument; AG

(b)	$\int x(2x+1)^8 \, dx$					
	$u = 2x + 1$			B1		OE
	$du = 2 \, dx$					
	$\int = \int \left( \frac{u-1}{2} \right) u^8 \left( \frac{du}{2} \right)$			M1		all in terms of $u$ . Condone omission of $du$
	$= \frac{1}{4} \int u^9 - u^8 \, du$					
	$= \frac{1}{4} \left[ \frac{u^{10}}{10} - \frac{u^9}{9} \right]$			B1		$p \frac{u^{10}}{10} + q \frac{u^9}{9}$
	$= \frac{(2x+1)^{10}}{40} - \frac{(2x+1)^9}{36} (+c)$			A1	4	OE; CAO SC: correct answer, no working/parts in $x$ (B1)



<p>4(a) <math>\int x \sin x \, dx \quad u = x</math></p> $\frac{dv}{dx} = \sin x$ $\frac{du}{dx} = 1 \quad v = -\cos x$ $\int = -x \cos x - \int -\cos x (dx)$ $= -x \cos x + \sin x (+c)$	<p>M1</p> <p>m1</p> <p>A1</p> <p>A1</p>	<p></p> <p></p> <p></p> <p>4</p>	<p>For differentiating one term and integrating other</p> <p>For correctly substituting their terms into parts formula</p> <p>CSO</p>
<p>(b) <math>u = x^2 + 5</math></p> $du = 2x \, dx$ $\int = \int \frac{1}{2} u^{\frac{1}{2}} (du)$ $= \frac{\frac{2}{3} u^{\frac{3}{2}}}{\frac{2}{2}}$	<p>M1</p> <p>A1</p> <p>A1<sup>✓</sup></p>	<p></p> <p></p> <p></p> <p></p>	<p><math>\int k u^{\frac{1}{2}} (du)</math> condone omission of <math>du</math> but M0 if <math>dx</math></p> <p><math>k = \frac{1}{2}</math> OE</p> <p>Ft <math>\int k u^{\frac{1}{2}} du</math></p>
<p>(c) <math>= \frac{1}{3} \sqrt{(x^2 + 5)^3} (+c)</math></p> $y = x^2 - 9$ $x^2 = y + 9$ $V = \pi \int x^2 \, dy$ $= \pi \int (y + 9) \, dy$ $= (\pi) \left[ \frac{y^2}{2} + 9y \right]_1^2 \text{ or } (\pi) \left[ \frac{(y+9)^2}{2} \right]_1^2$ $= (\pi) \left[ 20 - 9\frac{1}{2} \right]$ $= 10\frac{1}{2}\pi$	<p>A1</p> <p>B1</p> <p>M1</p> <p>m1</p> <p>A1</p>	<p>4</p> <p></p> <p></p> <p></p> <p>4</p>	<p>CSO</p> <p>SC <math>\frac{2}{6} \sqrt{(x^2 + 5)^3}</math> with no working B3</p> <p>Must have <math>\pi</math> and <math>x^2</math>, condone omission of <math>dy</math>, but B0 if <math>dx</math></p> <p><math>\int</math> "their <math>x^2</math>" <math>dy</math> integrated } <math>\pi</math> not necessary Limits 2 and 1 substituted in } correct order including - sign }</p> <p>CSO</p>
<p><b>Total</b></p>		<p><b>12</b></p>	



6(a)	$\int xe^{5x} dx$ $u = x \quad dv = e^{5x}$ $du = 1 \quad v = \frac{1}{5}e^{5x}$ $\int = \frac{1}{5}xe^{5x} - \int \frac{1}{5}e^{5x} dx$ $= \frac{1}{5}xe^{5x} - \frac{1}{25}e^{5x} (+c)$	M1 A1 A1 A1	4	integrate one term, differentiate one term
(b)(i)	$u = x^{\frac{1}{2}}$ $du = \frac{1}{2}x^{-\frac{1}{2}} dx$ $\int = \int \frac{1}{1+u} \times 2 du$	M1 A1	2	correct with no errors; AG
(ii)	$\int_1^9 dx = \int_1^3 \frac{2}{1+u} du$ $= [2 \ln(1+u)]_1^3$ $= 2 \ln 4 - 2 \ln 2$ $(= \ln 4)$	m1 M1 A1	3	correct limits used in correct expression, ignoring $k$ for $k \ln(1+u)$ ISW OE
<b>Total</b>			<b>9</b>	