



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2002

XVIII

1583

Time allowed
73 Minutes

Score

/61

Percentage

%

CHEMISTRY

**OCR
AS & A LEVEL**

Mark Scheme




Module 5: Physical chemistry and transition elements

www.exampaperspractice.co.uk



EXAM PAPERS PRACTICE

Question		Expected answers	Marks	Additional guidance
1	a	Co: $(1s^2 2s^2 2p^6) 3s^2 3p^6 3d^7 4s^2$ ✓ Co ³⁺ : $(1s^2 2s^2 2p^6) 3s^2 3p^6 3d^6$ ✓	2	ALLOW $(1s^2 2s^2 2p^6) 3s^2 3p^6 4s^2 3d^7$ (i.e. 4s before 3d) ALLOW upper case D, etc. and subscripts, e.g. [Ar]4S ₂ 3D ₇ If included, ALLOW 4s ⁰
	b	catalyst OR coloured ✓	1	IGNORE forms different oxidation states
	c	Donates an electron/lone pair to a metal ion OR forms a coordinate bond to a metal ion ✓	1	ALLOW donates an electron pair/lone pair to a metal/transition element ALLOW dative (covalent) bond for coordinate bond
	d	i		
		Co(OH) ₂ ✓ precipitation ✓	2	Mark independently ALLOW Co(OH) ₂ (H ₂ O) ₄ ALLOW precipitate (reaction)
		ii		
		CoCl ₄ ²⁻ ✓ ligand substitution ✓	2	Mark independently ALLOW ligand exchange DO NOT ALLOW just substitution

Question	Expected answers	Marks	Additional guidance
e		4	<p>ANNOTATIONS MUST BE USED CARE: Cl can be on any position, e.g. for B</p>  <p>complex ions in C and D can be other way around In one complex ion, the 2 Cls must be opposite one another In the other complex ion, the 2 Cls must be next to one another CARE: Cl atoms can be on any position, e.g. for C and D</p> 
	<p>Marking sequence See also Appendix 2 for examples</p> <ol style="list-style-type: none"> 1. Mark any correct complex ions first Do not look at these complex ions again 2. Mark with crosses any complex ions with incorrect ligands. This could include Cl in complex A, and NH₃Cl and NH₃⁺Cl⁻, but NOT NH₃----- connectivity on the LEFT only and NOT Cl⁻ and NOT just NH₃⁺ Do not look at these complex ions again 3. In the remaining complex ions, identify errors in ligands (See Appendix 2): e.g. <ul style="list-style-type: none"> • NH₃ ligands bonded to an H on the LEFT only: NH₃----- (<i>connectivity error</i>) • Cl⁻ • NH₃⁺ Mark these complex ions to maximise errors but treat any incorrectly bonded NH₃, Cl⁻ and NH₃⁺ as ECF 		



EXAM PAPERS PRACTICE

Question	Expected answers	Marks	Additional guidance
SEE APPENDIX 2 FOR EXAMPLES			
e ii	<p>143.4 OR 107.9 + 35.5 (g mol⁻¹) used <i>i.e. molar mass AgCl</i> OR amount of AgCl = 0.02(000) mol ✓</p> <p>Ratio ratio complex : Cl⁻ = 1 : 2 OR 0.01 : 0.02 ✓</p> <p>Identification – available from 1 : 2 ratio OR 2Cl⁻ Therefore the complex is B ✓</p>	3	<p>DO NOT ALLOW AgCl₂</p> <p>DO NOT ALLOW $\frac{2.868}{0.01}$ 0.01 linked to AgCl, not complex ALLOW this mark ONLY for evidence of Cl⁻</p> <p>Quality of Written Communication Identification as B is dependent on correct 1 : 2 ratio OR 2Cl⁻ for this mark</p>
Total		15	

For more help please visit our website www.exampaperspractice.co.uk



Question	Answer	Mark	Guidance
2 (a)	Fe: $(1s^2 2s^2 2p^6) 3s^2 3p^6 3d^6 4s^2$ ✓ Fe ²⁺ : $(1s^2 2s^2 2p^6) 3s^2 3p^6 3d^6$ ✓	2	ALLOW 4s before 3d, i.e. $(1s^2 2s^2 2p^6) 3s^2 3p^6 4s^2 3d^6$ ALLOW 4s ⁰ ALLOW subscripts IGNORE 1s ² 2s ² 2p ⁶ is written out a second time
(b)	coloured (compound/complex/precipitate/ions) OR catalyst ✓	1	IGNORE 'variable oxidation states' but ALLOW the idea that Fe ²⁺ can react to form an ion with a different charge/oxidation state. 'ion' is essential: 'atom' or 'metal' is not sufficient IGNORE partially filled d sub-shell/d orbital (question refers to property of Fe ²⁺)
(c)	Fe oxidised from +2 to +3 ✓ Cr reduced from +6 to +3 ✓	2	CHECK and credit oxidation numbers on equation ALLOW Fe ²⁺ oxidised to Fe ³⁺ ALLOW Cr ⁶⁺ reduced to Cr ³⁺ ALLOW + sign after number in oxidation number, <i>ie</i> 2+, etc ALLOW 1 mark only if oxidation numbers given with no identification of which species has been oxidised or reduced, <i>ie</i> Fe goes from +2 to +3 AND Cr goes from +6 to +3 Fe reduced from +2 to +3 AND Cr oxidised from +6 to +3 (<i>oxidation and reduction the wrong way around</i>) DO NOT ALLOW just 'Fe is oxidised and Cr reduced' IGNORE other oxidations numbers (even if wrong) IGNORE any references to electrons







Question	Answer	Mark	Guidance
2 (d)	$(K_{\text{stab}} =) \frac{[\text{Fe}(\text{NH}_3)_6]^{2+}}{[\text{Fe}(\text{H}_2\text{O})_6]^{2+} [\text{NH}_3]^6}$ <p>On top, ONLY $[\text{Fe}(\text{NH}_3)_6]^{2+}$ shown AND on bottom, $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ AND $[\text{NH}_3]^6$ shown ✓</p> <p>correct use of square brackets and double square brackets in expression ✓</p>	2	IGNORE state symbols ALLOW 1 mark if complete expression with correct use of double brackets is shown but upside down
(e) (i)	<p>O_2/oxygen bonds to Fe^{2+}/Fe(II)/Fe ✓</p> <p>When required, O_2 substituted OR O_2 released ✓</p>	2	ANNOTATE WITH TICKS AND CROSSES, etc ALLOW O_2 binds to Fe^{2+} OR O_2 donates electron pair to Fe^{2+} ALLOW O_2 bonds to metal ion/metal DO NOT ALLOW just O_2 bonds to haemoglobin OR O_2 bonds to complex



Question	Answer	Mark	Guidance
2 (e) (ii)	<p>(For complex) with CO, stability constant is greater (than with complex in O₂) OR with CO, stability constant is high ✓</p> <p>(Coordinate) bond with CO is stronger (than O₂) OR bond with CO is strong ✓</p>	2	<p>ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>Comparison of CO and O₂ is NOT required ALLOW stability constant with/of CO is greater IGNORE (complex with) CO is more stable</p> <p>ALLOW bond with CO is less likely to break OR bond with CO more likely to form OR 'CO cannot be removed' OR idea that attachment of CO is irreversible OR CO is a stronger ligand (than O₂) OR CO has greater affinity for ion/metal/haemoglobin (than O₂)</p> <p>IGNORE CO bonds more easily</p>
(f) (i)	Pt ²⁺ /Pt is +2/2+, 2 x Cl ⁻ -2 ✓	1	<p>DO NOT ALLOW response in terms of Cl₂ rather than Cl⁻ DO NOT ALLOW 'charges cancel' without the charges involved being stated</p>



Question	Answer	Mark	Guidance
2 (f) (ii)	<p>✓✓ For each structure</p> <p>Ligand donates an electron pair to metal (ion)/Pt²⁺/Pt OR forms a coordinate bond to the metal (ion)/Pt²⁺/Pt ✓</p>	3	<p>IGNORE any charge, ie Pt²⁺ OR Cl⁻, even if wrong IGNORE any angle, even if wrong ACCEPT bonds to H₃N (does not need to go to 'N')</p> <p>Assume that a solid line is in plane of paper Each structure must contain 2 'out wedges' AND 2 'in wedges' or dotted lines OR 4 solid lines at right angles (all in plane of paper)</p> <p>DO NOT ALLOW any structure that cannot be in one plane DO NOT ALLOW any structure with Cl₂ as a ligand DO NOT apply ECF from one structure to the other</p> <p>ALLOW coordinate bonds shown on diagrams provide that they start from a lone pair</p> <p>ALLOW 'dative covalent bond' or 'dative bond' as alternative for 'coordinate bond'</p> <p>IGNORE <i>cis</i> and <i>trans</i> labels (even if incorrect) IGNORE incorrect connectivity to NH₃, ie ALLOW NH₃—</p>
(iii)	platin binds to DNA (of cancer cells) OR platin stops (cancer) cells dividing/replicating ✓	1	

Question	Answer	Mark	Guidance
2 (g)	<p>1,1-cyclobutanedicarboxylate ion</p>  <p>Correct charge required (could also be 2- outside square brackets)</p> <p>carboplatin (<i>cis</i> isomer shown below)</p> 	2	<p>Must show cyclobutane ring with both COO⁻ groups bonded to same carbon</p> <p>ALLOW COO⁻ OR CO₂⁻ for each carboxylate ion ALLOW structures showing CH₂ or C atoms provided it is clear that C skeleton is shown, Note: H atoms are not required if C atoms shown, <i>ie</i></p>  <p>DO NOT ALLOW circle inside cyclobutane ring</p> <p>Two bonds from Pt to O atoms</p> <p>Any bonds from ligand MUST come from O OR from atom with lone pair</p> <p>IGNORE any charge shown Note: H atoms are not required if C atoms shown, (see ion in 1st structure)</p> <p>ALLOW ECF from 1st structure provided that the attached atoms are capable of forming coordinate bonds (<i>ie</i> they contain a lone pair of electrons) Example if 1st structure is as below, then ALLOW 1 mark ECF</p> 



Question	Answer	Mark	Guidance
3 (a) (i)	<p>amount S₂O₃²⁻ used $= 0.00100 \times \frac{24.6}{1000} = 2.46 \times 10^{-5} \text{ mol } \checkmark$</p> <p>amount O₂ in 25 cm³ sample $= \frac{2.46 \times 10^{-5}}{4} = 6.15 \times 10^{-6} \text{ mol } \checkmark$</p> <p>Concentration of O₂ in sample $= 6.15 \times 10^{-6} \times \frac{1000}{25} = 2.46 \times 10^{-4} \text{ (mol dm}^{-3}\text{)} \checkmark$</p> <p>mass concentration of O₂ in mg dm⁻³ $= 2.46 \times 10^{-4} \times 32 \text{ g} = 7.872 \times 10^{-3} \text{ (g dm}^{-3}\text{)}$ $= 7.872 \text{ (mg dm}^{-3}\text{)} \checkmark$</p>	4	<p>ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>ALLOW 0.0000246 (mol)</p> <p>ECF = $\frac{\text{answer above}}{4}$</p> <p>ALLOW 0.00000615 g</p> <p>ECF answer above $\times \frac{1000}{25}$</p> <p>ALLOW 0.000246 g</p> <p>ECF = answer above $\times 32 \times 1000$</p> <p>ALLOW 7.9 OR 7.87</p> <p>ALLOW 2 SF up to calculator value</p> <p>Must be in mg for mark</p> <p>Note: Candidate may work out steps 3 and 4 in the opposite order, ie</p> <p>mass of O₂ in sample $= 6.15 \times 10^{-6} \times 32 \times 1000 = 1.968 \times 10^{-1} \text{ mg}$</p> <p>mass concentration of O₂ in mg dm⁻³ $= 1.968 \times 10^{-1} \times \frac{1000}{25} = 7.872 \text{ (mg dm}^{-3}\text{)}$</p>
(ii)	<p>Comment 7.872 > 5 so fish can survive \checkmark</p>	1	<p>ECF If final answer > 5 fish can survive If final answer < 5 fish cannot survive</p>
(b) (i)	NO \checkmark	1	ALLOW N ₂ H ₂



Question		er	Mark	Guidance
(b)	(ii)	$2\text{H}_2\text{O} + 2\text{I}^- + 2\text{NO}_2^- \longrightarrow 2\text{NO} + \text{I}_2 + 4\text{OH}^-$ OR $2\text{H}^+ + \text{I}^- + 2\text{NO}_2^- \longrightarrow 2\text{NO} + \text{I}_2 + 2\text{OH}^-$ species ✓ balance ✓	2	IGNORE state symbols ALLOW multiples For species ONLY, IGNORE any extra H_2O or e^- on either side of the equation ALLOW on LHS: $2\text{HI} + 2\text{NO}_2^-$ OR $2\text{I}^- + 2\text{HNO}_2$ ALLOW species and equation involving N_2H_2 : $6\text{H}_2\text{O} + 8\text{I}^- + 2\text{NO}_2^- \longrightarrow \text{N}_2\text{H}_2 + 4\text{I}_2 + 10\text{OH}^-$ OR $6\text{H}^+ + 8\text{I}^- + 2\text{NO}_2^- \longrightarrow \text{N}_2\text{H}_2 + 4\text{I}_2 + 4\text{OH}^-$ species ✓ balance ✓
Total			8	



Question		Answer	Marks	Guidance
4	(a)	$\text{MnO}_2 + 4\text{OH}^- \longrightarrow \text{MnO}_4^{2-} + 2\text{H}_2\text{O} + 2\text{e}^- \checkmark$ $3\text{H}_2\text{O} + \text{ClO}_3^- + 6\text{e}^- \checkmark \longrightarrow 6\text{OH}^- + \text{Cl}^-$	2	ALLOW 'e': i.e. – sign not required
	(b)	<p>Role of CO₂ CO₂ reacts with H₂O forming an acid OR carbonic acid/H₂CO₃ forms OR CO₂ is acidic ✓</p> <p>Equation involving OH⁻ H₂CO₃ + OH⁻ → H₂O + HCO₃⁻ OR H₂CO₃ + 2OH⁻ → 2H₂O + CO₃²⁻ OR CO₂ + OH⁻ → CO₃²⁻ + H⁺ OR CO₂ + OH⁻ → HCO₃⁻ OR CO₂ + 2OH⁻ → CO₃²⁻ + H₂O OR H⁺ + OH⁻ → H₂O ✓</p> <p>Effect on equilibrium with reason equilibrium shifts to right AND to restore OH⁻ ✓</p>	3	<p>ANNOTATIONS MUST BE USED</p> <p>-----</p> <p>ALLOW equation: CO₂ + H₂O → H₂CO₃ OR CO₂ + H₂O → H⁺ + HCO₃⁻ OR CO₂ + H₂O → 2H⁺ + CO₃²⁻</p> <p>ALLOW for 'restores OH⁻' the following: 'makes more OH⁻', 'OH⁻ has been used up' DO NOT ALLOW just 'equilibrium shifts to right'</p>



Question	Answer	Marks	Guidance
(c)	<p>FOLLOW through stages to mark</p> <hr/> <p>Moles in titration</p> $n(\text{KMnO}_4) = 0.0200 \times \frac{26.2}{1000} = 5.24 \times 10^{-4} \text{ mol } \checkmark$ $n(\text{SO}_3^{2-}) = 1.31 \times 10^{-3} \text{ mol } \checkmark$ <p>Scaling</p> $n(\text{SO}_3^{2-}) \text{ in original } 100 \text{ cm}^3$ $= 4 \times 1.31 \times 10^{-3} = 5.24 \times 10^{-3} \text{ mol } \checkmark$ <p>Mass</p> <p>Mass of Na_2SO_3 in sample</p> $= 126.1 \times 5.24 \times 10^{-3} \text{ g} = 0.660764 \text{ g } \checkmark$ <p>Percentage</p> $\% \text{ Na}_2\text{SO}_3 = \frac{0.660764}{0.720} \times 100 = 91.8\% \checkmark$	5	<p>ANNOTATIONS MUST BE USED</p> <p>AT LEAST 3 SF for each step</p> <hr/> <p>ECF 2.5 x answer above</p> <p>ECF 4 x answer above</p> <p>ECF 126.1 x answer above</p> <p>ALLOW 0.661 g up to calculator value</p> <p>ECF $\frac{\text{calculated mass above}}{0.720} \times 100$</p> <p>ALLOW 91.8% (1 DP) up to calculator value of 91.77277778</p> <p>i.e. DO NOT ALLOW 92%</p> <p>COMMON ERRORS:</p> <p>36.8(1)% 4 marks no 2.5 factor</p> <p>22.9(4)% 4 marks no scaling by 4</p> <p>9.18% 3 marks no 2.5 and no x 4</p> <p>Watch for random ECF %s for % from incorrect $M(\text{Na}_2\text{SO}_3)$, e.g. use of $M(\text{SO}_3^{2-}) = 80.1$ giving 58.3%</p>
	<p>ALLOW alternative approach based on theoretical content of Na_2SO_3 for last 2 marks</p> <p>Theoretical amount, in moles, of Na_2SO_3 in sample</p> $n(\text{Na}_2\text{SO}_3) = \frac{0.720}{126.1} = 5.71 \times 10^{-3} \text{ mol } \checkmark$ <p>Percentage</p> $\% \text{ Na}_2\text{SO}_3 = \frac{5.24 \times 10^{-3}}{5.71 \times 10^{-3}} \times 100 = 91.8\% \checkmark$		
	Total	10	



Question	Expected answers	Marks	Additional guidance
5 a	$\text{Fe}_2\text{O}_3 + 6\text{H}^+ \longrightarrow 2\text{Fe}^{3+} + 3\text{H}_2\text{O} \checkmark$	1	ALLOW $\text{Fe}_2\text{O}_3 + 6\text{HCl} \longrightarrow 2\text{FeCl}_3 + 3\text{H}_2\text{O}$ OR $\text{Fe}_2\text{O}_3 + 6\text{HCl} \longrightarrow 2\text{Fe}^{3+} + 6\text{Cl}^- + 3\text{H}_2\text{O}$ ALLOW correct multiples IGNORE state symbols DO NOT ALLOW Fe_2Cl_6 as a product
b	$\text{Sn}^{2+} + 2\text{Fe}^{3+} \longrightarrow \text{Sn}^{4+} + 2\text{Fe}^{2+} \checkmark$ $6\text{Fe}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \longrightarrow$ $6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O} \checkmark$	2	IGNORE state symbols ALLOW overall equations: $\text{SnCl}_2 + 2\text{FeCl}_3 \longrightarrow \text{SnCl}_4 + 2\text{FeCl}_2$ $6\text{FeCl}_2 + \text{K}_2\text{Cr}_2\text{O}_7 + 14\text{HCl} \rightarrow$ $6\text{FeCl}_3 + 2\text{CrCl}_3 + 2\text{KCl} + 7\text{H}_2\text{O}$ ALLOW correct multiples



Question	Expected answers	Marks	Additional guidance
c	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 54.6%, award 5 marks</p> <hr/> <p>Amount Fe²⁺ in 250 cm³ solution – 3 marks amount Cr₂O₇²⁻ used = $0.0200 \times \frac{26.5}{1000}$ = 5.30×10^{-4} (mol) ✓</p> <p>amount Fe²⁺ = $6 \times 5.30 \times 10^{-4}$ = 3.18×10^{-3} mol ✓</p> <p>amount Fe²⁺ in original 250 cm³ = $10 \times 3.18 \times 10^{-3}$ = 3.18×10^{-2} (mol) ✓</p>		<p>ANNOTATIONS MUST BE USED IF there is an alternative answer, 1st check common errors below. Then see if there is any ECF credit possible using working below</p> <hr/> <p>Working must be to at least 3 SF throughout BUT ignore trailing zeroes, <i>i.e.</i> for 0.490 allow 0.49</p> <p>ALLOW ECF from different Fe²⁺ ratio in equation from 8(b) BUT still ALLOW 6 : 1 even from different ratio in equation If no equation use actual 6 : 1 ratio DO NOT AWARD 'ratio mark' at all for use of 1 : 1 ratio – <i>makes problem easier</i></p> <p>ECF 10 × answer above</p>
	<p>% Fe in ore – 2 marks mass of Fe in ore = $55.8 \times 3.18 \times 10^{-2}$ g = 1.77444 g ✓</p>		<p>ECF 55.8 × answer above</p> <p>IF answer above has not been used AND × 55.8, DO NOT ALLOW this mark but do ALLOW final %</p> <p>IF answer above AND 55.8 are BOTH not used, then DO NOT ALLOW ANY further marks</p>
	<p>percentage Fe in ore = $\frac{1.77444}{3.25} \times 100$ = 54.6% ✓</p>	5	<p>ECF $\frac{\text{answer above}}{3.25} \times 100$</p> <p>ALLOW 54.5% (from 1.77 g) AND any answer with > 1 decimal place that rounds back to 54.5 OR 54.6</p>
			<p>COMMON ERRORS</p> <p>5.46 ✓✓✓✓ × 10 omitted 51.5 ✓✓✓✓ titre taken as 25.0 156.2 ✓✓✓✓ × 159.6 instead of 55.8 15.62 ✓✓✓ × 159.6 and × 10 omitted 45.5 ✓✓✓✓ 5 : 1 ratio 1.52 ✓✓✓✓ ÷ 6 instead of × 6</p>



Question	Expected answers	Marks	Additional guidance
d	E^\ominus for MnO_4^- is more positive/greater than Cl_2 OR E^\ominus for $\text{Cr}_2\text{O}_7^{2-}$ is less positive/smaller than Cl_2 ✓ MnO_4^- reacts with Cl^- OR HCl (forming Cl_2 gas) OR $\text{Cr}_2\text{O}_7^{2-}$ does not react with Cl^- ions ✓	2	ORA: E^\ominus for Cl_2 is less positive/smaller than MnO_4^- OR E^\ominus for Cl_2 is more positive/greater than $\text{Cr}_2\text{O}_7^{2-}$
Total		10	