

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you



Time allowed 67 Minutes

Score

/56

Percentage

%

CHEMISTRY

OCR AS & A LEVEL

Topic Questions

Module 6: Organic chemistry and analysis

www.exampaperspractice.co.uk



F324: Rings, Polymers and Analysis 4.1.2 Carbonyl Compounds

1.	•	oxyethanal, HOCH ₂ CHO, is sometimes referred to as the 'first sugar' as it is the lest possible molecule that contains both an aldehyde group and an alcohol group.	
		ochemist investigated some redox reactions of hydroxyethanal and found that eral different products were produced.	
	(a)	The biochemist reacted hydroxyethanal with Tollens' reagent.	
		(i) State what the biochemist would see when hydroxyethanal reacts with Tollens' reagent.	
		(ii) Write the structural formula of the organic product formed when hydroxyethanal reacts with Tollens' reagent.	[1]
			[1]
	(b)	The biochemist also reacted hydroxyethanal with acidified dichromate by heating under reflux.	
		Write an equation for this oxidation.	
		Use [O] to represent the oxidising agent.	
			[2]
	(c)	The biochemist then reduced hydroxyethanal using aqueous NaBH ₄ .	
		(i) Write the structural formula of the organic product.	

[1]



(ii) Outline the mechanism for this reduction.

Use curly arrows and show any relevant dipoles.

[4]

[Total 9 marks]

2. Propanal, CH₃CH₂CHO, can be used in the synthesis of organic compounds.

CH₃CH₂CHO reacts with NaBH₄ in a nucleophilic addition reaction. The nucleophile can be represented as a hydride ion, H⁻. A mechanism for the reaction is shown below.

(i) Add 'curly arrows' to the mechanism to show how the intermediate reacts with the water molecule in **step 2**.

[2]

(ii) Draw the structure of the organic product in the box above.

[1]

(iii) What is meant by the term nucleophile?

[1]



	(iv)		cribe, in wor				pening	to the el	ectror	n pairs	s and	bond	s in	
													⁻ otal 7 m	[3] arks]
3.			vn colourles ut-3-en-1-ol	-		ecular	formula	C ₄ H ₈ O	was t	hougl	ht to b	e one	e of	
	н—	H - -C H	H H 	OH	H C=	=C-	H C -C)Н С—Н	н—	H -C- H	H -C- H	O = -C-	H -C	Ⅎ
		b	utanal			but-3-	en-1-ol			I	butan	one		
	(a)	State	e a simple of butanal or reagent observation	nly;										
			organic pr	oduct										[3]



	(ii)	but-3-en-1-ol only .	
		reagent	
		observation	
		type of reaction	
			[3]
(b)		anal and butanone both react with 2,4-dinitrophenylhydrazine to produce ures containing orange precipitates.	
		ine how the mixtures containing these orange precipitates can be used to nguish between butanal and butanone.	0
			[3]
		דן	otal 9 marks]
But-2	2-enal	I, CH ₃ CH=CHCHO, is a pale yellow, flammable liquid with an irritating od	lour.
(a)	(i)	Describe a simple chemical test that would show that but-2-enal is an aldehyde.	
			[2]

4.



	(11)	Explain why this test gives a different result with aldehydes than it does with ketones.	
			[1]
(b)	But-	2-enal also reacts with sodium borohydride, NaBH ₄ .	
	(i)	Identify the organic compound formed in this reaction.	
			[1]
	(ii)	State the type of chemical reaction occurring.	
			[1]
			۲۰.
(c)	Pred	cautions must be taken to prevent but-2-enal catching fire.	
	Con C ₄ H	struct a balanced equation for the complete combustion of but-2-enal,	
	0411	o♥.	

[1]

[Total 6 marks]



5. (a) The reaction of trichloroethanal with water is a nucleophilic addition reaction. It can be catalysed by small amounts of hydroxide ions, OH⁻.

Complete the diagram below to suggest a mechanism for this reaction. Show all the relevant dipoles and curly arrows.

$$C_{l} = C_{l} = C_{l$$

(b) The recommended adult dose of chloral hydrate as a sedative is 250 mg, three times a day.

Calculate the mass of **trichloroethanal** you would need to react with water to make one week's supply of chloral hydrate for an adult, assuming a 60% yield.

 $M_{\rm r}$: chloral hydrate, 165.5; trichloroethanal, 147.5

mass of trichloroethanal =g

[3]

[5]

[Total 8 marks]



6.	Chloral hydrate is broken down in the body after several hours. One reaction is
	oxidation to trichloroethanoic acid.

Complete the equation for this reaction below.

$$Cl_3CCH(OH)_2 + [O]$$

[Total 1 mark]

7. The reducing agent, NaBH₄, is used widely in organic chemistry. One example is for the reduction of diphenylethanedione, $C_{14}H_{10}O_2$, shown below.

diphenylethanedione

(i) Draw a displayed formula to show the structure of the organic product that would be formed by reducing diphenylethanedione with excess NaBH₄.

[1]

(ii) Complete and balance the equation for this reaction, using [H] to represent the reducing agent.


$$C_{14}H_{10}O_2 \longrightarrow$$

[1]

[Total 2 marks]




8.



Mild oxidation of geraniol gives an aldehyde Y.

(i) Draw the skeletal formula of aldehyde **Y** below.



aldehyde Y

[2]

(ii) Complete the equation for the oxidation of geraniol to aldehyde Y.

$$C_9H_{15}CH_2OH + [O] \rightarrow$$

[2]

[Total 4 marks]

- **9.** Cinnamaldehyde can be reduced using sodium borohydride, NaBH4.
 - (i) State which functional group reacts with the sodium borohydride.

[1]



Complete and balance the equation for this reaction.

(ii)

	$C_6H_5CH=CHCHO$ + [H] \rightarrow
	[1]
	[Total 2 marks]
10.	In this question, one mark is available for spelling, punctuation and grammar.
	Tollens' reagent can be used to identify the aldehyde group in cinnamaldehyde.
	 Describe how you would make Tollens' reagent and carry out this test in the laboratory.
	 Explain what happens to both the Tollens' reagent and the cinnamaldehyde in this reaction. Identify the organic product.
	[7] Quality of Written Communication. [1]

[Total 8 marks]