



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

2002

XVIII

1583

Time allowed
74 Minutes

Score

/62

Percentage

%

CHEMISTRY

**OCR
AS & A LEVEL**


Topic Questions

Module 3: Periodic table and energy


www.exampaperspractice.co.uk

- 1 Butan-2-ol can be prepared using two different methods.

Method 1




$\text{H}_2\text{O(g)}$



butan-2-ol

Method 2



NaOH(aq)

- (a) Comment on the atom economy of each method, giving your reasons.

.....

.....

.....

.....

..... [2]

- (b) State the catalyst required for **Method 1**.

..... [1]

- (c) Average bond enthalpies can be used to calculate enthalpy changes.

- (i) What is meant by the term *average bond enthalpy*?

.....

.....

.....

..... [2]

- (ii) Calculate the enthalpy change of reaction, ΔH_r , for preparing 1 mol of butan-2-ol by **Method 1**.

Average bond enthalpies are given below.

Bond	Average bond enthalpy / kJ mol^{-1}
O–H	464
C–H	413
C–C	347
C–O	358
C=C	612

$$\Delta H_r = \dots\dots\dots \text{kJ mol}^{-1} \quad [3]$$

- (d) A student uses **Method 2** to prepare 3.552 g of butan-2-ol from 2-bromobutane.
The percentage yield of butan-2-ol is 80.0%.

Calculate the mass of 2-bromobutane that the student uses.
Give your answer to **three** significant figures.


$$\text{mass of 2-bromobutane} = \dots\dots\dots \text{g} \quad [3]$$

[Total: 11]

2 A branched-chain alcohol **J** is a liquid and has the molecular formula $C_5H_{12}O$.

(a) A student does an experiment to measure the enthalpy change of combustion, ΔH_c , of alcohol **J**.

(i) The student burns alcohol **J** using the apparatus below.



The student found that combustion of 1.54 g of alcohol **J** changes the temperature of 180 g of water from 22.8°C to 75.3°C .

The specific heat capacity of water is $4.18\text{ J g}^{-1}\text{ K}^{-1}$.

- Calculate the amount, in mol, of alcohol **J** that burns.
- Calculate the enthalpy change of combustion, ΔH_c , of alcohol **J**, in kJ mol^{-1} .

Give your final answer to **three** significant figures.

$\Delta H_c = \dots\dots\dots \text{kJ mol}^{-1}$ [4]

- (ii) The calculated value of ΔH_c from this experiment is different from the value obtained from data books.

Apart from heat loss, suggest **two** reasons for the difference.

Assume that the calculation has been carried out correctly.

.....


 [2]

- (b) The enthalpy change of combustion for alcohol **J** can also be determined indirectly from standard enthalpy changes of formation.

- (i) Write an equation, including state symbols, for the chemical change that represents the standard enthalpy change of formation of the liquid alcohol **J**, $C_5H_{12}O$.

..... [1]

- (ii) The equation for the complete combustion of alcohol **J** is shown below.




Enthalpy changes of formation, ΔH_f , are shown in the table.

Substance	$C_5H_{12}O(l)$	$CO_2(g)$	$H_2O(l)$
$\Delta H_f / kJ mol^{-1}$	-366	-394	-286

Calculate the enthalpy change of combustion, ΔH_c , of alcohol **J** from the information given above.

$$\Delta H_c = \dots\dots\dots kJ mol^{-1} \quad [3]$$

- (c) The branched-chain alcohol **J**, $\text{C}_5\text{H}_{12}\text{O}$, was heated under reflux with excess $\text{H}_2\text{SO}_4/\text{K}_2\text{Cr}_2\text{O}_7$ to form an organic compound **K** with the infrared spectrum below.



- Determine the structures for the branched-chain alcohol **J** and compound **K**. Your answer should explain all your reasoning using the evidence given.
- Write an equation for the reaction of **J** when heated under reflux with excess $\text{H}_2\text{SO}_4/\text{K}_2\text{Cr}_2\text{O}_7$ to form **K**. Use [O] to represent the oxidising agent.



Your answer needs to be clear and well organised using the correct terminology.

[6]

(d) The alcohol **J** is soluble in water.

Explain why alcohol **J** is soluble in water.

Use a labelled diagram to support your answer.

Include relevant dipoles and lone pairs.

.....

.....

..... [1]

[Total: 17]

- 3 Hydrogen iodide, HI, is a colourless gas that can be made from the reaction of hydrogen, H₂, and iodine, I₂.

This reversible reaction is shown in **equilibrium 3.1** below.




The activation energy for the forward reaction is 173 kJ mol⁻¹.

- (a) Complete the enthalpy profile diagram below for the forward reaction in **equilibrium 3.1**.

On your diagram:

- Label the activation energy, E_a
- Label the enthalpy change of reaction, ΔH
- Include the formulae of the reactants and products.



[2]

- (b) Calculate the activation energy, E_a , for the reverse reaction.

$$E_a \text{ (reverse reaction)} = \dots\dots\dots \text{ kJ mol}^{-1} \quad [1]$$

- (c) When the reverse reaction takes place hydrogen iodide, HI, decomposes to form iodine and hydrogen.

Calculate the enthalpy change when 336 dm³ of hydrogen iodide, measured at room temperature and pressure, decomposes.

Include the sign for enthalpy change in your answer.

enthalpy change kJ [2]

- (d) A student mixes hydrogen and iodine at room temperature and pressure and allows the mixture to reach dynamic equilibrium.



- (i) A closed system is required for dynamic equilibrium to be established.

State **one** other feature of this dynamic equilibrium.

.....
 [1]

- (ii) The student heats the equilibrium mixture keeping the volume constant.

Predict how the composition of the equilibrium mixture changes on heating.

Explain your answer.

.....

 [2]

- (iii) Predict and explain what effect, if any, an increase in the pressure would have on the position of the equilibrium.

effect
 explanation
 [1]

- (e) Calculate the bond enthalpy for the H–I bond in **equilibrium 3.1**, given the following information.


Bond	Bond Enthalpy / kJ mol ⁻¹
H–H	436
I–I	151

bond enthalpy kJ mol⁻¹ [2]

[Total: 11]

4 This question is about the determination of enthalpy changes.

- (a) A student carries out an experiment to find the enthalpy change of reaction, ΔH_r , for the reaction below.



In the experiment, 3.18 g of Na_2CO_3 are added to 50.0 g of 2.00 mol dm^{-3} HCl , an excess. The temperature of the reaction mixture increases by 5.5°C .


Calculate ΔH_r , in kJ mol^{-1} .

Give your answer to **three** significant figures.

The specific heat capacity, c , of the reaction mixture is $4.18 \text{ J g}^{-1} \text{ K}^{-1}$.

$\Delta H_r = \dots\dots\dots \text{ kJ mol}^{-1}$ [4]

- (b) The booster rocket of a spacecraft uses a mixture of aluminium and ammonium chlorate(VII), NH_4ClO_4 , as a fuel. The equation and some enthalpy changes are shown below.



Substance	Standard enthalpy change of formation, $\Delta H_f^\circ / \text{kJ mol}^{-1}$
$\text{NH}_4\text{ClO}_4\text{(s)}$	-295
$\text{Al}_2\text{O}_3\text{(s)}$	-1676
$\text{AlCl}_3\text{(s)}$	-704
$\text{H}_2\text{O(g)}$	-242

- (i) What is meant by the term *standard enthalpy change of formation*?

Give the standard conditions.

.....

 [3]

- (ii) Write the equation, including state symbols, for the reaction that represents the standard enthalpy change of formation of $\text{NH}_4\text{ClO}_4\text{(s)}$.


..... [2]

- (iii) Calculate the enthalpy change of formation of NO(g) using the data above.

enthalpy change of formation of NO(g) = kJ mol^{-1} [3]
[Total: 12]

5 Nitrogen forms several oxides including N_2O_4 , N_2O and NO .

(a) A rocket uses the reaction between N_2O_4 and methylhydrazine, CH_3NHNH_2 , **equation 5.1**, to release a large amount of energy.




Some enthalpy changes of formation, ΔH_f , are shown in the table.

Substance	$\Delta H_f / \text{kJ mol}^{-1}$
$\text{CH}_3\text{NHNH}_2(\text{l})$	+54
$\text{N}_2\text{O}_4(\text{l})$	−20
$\text{CO}_2(\text{g})$	−394
$\text{H}_2\text{O}(\text{g})$	−242

Using the enthalpy changes of formation, ΔH_f , calculate the enthalpy change of reaction in **equation 5.1**.

enthalpy change of reaction = kJ mol^{-1} [3]

- (b) Under certain conditions nitrogen reacts with oxygen to make N_2O .



The enthalpy profile diagram for this reaction is shown in **Fig. 5.3**.




Fig. 5.3

- (i) Calculate the enthalpy change when 240 dm^3 of $\text{N}_2\text{O}(\text{g})$, measured at room temperature and pressure, is formed from N_2 and O_2 .

enthalpy change = kJ [2]

- (ii) What is the enthalpy change of formation, ΔH_f , of $\text{N}_2\text{O}(\text{g})$?

$\Delta H_f = \dots\dots\dots \text{ kJ mol}^{-1}$ [1]





Fig. 5.3 (repeated)

- (iii) The reaction in **equation 5.2** is reversible.



Calculate the activation energy, E_a , for the reverse reaction.

$$E_a \text{ (reverse reaction)} = \dots\dots\dots \text{ kJ mol}^{-1} \quad [1]$$

- (c) Describe and explain, using equations, how the concentration of ozone in the stratosphere is maintained.

.....

.....

.....

..... [2]

- (d) In the stratosphere, NO catalyses the breakdown of ozone.

Write **two** equations to show how NO catalyses this breakdown.

.....

..... [2]

[Total: 11]