



**EXAM PAPERS PRACTICE**

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

## Transition metals 1

2002

XVIII

1583

---

# CHEMISTRY

**AQA**

**AS & A LEVEL**

**Mark Scheme**


**Inorganic Chemistry**

For more help, please visit our website [www.exampaperspractice.co.uk](http://www.exampaperspractice.co.uk)



## Mark schemes


1

*Allow equation with  $\text{OH}^-$  provided equation showing formation of  $\text{OH}^-$  from  $\text{NH}_3$  given*

1

Green precipitate

1



1

Green precipitate

*effervescence incorrect so loses M4*

1

- (b) (i) Colourless / (pale) green changes to pink / purple (solution)

*Do not allow pale pink to purple*

1

Just after the end-point  $\text{MnO}_4^-$  is in excess / present

1



1

$$\text{Moles KMnO}_4 = 18.7 \times 0.0205 / 1000 = (3.8335 \times 10^{-4})$$

*Process mark*

1

$$\text{Moles Fe}^{2+} = 5 \times 3.8335 \times 10^{-4} = 1.91675 \times 10^{-3}$$

*Mark for  $M2 \times 5$* 

1

$$\text{Moles Fe}^{2+} \text{ in } 250 \text{ cm}^3 = 10 \times 1.91675 \times 10^{-3} = 0.0191675 \text{ moles in } 50 \text{ cm}^3$$

*Process mark for moles of iron in titration ( $M3$ )  $\times 10$* 

1

$$\text{Original conc Fe}^{2+} = 0.0191675 \times 1000 / 50 = 0.383 \text{ mol dm}^{-3}$$

*Answer for moles of iron ( $M4$ )  $\times 1000 / 50$* *Answer must be to at least 2 sig. figs. (0.38)*

1

[11]



2



1

 Mn<sup>2+</sup> OR Mn<sup>3+</sup>
*If catalyst incorrect can only score M1 and M3*

1

 (Possible because) Mn can exist in variable oxidation states

1

 E<sub>a</sub> lowered because oppositely charged ions attract

*These marks can be gained in any order*

1

 Mn<sup>3+</sup> (reduced) to Mn<sup>2+</sup> by C<sub>2</sub>O<sub>4</sub><sup>2-</sup> / equation


*M5 may appear before M2*

1

 Mn<sup>2+</sup> (oxidised (back)) to Mn<sup>3+</sup> by MnO<sub>4</sub><sup>-</sup> / equation

*M5 and M6 can be scored in unbalanced equations or in words showing:*


1

 (b) **Graph marks**


S-shaped curve must not rise significantly and must not fall rapidly initially.

 Starts on concentration axis **and** is levelling out (can level out on time axis or above but parallel to time axis)

*Cannot score graph marks (M1 and M2) if no axes and / or no labels*

1

1

### Explanation marks

Slope / rate increases as catalyst (concentration) forms

1

Slope / rate decreases as (concentration) of  $\text{MnO}_4^-$  ions / reactant(s) decreases (OR reactants are being used up)

*Explanation marks can be awarded independent of graph.*

1

[10]

3

- (a) 164.0

*Must be 1 decimal place*

1

- (b)  $17.1(\%) (= 28.0 \times 100 / Q_a)$

*Consequential on their (a)*

*Ignore precision but must be to at least 2 sig fig.*

*(i.e. accept 17 or 17.07)*

1

- (c) (i) Absorption depends on (proportional to) path length / distance travelled through solution

*Do not allow size.*

1

- (ii) To select the colour / frequency / wavelength that is (most strongly) absorbed (by the sample)

*Allow the filter is chosen to complement the colour of the solution*

1

- (iii) Quicker to analyse extracted samples than by titration / uses smaller volumes of solution

1

[5]

4

- (a) This question is marked using levels of response. Refer to the Mark Scheme Instructions for Examiners for guidance on how to mark this question.

All stages are covered and the explanation of each stage is generally correct and virtually complete.

Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 then stage 3.

**Level 3**  
5 – 6 marks

All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete.

Answer is mainly coherent and shows progression from stage 1 to stage 3.

**Level 2**  
3 – 4 marks



## EXAM PAPERS PRACTICE

Two stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete

Answer includes isolated statements but these are not presented in a logical order or show confused reasoning.

Level 1  
1 – 2 marks

Insufficient correct chemistry to gain a mark.

Level 0  
0 marks

### Indicative chemistry content

#### Stage 1: Electrons round P

- P has 5 electrons in the outside shell
- With 3 electrons from 3 fluorine, there are a total of 8 electrons in outside shell
- so 3 bond pairs, 1 non-bond pair

#### Stage 2: Electron pair repulsion theory

- Electron pairs repel as far as possible
- Lone pair repels more than bonding pairs

#### Stage 3: Conclusions

- Therefore, tetrahedral / trigonal pyramidal shape
- With angle of  $109.5^\circ$  decreased to  $107^\circ$

6

- (b)  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^7$

*Allow correct numbers that are not superscripted*

1

- (c) Too many electrons in d sub-shell / orbitals

1

- (d) Tetrahedral (shape)

1

$109.5^\circ$

*Allow  $109^\circ$*

1

[10]

5

- (a) Y

1

- (b) X

1

- (c) Jump in trend of ionisation energies after removal of fifth electron

Fits with an element with 5 outer electrons ( $4s^2 3d^3$ ) like V

1



- (d) Explanation: Two different colours of solution are observed

1

Because each colour is due to vanadium in a different oxidation state

1

- (e) **Stage 1:** mole calculations in either order

$$\text{Moles of vanadium} = 50.0 \times 0.800 / 1000 = 4.00 \times 10^{-2}$$

*Extended response*

*Maximum of 5 marks for answers which do not show a sustained line of reasoning which is coherent, relevant, substantiated and logically structured.*

1

$$\text{Moles of SO}_2 = pV / RT = (98\,000 \times 506 \times 10^{-6}) / (8.31 \times 293)$$

$$= 2.04 \times 10^{-2}$$

1

**Stage 2:** moles of electrons added to  $\text{NH}_4\text{VO}_3$

When  $\text{SO}_2$  (sulfur(IV) oxide) acts as a reducing agent, it is oxidised to sulfate(VI) ions so this is a two electron change

1

$$\text{Moles of electrons released when SO}_2 \text{ is oxidised} = 2.04 \times 10^{-2} \times 2$$

$$= 4.08 \times 10^{-2}$$

1

**Stage 3:** conclusion

But in  $\text{NH}_4\text{VO}_3$  vanadium is in oxidation state 5

1

$4.00 \times 10^{-2}$  mol vanadium has gained  $4.08 \times 10^{-2}$  mol of electrons  
therefore 1 mol vanadium has gained  $4.08 \times 10^{-2} / 4.00 \times 10^{-2} = 1$  mol  
of electrons to the nearest integer, so new oxidation state is  $5 - 1 = 4$

1

[11]

6

- (a)  $2\text{MnO}_4^- + 16\text{H}^+ + 5\text{C}_2\text{O}_4^{2-} \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 10\text{CO}_2$

*For all species correct / moles and species correct but charge incorrect*

1

*For balanced equation including all charges (also scores first mark)*

1

- (b) Manganate(VII) ions are coloured (purple)

1



## EXAM PAPERS PRACTICE

All other reactants and products are **not** coloured (or too faintly coloured to detect)

*Allow (all) other species are colourless*

*Allow  $Mn^{2+}$  are colourless / becomes colourless / pale pink*

1

- (c) The catalyst for the reaction is a reaction product

1

Reaction starts off slowly / gradient shallow

1

Then gets faster/rate increases / gradient increases

*Allow concentration of  $MnO_4^-$  decreases faster / falls rapidly*

1

- (d)  $Mn^{2+}$  ions


*Allow  $Mn^{2+}$  ions*

1

- (e)  $MnO_4^- + 8H^+ + 4Mn^{2+} \rightarrow 5Mn^{3+} + 4H_2O$

*Allow multiples*

1



1

[10]

7

- (a) Ti(IV) [Ar]

*Or  $1s^2 2s^2 2p^6 3s^2 3p^6$*

1

Ti(III) [Ar]3d<sup>1</sup>

*Or  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^1$*

1

Ti(III) has a d electron that can be excited to a higher level

*Allow idea that d electrons can be excited to another level (or move between levels)*

1

Absorbs one colour of light from white light

*Allow idea that light is absorbed*

1

Ti(IV) has no d electron so no electron transition with energy equal to that of visible light

*Allow Ti(IV) has no d electrons*



1

(b)	$[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	1
	$[\text{Cr}(\text{OH})_6]^{3-}$	1
	$[\text{CuCl}_4]^{2-}$	1
(c)	(i) Rapid determination of concentration <i>Or easy to get many readings</i>	1
	Does not use up any of the reagent/does not interfere with the reaction <i>Or possible to measure very low concentrations</i>	1
	(ii) Curve starts with small gradient (low rate)	1
	Because negative ions collide so $E_a$ high	1
	Curve gets steeper	1
	Because autocatalyst ( $\text{Mn}^{2+}$ ) formed	1
	Curve levels out approaching time axis <i>Can score this mark and next one ONLY with simple curve (that is curve with gradually decreasing gradient)</i>	1
	Because $\text{MnO}_4^-$ ions used up <i>5 max</i>	1

[15]



8



1

(ii)  $PV = nRT$   $n = PV/RT$   
(allow either formula but penalise contradiction)

1

$$n = \frac{110000 \times 102 \times 10^{-6}}{8.31 \times 298}$$

1

$$= 4.53 \times 10^{-3} \text{ (mol)}$$

(answer must have at least 3 sig. figs. Ignore units)

1

(iii) Moles of iron =  $4.5(3) \times 10^{-3} \text{ mol}$   
(allow consequent on (a)(ii))

(or =  $4.2(5) \times 10^{-3}$  if candidate uses given moles of hydrogen)

1

$$\text{Mass of iron} = 4.53 \times 10^{-3} \times 55.8 = 0.253 \text{ g}$$

(mark is for method mass = moles  $\times A_r$ )


(Mass of iron can be 56)

1

(iv)  $0.253 \times 100 / 0.263 = 96.1 \%$  (mark is for answer to 2 sig. figs.)  
(allow consequent on mass of iron. E.g. = 90% from  
 $4.2(5) \times 10^{-3}$  moles of  $\text{H}_2$  and Fe)  
(Do not allow answers greater than or equal to 100%)

1






*(ignore state symbols)*

1



1



1



$$= 4.53 \times 10^{-3}/6 = 7.55 \times 10^{-4}$$

*(Allow conseq, mark is for method (a)(iii)/6)*

1

Volume of dichromate = moles/concentration

$$= (7.55 \times 10^{-4} \times 1000)/0.0200$$

*(mark is for this method)*

1

$$V = 37.75 \text{ (cm}^3\text{)}$$

*(allow 37.7 to 37.8, allow no units but penalise wrong units)*

*(allow conseq on moles of dichromate)*


*(if value of  $3.63 \times 10^{-3}$  used answer is 30.2 to 30.3,*

*otherwise ans = moles  $\text{Fe}^{2+}/0.00012$ )*

*(if mole ratio wrong and candidate does not divide by 6,*

*max score is ONE for volume method)*

1



1

**[14]**



9

- (a) effect on reaction rate: catalyst provides an alternative reaction route.;

1

with a lower  $E_a$ ;

1

more molecules able to react or rate increased;

1

equilibrium: forward and backward rates changes by the same amount;

1

hence concentration of reactants and products constant or yield unchanged;

1

- (b) heterogeneous: catalyst in a different phase or state to that of the reactants;

1

active site: place where reactants adsorbed or attached or bond etc.;

1

reaction occurs or an explanation of what happens;

(allow absorbed)

1

reasons: large surface area;  
reduce cost or amount of catalyst;

2

catalyst poison: lead adsorbed;  
lead not desorbed or site blocked;  
(lead adsorbed irreversibly scores both of these marks)

2

- (c) reaction slow as: both ions negatively charged or ions repel;

1



Species;  
Balanced;

2



Species ;  
Balanced;

2

[17]



10

- (a) oxidation state of N in  $\text{Cu}(\text{NO}_3)_2$ : +5;

1

oxidation state of N in  $\text{NO}_2$ : +4;

1

oxidation product: oxygen;

1

- (b) copper-containing species:  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ ;

1

shape: octahedral;

1

- (c) (i) precipitate B:  $\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2$  or  $\text{Cu}(\text{OH})_2$  or name;


1

equation:  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow \text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2 + 2\text{NH}_4^+$

OR



and



1

- (ii)  $\text{NH}_3$  accepts a proton;

1


- (d) (i) identity:  $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$ ;

1

colour: deep blue;

1

equation:



1

- (ii)  $\text{NH}_3$  is an electron pair donor;

1

- (e) identity:  $[\text{CuCl}_4]^{2-}$ ;

1

colour: yellow-green;

1

shape: tetrahedral;

1



- (f) (i)  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10}$ ; 1
- (ii) role of Cu: a reducing agent; 1

[17]

11

- (a) most powerful reducing agent: Zn; 1
- (b) (i) reducing species:  $Fe^{2+}$  1
- (ii) oxidising species:  $Cl_2$ ; 1
- (c) (i) standard electrode potential 1.25 V; 1
- (ii) equation:  $Tl^{3+} + 2 Fe^{2+} \rightarrow 2Fe^{3+} + Tl$  + balanced; 1
- correct direction; 1
- (d) (i) moles  $KMnO_4 = 16.2 \times 0.0200 \times 10^{-3} = 3.24 \times 10^{-4}$ ; 1
- moles  $H_2O_2 = \text{Moles } KMnO_4 \times 5 / 2 = 8.10 \times 10^{-4}$ ; 1
- $8.10 \times 10^{-4}$  moles  $H_2O_2$  in  $25 \text{ cm}^3$
- $8.10 \times 10^{-4} \times 1000 / 25$  in  $1000 \text{ cm}^3 = 0.0324 \text{ mol dm}^{-3}$ ; 1
- hence  $\text{g dm}^{-3} = \text{mol dm}^{-3} \times M_r = 0.0324 \times 34 = 1.10$ ;  
 (penalise use of an incorrect  $H_2O_2$  to  $KMnO_4$  ratio by two marks) 1



- (ii)  $PV = nRT$ ; 1
- hence  $V = nRT / P$
- $= 8.10 \times 10^{-4} \times 8.31 \times 298 / 98000$ ; 1
- $= 2.05 \times 10^{-5}$ ; 1
- units  $\text{m}^3$ ;
- (mark consequentially to answers in (c)(i))*
- (allow correct answers with other units)*
- (answers to (c)(i) and (ii) must be to 3 significant figures; penalise once only)* 1
- [14]

12 [1]

13 [1]

14 (a)  $\text{FeCl}_3$  accepts electron pairs from water 1

Hence acts as a Lewis acid 1

$[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  donates protons 1

Hence acts as a Bronsted-Lowry acid 1

(b) The  $\text{Fe}^{2+}$  ion has a smaller charge to size ratio 1

Hence less polarising than  $\text{Fe}^{3+}$  or less weakening effect on O-H bonds 1


(c) (i)  $\text{V}_2\text{O}_5 + \text{SO}_2 \rightarrow \text{V}_2\text{O}_4 + \text{SO}_3$  1

$\text{V}_2\text{O}_4 + \text{O}_2 \rightarrow \text{V}_2\text{O}_5$  1



- (ii) Both ions are negative or ions repel

1




Species

1

Balanced

1



Species

1

Balanced

1

[13]

15

- (a)
- $\text{Fe} + \text{H}_2\text{SO}_4 \rightarrow \text{FeSO}_4 + \text{H}_2$

1

- (b)
- $\text{MnO}_4^- + 8\text{H}^+ + 5\text{Fe}^{2+} \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O} + 5\text{Fe}^{3+}$

1

- (c) Moles
- $\text{MnO}_4^-$
- in
- $19.6 \text{ cm}^3$

$$= 19.6 \times 0.022 \times 10^{-3} = 4.312 \times 10^{-4}$$

1

Moles  $\text{Fe}^{2+}$  in  $25 \text{ cm}^3$ 

$$= 5 \times 4.312 \times 10^{-4} = 2.156 \times 10^{-3}$$

1

Moles  $\text{Fe}^{2+}$  in  $250 \text{ cm}^3$ 

$$= 10 \times 2.156 \times 10^{-3} = 2.156 \times 10^{-2}$$

1

Mass  $\text{Fe}^{2+} = \text{moles} \times A_r$ 

$$A_r = 2.156 \times 10^{-2} \times 55.8 = 1.203 \text{ g}$$

1

Percentage by mass of carbon

$$= (1.270 - 1.203) \times 100 / 1.270$$

$$= 5.28\%$$

1

- (d) Repeat the titration and take an average of the concordant results

1



- (e) Analyse several samples from different parts of the molten iron

1

[9]

16

- (a) Equation:



Species

1

Balance

1

Colours:

e.g.  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  blue

1

e.g.  $[\text{CuCl}_4]^{2-}$  yellow/green

1

- (b) (i)  $\Delta E$ : The energy absorbed

1

h: Planck's constant

1

- (ii) Factor 1 Change of ligand

1

Factor 2 Change in oxidation state

1

Factor 3 Change in co-ordination number

1

[9]

17

- (a) Species  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$

1

Precipitate  $\text{Co}(\text{H}_2\text{O})_4(\text{OH})_2$

1

- (b)  $[\text{Co}(\text{NH}_3)_6]^{2+}$

1





(c)	Reaction	Oxidation	1
	Reactant	Oxygen in the air	1
(d)	R	Iodine	1
	Explanation	$[\text{Co}(\text{H}_2\text{O})_6]^{3+}$ oxidises $\text{I}^-$ to $\text{I}_2$	1

[7]

18

(a)	$[\text{Co}(\text{H}_2\text{O})_6]^{2+}$	1
	octahedral	
	<i>Only allow if species has 6 ligands but allow if M1 not given because charge missing</i>	1
(b)	$\text{CoCO}_3$	
	<i>Mark independently</i>	1
	Purple solid (allow pink) <i>Allow pink precipitate</i>	1
(c)	$[\text{Co}(\text{H}_2\text{O})_6]^{2+} + 6\text{NH}_3 \rightarrow [\text{Co}(\text{NH}_3)_6]^{2+} + 6\text{H}_2\text{O}$	
	<i>Allow <math>[\text{Co}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}</math></i>	
	Formula of product	1
	Balanced equation	1
(d)	$[\text{Co}(\text{NH}_3)_6]^{3+}$	
	<i>Allow <math>[\text{Co}(\text{NH}_3)_5\text{H}_2\text{O}]^{3+}</math></i>	1
	Oxidising agent	1



- (e)  $[\text{Co}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_3]^{2+}$

*Allow use of en  $[\text{Coen}_3]^{2+}$*

1

Entropy change for reaction is positive

*Mark independently*

1

Because 4 mol reactants form 7 mol products  
(or increase in number of particles)

*Or bidentate replaces unidentate*

1

- (f)  $[\text{CoCl}_4]^{2-}$

1

$\text{Cl}^-$  ligand too big to fit more than 4 round  $\text{Co}^{2+}$

*Allow  $\text{Cl}^-$  is bigger*

*Allow chlorine and Cl but NOT chlorine molecules.*

1

[13]

19

- (a) Partially filled/incomplete d sub-shell/orbital/shell

*Ignore reference to f orbitals*

*Do **not** allow d block*

*Do **not** allow half-filled d orbitals*

1

- (b) Has ligand(s)

*Allow molecules/ions with lone pairs*

1

linked by co-ordinate bonds

*Allow dative/donation of lone pair*

1

- (c) (Blue) light is absorbed (from incident white light)

1

Due to electrons moving to higher levels/electrons excited

*Allow  $d \rightarrow d$  transitions*

1

Red light (that) remains (is transmitted)/light that remains  
(transmitted light) is the colour observed

*Allow red light reflected*


1



- (d) (i) Circle round any O<sup>-</sup>  
*List principle*  
1
- Circle round either N  
1
- (ii)  $\text{EDTA}^{4-} + [\text{Co}(\text{H}_2\text{O})_6]^{2+} \rightarrow [\text{CoEDTA}]^{2-} + 6\text{H}_2\text{O}$   
*Allow missing square brackets*  
*Ignore state symbols*  
1
- (iii) Increase in entropy/ $\Delta S$  positive  
*Or increase in disorder*  
1
- Because 2 mol (of particles/molecules/species/entities) form 7 mol  
*Allow 'increase in number' as stated in words or as shown by any numbers deduced correctly from an incorrect equation*  
*Do not allow increase in ions/atoms*  
1
- (e) (i) Co-ordinate/dative/dative covalent bond  
*Allow pair of electrons donated by nitrogen/ligand*  
*Do not allow pair of electrons donated from Iron/Fe*  
1
- Covalent bond  
*Shared electron pair*  
1
- (ii) Transport of oxygen/O<sub>2</sub>  
*Allow any statement that implies oxygen carried (around the body)*  
*Do not allow transport of carbon dioxide (CO<sub>2</sub>). This also contradicts the mark (list principle)*  
1
- (iii) Because it bonds to the iron/haemoglobin  
*Allow blocks site*  
*/CO has greater affinity for haemoglobin*  
*/carboxyhaemoglobin more stable than oxyhaemoglobin*  
1
- Displaces oxygen  
*Or prevents transport of oxygen*  
*QoL*  
1



20



Correct product

1

Balanced equation

1

- (b) (i) Hydrogen bond

1

Oxygen (or nitrogen)

*Only score this mark if type of bond is correct*

1

- (ii) Co-ordinate

1

Nitrogen (or oxygen)

*Bond type must be correct to score this mark but allow M2 if bond is covalent*

1

- (c) Killing them or causing damage (medical side effects)

*Allow any correct side effect (e.g. hair loss)*

*Allow kills healthy (or normal) cells*

1

May attach to DNA in normal cells

1

[8]

21

- (a) Alternative route

*Allow mechanism outlined*

*allow forms intermediate species*

1

Lower activation energy

1

- (b) Variable oxidation state

*allow changes oxidation states*

1



*allow  $2\text{VO}_2$  instead of  $\text{V}_2\text{O}_4$*

1



1



- (ii) Poison attaches to surface  
*Allow blocks active site/surface*  
*Decreases surface area* 1
- (iii) Purify reactants  
*Allow remove impurities* 1
- [7]

22

- (a) (i) Propanone evaporates (or similar) 1
- Removes water (from the precipitate)  
*Accept 'removes impurities / excess reagents'.*  
*Accept 'salt insoluble in propanone'.* 1
- (ii) Add NaOH / NH<sub>3</sub> / Na<sub>2</sub>CO<sub>3</sub> 1
- No green ppt  
*Accept 'no visible change'.*  
*Must have correct reagent to score this mark.* 1
- (iii) Some salt dissolves (in propanone) **or** some lost in filtration **or** some Fe<sup>2+</sup> gets oxidised (to Fe<sup>3+</sup> in air)  
*Do not accept 'reaction reversible' or 'incomplete reaction' or similar.* 1
- (iv) Moles Fe<sup>2+</sup> =  $2.50 \times 10^{-2}$   
*Accept  $2.5 \times 10^{-2}$*  1
- $M_r$  of salt = 179.8  
*Allow 180*  
*Allow if 179.8 or 180 appears in a calculation.* 1
- Mass of salt =  $179.8 \times 2.5 \times 10^{-2} \times 0.95 = 4.27$  (g)  
*Correct answer with no working scores this mark only.*  
*Allow range 4.2 to 4.3 (g)* 1
- (v) 1.67 mol or correct ratio of 5FeC<sub>2</sub>O<sub>4</sub> : 3MnO<sub>4</sub><sup>-</sup> 1



*Accept multiples.*

1

- (c) (Insoluble) calcium ethanedioate coats surface

*Allow 'calcium ethanedioate is insoluble'.*

*Do not allow answers based on ethanedioic acid being a weak acid.*

*Do not accept 'acid used up' or 'reaction very fast'.*

1

- (d) Small amount of tea used **or** concentration of the acid in tea is low

*Accept 'high temperature decomposes the acid'.*

*Accept 'calcium ions in milk form a precipitate with the acid'.*

*Do not accept 'do not drink tea often' or similar.*

1

- (e) Mass of acid = 180.0 and mass of reagents = 450.0

*Accept 180 and 450.*

1

$$(180 / 450 \times 100 =) 40.0\%$$


*Do not penalise precision.*

*Correct answer without working scores this mark only.*

1

[14]

23



*Accept multiples.*

*Equation must have cobalt(II) hexaaqua ion.*

1

- (b) Ethanedioate ion reduces iron(III) ion **or**  
iron(III) ion oxidises ethanedioate ion

*Allow answer using equations.*

1

$E^\ominus (\text{CO}_2 / \text{C}_2\text{O}_4^{2-})$  more negative than  $E^\ominus (\text{Fe}^{3+} / \text{Fe}^{2+})$  **or**

$E^\ominus (\text{Fe}^{3+} / \text{Fe}^{2+}) > E^\ominus (\text{CO}_2 / \text{C}_2\text{O}_4^{2-})$

**or** e.m.f. positive **or** cell voltage = +1.26

1

[3]

24

- (a) An electron pair on the ligand

Is donated from the ligand to the central metal ion

1

1

- (b) Blue precipitate

Dissolves to give a dark blue solution

1


1



1



1




1



1

And the same number of bonds broken and made

1



1

Therefore, the free-energy change is negative

*M2 can only be awarded if M1 is correct*

1

[11]

25




1

Yellow-green/yellow/green

*Not necessary to indicate solution*


*Do not allow precipitate/solid*

1



*Allow + 4HCl → 4H<sup>+</sup>*

1




*Allow Cu(OH)<sub>2</sub>/copper hydroxide*

1

Blue precipitate/solid

*Ignore shades*

1



*Allow any balanced equation/equations leading to this hydroxide or Cu(OH)<sub>2</sub>*

*But must use ammonia*

1






- (c) Y is  $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$

1

Deep/dark/royal blue solution

*QoL*

1



*Accept equation for formation from  $\text{Cu}(\text{OH})_2$*

1

- (d) Z is  $\text{CuCO}_3$


*Allow copper carbonate*

1

Green solid/precipitate

*Allow blue-green precipitate*

1



1

- (e) (i)  $\text{Cu}^{2+}(\text{aq}) + \text{Fe}(\text{s}) \rightarrow \text{Cu}(\text{s}) + \text{Fe}^{2+}(\text{aq})$

*Allow hydrated ions*

*State symbols not essential but penalise if wrong*

1

Blue

*Do not allow description of solids*

1

Green

*Allow yellow/(red-)brown/orange*

1

- (ii) Any two correct points about copper extraction from two of these three categories:

Any relevant mention of lower energy consumption

*Do not allow reference to electricity alone or to temperature alone.*

Any relevant mention of benefits of less mining (of copper ore)

*Allow avoids depletion of (copper ore) resources*

Less release of  $\text{CO}_2$  (or CO) into the atmosphere

*Not just greenhouse gases. Must mention  $\text{CO}_2$  or CO*

Max 2

[17]


**26**

- (a) Same phase/state 1
- (b) Because only exist in one oxidation state  
*Allow do not have variable oxidation states* 1
- (c)  $2\text{I}^- + \text{S}_2\text{O}_8^{2-} \rightarrow \text{I}_2 + 2\text{SO}_4^{2-}$   
*Ignore state symbols*  
*Allow multiples* 1
- (d) Both (ions) have a negative charge  
*Or both have the same charge*  
*Or (ions) repel each other*  
*Do not allow both molecules have the same charge (contradiction)* 1
- (e)  $2\text{Fe}^{2+} + \text{S}_2\text{O}_8^{2-} \rightarrow 2\text{Fe}^{3+} + 2\text{SO}_4^{2-}$  1
- $2\text{Fe}^{3+} + 2\text{I}^- \rightarrow 2\text{Fe}^{2+} + \text{I}_2$  1
- Equations can be in any order*
- Positive and negative (ions)/oppositely charged (ions)  
*Mark independently* 1
- (f) Equations 1 and 2 can occur in any order  
*Allow idea of  $\text{Fe}^{3+}$  converted to  $\text{Fe}^{2+}$  then  $\text{Fe}^{2+}$  converted back to  $\text{Fe}^{3+}$*  1

**[8]**



27



1

- (b) (i) Enthalpy change for formation of 1 mol of substance

*Allow heat energy change. NOT energy*

1

From its elements

1


Reactants and products/all substances in their standard states

*Or normal states at 298 K, 1 bar (100 kPa)*

1



1

*Or labelled diagram*

1

$$= 193 + 590 + 1150 + 158 + (2 \times -348) - 2602$$

1

$$= -1207 \text{ kJ mol}^{-1}$$

*Correct answer scores 3**-842 scores 2 (transfer error)**-859 scores 1 only (using one E.A.)**Units not required, wrong units lose 1 mark*

1

- (c) Electrostatic attraction stronger/ionic bonding stronger/attraction between ions stronger/more energy to separate ions

*Molecular attraction/atoms/intermolecular forces CE=0*

1

Because fluoride (ion) smaller than chloride

*Do not allow F or fluorine*

1

*Can be on cycle/diagram*

1

$$= -141 \text{ kJ mol}^{-1}$$

*Correct answer scores 2**Units not required, wrong units lose 1 mark*

1



(ii) Decreases

*If ans to (d)(i) positive allow increases*

1

Reaction exothermic/ $\Delta H$  -ve

*If (d)(i) +ve allow endothermic/ $\Delta H$  + ve*

1

(Equilibrium) shifts to left/backwards

(as temperature rises)/equilibrium

opposes the change

*If (d) (i) +ve allow shifts to right/forwards/equilibrium opposes the change*

*If no answer to (d) (i) assume -ve  $\Delta H$  used*

*If effect deduced incorrectly from any  $\Delta H$  CE = 0 for these 3 marks*

1

(e) u.v. absorbed: electrons/they move to higher energy  
(levels)/electrons excited

1

visible light given out: electrons/they fall back down/move to  
lower energy (levels)

*Must refer to absorbing u.v. NOT visible light or this must be implied.*

1

[17]

28

(a) (i) Ammonia

*If reagent is missing or incorrect cannot score M3*

1

Starts as a pink (solution)

1

Changes to a yellow/straw (solution)

*Allow pale brown*

*Do not allow reference to a precipitate*

1

(ii) (dark) brown

*Do not allow pale/straw/yellow-brown (i.e. these and other shades except for dark brown)*

1



- (b) (i) Ruby/red-blue/purple/violet/green


*Do not allow red or blue**If ppt mentioned contradiction/CE = 0*

1

Green

*If ppt mentioned contradiction/CE = 0*

1



1


Formula of product

*Can score this mark in (b) (ii)*

1

- (ii)
- $\text{H}_2\text{O}_2 + 2\text{e}^- \rightarrow 2\text{OH}^-$

1

*Allow 1 mark out of 2 for a balanced half-equation such as  $\text{Cr(III)} \rightarrow$*  *$\text{Cr(VI)} + 3\text{e}^-$* *or  $\text{Cr}^{3+} + 4\text{H}_2\text{O} \rightarrow \text{CrO}_4^{2-} + 8\text{H}^+ + 3\text{e}^-$  etc**also for  $2\text{Cr(III)} + 3\text{H}_2\text{O}_2 \rightarrow 2\text{CrO}_4^{2-}$  (unbalanced)*

2

Yellow

*Do not allow orange*

1

- (c)
- $2\text{MnO}_4^- + 6\text{H}^+ + 5\text{H}_2\text{O}_2 \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{O}_2$

*if no equation and uses given ratio can score M2, M3, M4 & M5*

1

$$\text{Moles MnO}_4^- = (24.35/1000) \times 0.0187 = \underline{4.55 \times 10^{-4}}$$

*Note value must be quoted to at least 3 sig. figs.**M2 is for  $4.55 \times 10^{-4}$* 

1

$$\text{Moles H}_2\text{O}_2 = (4.55 \times 10^{-4}) \times \underline{5/2} = 1.138 \times 10^{-3}$$

*M3 is for  $\times 5/2$  (or  $7/3$ )**Mark consequential on molar ratio from candidate's equation*

1



Moles  $\text{H}_2\text{O}_2$  in  $5 \text{ cm}^3$  original

*M4 is for  $\times 10$*

1

$$= (1.138 \times 10^{-3}) \times \underline{10} = 0.01138$$

$$\text{Original } [\text{H}_2\text{O}_2] = 0.01138 \times \underline{(1000/5)} = 2.28 \text{ mol dm}^{-3}$$

(allow 2.25-2.30)

*M5 is for consequentially correct answer from (answer to mark 4)  $\times$  (1000/5)*

*Note an answer of between 2.25 and 2.30 is worth 4 marks*

*If candidate uses given ratio 3/7 max 4 marks:*

**M1:** Moles of  $\text{MnO}_4^- = \underline{4.55 \times 10^{-4}}$

**M2:** Moles  $\text{H}_2\text{O}_2 = (4.55 \times 10^{-4}) \times \underline{7/3} = 1.0617 \times 10^{-3}$

**M3:** Moles  $\text{H}_2\text{O}_2$  in  $5 \text{ cm}^3$  original

$$= (1.0617 \times 10^{-3}) \times 10 = 0.01062$$

**M4:** Original  $[\text{H}_2\text{O}_2] = 0.01062 \times (1000/5) = 2.12 \text{ mol dm}^{-3}$

(allow 2.10 to 2.15)

1

[17]

29

(a) Brown ppt/solid

1

Gas evolved/effervescence

1



*Must be stated, Allow  $\text{CO}_2$  evolved. Do not allow  $\text{CO}_2$  alone*

*Correct iron product (1) allow  $\text{Fe}(\text{OH})_3$  and in equation*

*Balanced equation (1)*

2


(b) White ppt/solid

1


Colourless Solution

*Only award M2 if M1 given or initial ppt mentioned*

1



1



*Allow formation of  $[\text{Al}(\text{H}_2\text{O})_{6-x}(\text{OH})_x]^{(x-3)-}$  where  $x = 4, 5, 6$*

*Allow product without water ligands*

*Allow formation of correct product from  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$*

1



(c) Blue ppt/solid

1



(Dissolves to give a) deep blue solution

*Only award M2 if M1 given or initial ppt mentioned*

1




1



1

(d) Green/yellow solution

1



1

[14]





30

- (a) (i) absorbs (certain frequencies of) (white) light / photons  
*not absorbs white / u.v. light*

1

d electrons excited / promoted  
*or d electrons move between levels / orbitals*  
*d electrons can be implied elsewhere in answer*

1

the colour observed is the light not absorbed / light  
 reflected / light transmitted  
*allow blue light transmitted*  
*penalise emission of light in M3*

1

- (ii)  $\Delta E$  is the energy gained by the (excited) electrons (of  $\text{Cu}^{2+}$ )  
*allow:*
- energy difference between orbitals / sub-shells
  - energy of photon / light absorbed
  - change in energy of the electrons energy lost by excited electrons
  - energy of photon / light emitted

1

$h$  (Planck's) constant

1

$\nu$  frequency of light (absorbed by  $\text{Cu}^{2+}(\text{aq})$ )  
*do not allow wavelength*  
*If energy lost / photon lost / light emitted in M1 do not penalised*  
*light emitted*

1

- (iii)  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 4\text{Cl}^- \rightarrow [\text{CuCl}_4]^{2-} + 6\text{H}_2\text{O}$   
*note that  $[\text{CuCl}_4]^{-2-}$  is incorrect*  
*penalise charges shown separately on the ligand and overall*  
*penalise HCl*

1


tetrahedral

1

$\text{Cl}^-$  / Cl / chlorine too big (to fit more than 4 round Cu)  
*allow*  
*water smaller than  $\text{Cl}^-$*   
*explanation that change in shape is due to change in*  
*co-ordination number*

1

(b)



*allow:*


- ion drawn with any bond angles
- ion in square brackets with overall / 2- charge shown outside the brackets
- ion with delocalised O=C-O bonds in carboxylate group(s)

1

lone pair(s) on O<sup>-</sup> / O

*allow position of lone pair(s) shown on O in the diagram even if the diagram is incorrect.*

1



product correct

1

equation balanced

1

6

*note can only score M3 and M4 if M1 awarded or if complex in equation has 2 waters and 2 ethanedioates*


1

octahedral

*If this condition is satisfied the complex can have the wrong charge(s) to allow access to M3 and M4 but not M1*

1

(ii)



ignore charges

diagram must show both ethanedioates with correct bonding

ignore water

1

90°

allow 180°

 mark bond angle independently but penalise if angle incorrectly  
labelled / indicated on diagram

1

[17]

31

- (a) Incomplete (or partially filled) d orbitals/sub-shells

Do not allow d shell

1

- (b) Variable oxidation states

1

- (c) (i)
- $[\text{H}_3\text{N}-\text{Ag}-\text{NH}_3]^+$

 Allow  $[\text{Cl}-\text{Ag}-\text{Cl}]^-$  or similar Cu(I) ion

Allow compounds in (i), (ii) and (iii) (eg Cl-Be-Cl)

Allow no charge shown, penalise wrong charge(s)

1

- (ii) Cis platin drawn out as square planar

 Allow  $\text{NiX}_4^{2-}$  etc

1

- (iii)
- $[\text{CuCl}_4]^{2-}$
- drawn out as tetrahedral ion

 Or  $[\text{CoCl}_4]^{2-}$  drawn out

1

- (d) (i)
- $\text{SO}_2 + 1/2\text{O}_2 \rightarrow \text{SO}_3$

Allow multiples

 Allow  $\text{SO}_2 + 1/2\text{O}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$ 

ignore state symbols

1

- (ii) In a different phase/state (from the reactants)

1



*can be in either order*

1



*allow multiples*

1

(iv) Surface area is increased

1

By use of powder or granules or finely divided

*Allow suspending/spreading out onto a mesh or support*

1

(e) (i) Forms two or more co-ordinate bonds

*Allow more than one co-ordinate bond or donates more than 1 electron pair.*

*Do not allow "has more than one electron pair"*

*Allow uses more than one atom to bond (to TM)*

1

(ii) Number of product particles > Number of reactant particles

*Allow molecules/entities instead of particles*

*Penalise incorrect numbers (should be 2→5)*

1

Disorder increases or entropy increases

(or entropy change is positive)

*Allow  $\Delta G$  must be negative because  $\Delta H = 0$  and  $\Delta S$  is +ve*

1

(iii) 6

1

Cyanide strongly bound to Co (by co-ordinate/covalent bond)

1

[16]

32

(a) Plots all of the points correctly  $\pm$  one square

1

Straight line through the points is best fit

*Candidate does not have to extrapolate line to the origin.*

*Line must pass through the origin  $\pm$  1 square.*

*Lose this mark if the candidate's line is doubled or kinked.*

*Allow line that doesn't pass through the origin if one or more points are misplotted.*

1



- (b)  $7.6 \pm 0.1 \times 10^{-2} \text{ (mol dm}^{-3}\text{)}$

*Do not penalise precision, but at least 2 significant figures.*

1

[3]

33

$\text{MnO}_4^-$  will oxidise the chloride ion / reaction of  $\text{MnO}_4^-$  and  $\text{Cl}^-$  feasible

*Accept converse argument with  $\text{Cr}_2\text{O}_7^{2-}$ .*

*Accept calculations of overall  $E^\circ$  values.*

1

Larger volume needed

1

[2]

34

(a) (i)  $\text{EDTA}^{4-} + [\text{Cu}(\text{H}_2\text{O})_6]^{2+} \rightarrow [\text{Cu}(\text{EDTA})]^{2-} + 6\text{H}_2\text{O}$

1

(ii)  $(\text{Mol EDTA} = (6.45/1000) \times 0.015 = )9.68 \times 10^{-5} \text{ mol Cu(II)}$

1

$\text{Conc. Cu(II)} = ((9.68 \times 10^{-5}) / 0.025 =) 0.00387 \text{ mol dm}^{-3}$

*Correct answer without working gains M2 only.*

1

(b) Samples may not be consistent throughout the river  
OR

Concentration may vary over time

*Ignore comments on technique.*

1

(c)  $[\text{Ag}(\text{NH}_3)_2]^+$

*Accept name eg diamminesilver(I) ion.*

1

aldehyde

*Allow CHO.*

1

[6]

35

(a) (ligand) substitution

*Allow 'ligand exchange'.*

1

(b) To displace the equilibrium to the right


*To ensure reaction goes to completion.*

1

To improve the yield

*Allow 'to replace all chlorines'.*

1



*Allow correct ionic equations  $\text{PtCl}_4^{2-} + 4\text{I}^- \rightarrow \text{PtI}_4^{2-} + 4\text{Cl}^-$*

*Allow multiples and fractions.*

1

(ii)  $= (780.9) \times 100 / (415.3 + 664)$

*Working must be clearly shown.*

*Allow one mark for correct relationship even if  $M_r$  values are incorrect eg using values from ionic equation.*

1

$= 72.4$

*Allow 72%*

1



*Ignore state symbols even if incorrect.*

*This equation only.*

1

(ii) Stops the reverse reaction / equilibrium displaced to the right

1

(e) Number of steps in the process

*Allow 'equilibrium may lie on the reactant side' / side reactions / isomer formation.*

1

Losses at each stage of the synthesis

*Equilibrium losses or practical losses or yield not 100% for each step.*

1

(f) Minimum amount of hot solvent

*Accept 'small' for minimum.*

*Accept water.*

1

Cool / crystallise

1

Filter

1

(g) (i) Small amounts are more likely to kill cancer cells rather than the patient

1

(ii) Wear gloves / wash hands after use


*Ignore masks.*

*Apply the list principle if more than one answer.*


1



36



1



1

two negative ions repel / lead to reaction that is slow / lead to reaction that has high  $E_a$


1

iron able to act because changes its oxidation state  
*allow iron has variable oxidation state*

1

With iron ions have alternative route / route with lower activation energy

1



*can have  $\text{H}_2\text{O}$  on LHS and  $\text{H}_3\text{O}^+$  on R*

*do not penalise further hydrolysis equations*

*allow high charge density*

1

$\text{Fe}^{3+}$  ion has higher charge (to size ratio) (than  $\text{Fe}^{2+}$ )

1

increases polarisation of co-ordinated water / attracts O  
releasing an  $\text{H}^+$  ion / weakens O—H bond

1



- (ii)  $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 6\text{Fe}^{3+}$   
 or 6 mol Fe(II) react with 1 mol dichromate  
 If factor of 6 not used max = 3 for M2, M4 and M5  
 e.g. 1:1 gives ans = 8.93 to 8.98% (scores 3)

1

$$\text{moles dichromate} = 23.6 \times 0.218/1000 = 5.14 \times 10^{-4}$$

1

$$\text{moles iron} = 5.14 \times 10^{-4} \times 6 = 0.00309$$

M3 also scores M1

1

$$\text{mass iron} = 0.00309 \times 55.8 = 0.172$$

Mark is for moles of iron  $\times$  55.8 conseq

Allow use of 56 for iron

1

$$\% \text{ by mass of iron} = 0.172 \times 100/0.321 = 53.7\%$$

Answer must be to at least 3 sig figures allow 53.6 to 53.9

Mark is for mass of iron  $\times$  100/0.321 conseq

1

- (c) brown precipitate / solid

Allow red-brown / orange solid

Not red or yellow solid

1

bubbles (of gas) / effervescence/ fizz


Allow gas evolved / given off

Do not allow just gas or  $\text{CO}_2$  or  $\text{CO}_2$  gas


1



Allow



Use of  $\text{Na}_2\text{CO}_3$



1

[16]





37

- (a)
- $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10}$

*allow [He] 2s<sup>2</sup> . or [Ne] 3s<sup>2</sup>.or [Ar]3d<sup>10</sup>*

1

d sub-shell / shell / orbitals / sub-level full (or not partially full)

*can only score M2 if d<sup>10</sup> in M1 correct**allow 'full d orbital' if d<sup>10</sup> in M1**do not allow d block*

1

- (b) atom or ion or transition metal bonded to / surrounded by one or more ligands

*Allow Lewis base instead of ligand*

1

by co-ordinate / dative (covalent) bonds / donation of an electron pair

*can only score M2 if M1 correct*

1

- (c) H
- <sub>2</sub>
- / hydrogen

*do not allow H*

1

no lone / spare / non-bonded pair of electrons

*only score M2 if M1 correct or give 'H' in M1*

1

- (d) (i) +2 or 2+ or Pd
- <sup>2+</sup>
- or II or +II or II+ or two or two plus

1

- (ii) tetrahedral

*these shapes can be in any order*

1

square planar

*allow phonetic spelling e.g. tetrahydral*

1

[9]

38

- (a) Electron
- pair
- donor

*Allow lone pair donor*

1

- (b)
- $[\text{Cu}(\text{H}_2\text{O})_6]^{2+} + 2\text{NH}_3 \rightarrow \text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2 + 2\text{NH}_4^+$

1



# EXAM PAPERS PRACTICE

(Blue solution) gives a (pale) blue precipitate/solid

*M2 only awarded if M1 shows Bronsted-Lowry reaction*

1



*Allow formation in two equations via hydroxide*

1

(Blue solution) gives a dark/deep blue solution

*If (b) and (c) are the wrong way around allow one mark only for each correct equation with a correct observation (max 2/4)*

*M2 only awarded if M1 shows Lewis base reaction*

1

(d) (Start with) green (solution)

1

Green precipitate of  $\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2$  /  $\text{Fe}(\text{OH})_2$  / iron(II) hydroxide

*Do not allow observation if compound incorrect or not given*

1

Slowly changes to brown solid

*Allow red-brown ppt*

*Allow turns brown or if precipitate implied*

*Can only score M3 if M2 scored*


1

(Iron(II) hydroxide) oxidised by air (to iron(III) hydroxide)

*Allow  $\text{Fe}(\text{OH})_2$  oxidised to  $\text{Fe}(\text{OH})_3$  by air /  $\text{O}_2$*

*Ignore equations even if incorrect*

1



*For correct Al species*

1

*For correct balanced equation*


*Allow equation with formation of  $3[\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_3] +$  from 1 mol*

*$[\text{Al}(\text{H}_2\text{O})_6]^{3+}$*

1

White precipitate

1



1



## EXAM PAPERS PRACTICE

Complex with 3 en showing 6 correct bonds from N to Co

*Ignore charge*

*Accept N – N for ligand*

*Ignore incorrect H*


*If C shown, must be 2 per ligand*

1

Co-ordinate bonds (arrows) shown from N to Co

*Can only score M3 if M2 correct*

1



*For Co(III) species*

1

*For balanced equation (others are possible)*



*If en used can score M4 and M5 only*

*If Cu not Co, can only score M2 and M3*

*Allow  $\text{N}_2\text{C}_2\text{H}_8$  in equations*

1

[17]

39

(a) Variable oxidation state

1

eg Fe(II) and Fe (III)

*Any correctly identified pair*

*Allow two formulae showing complexes with different oxidation states even if oxidation state not given*

1

(Characteristic) colour (of complexes)

1

eg  $\text{Cu}^{2+}(\text{aq})$  /  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  is blue


*Any correct ion with colour scores M3 and M4*

*Must show (aq) or ligands OR identified coloured compound e.g.  $\text{CoCO}_3$*

1

(b) Tetrahedral

1




*Any correct complex*

*(Note charges must be correct)*

Square planar

1



*Any correct complex*

1

Linear

*Do not allow linear planar*

1



*$[\text{AgCl}_2]^-$  etc*

1



*If equation does not show increase in number of moles of particles*

*CE = 0/3 for (c)(ii)*

*If no equation, mark on*

1

(ii) 2 mol of reactants form 7 mol of products

*Allow more moles/species of products*

*Allow consequential to (c)(i)*

1

Therefore disorder increases

1

Entropy increases / +ve entropy change / free-energy change is negative

1

(iii) Moles EDTA =  $6.25 \times 0.0532 / 1000 = (3.325 \times 10^{-4})$

1

Moles of  $\text{Ca}^{2+}$  in  $1 \text{ dm}^3 = 3.325 \times 10^{-4} \times 1000 / 150 = (2.217 \times 10^{-3})$

*Mark is for  $M1 \times 1000 / 150$  OR  $M1 \times 74.1$*

*If ratio of  $\text{Ca}^{2+} : \text{EDTA}$  is wrong or  $1000 / 150$  is wrong, CE and can score M1 only*

*This applies to the alternative*

1

Mass of  $\text{Ca}(\text{OH})_2 = 2.217 \times 10^{-3} \times 74.1 = 0.164 \text{ g}$

*$M1 \times 74.1 \times 1000 / 150$*

*Answer expressed to 3 sig figs or better*

*Must give unit to score mark*

*Allow 0.164 to 0.165*

1




40

## (a) Reaction 1

**General principles in marking this question***Square brackets are not essential**Penalise charges on individual ligands rather than on the whole complex**Reagent and species can be extracted from the equation**Ignore conditions such as dilute, concentrated, excess**Reagent must be a compound NOT just an ion**Equations must start from  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  except in part (b)**Mark reagent, species and equation independently*ammonia ( $\text{NH}_3$ ) (solution) /  $\text{NaOH}$ 

1

*Do not allow  $\text{OH}^-$  for reagent**Product 1, balanced equation 1**Allow either equation for ammonia*

2

## (b) Reaction 2

Ammonia (conc / xs)

1



*Product 1, balanced equation 1**Note that the equation must start from the hydroxide* *$[\text{Cu}(\text{H}_2\text{O})_4(\text{OH})_2]$* 

2


## (c) Reaction 3

 $\text{Na}_2\text{CO}_3$  / any identified soluble carbonate /  $\text{NaHCO}_3$ *Do not allow  $\text{NaCO}_3$  or any insoluble carbonate but mark on*

1



OR with  $\text{NaHCO}_3$



*Product 1, balanced equation 1*


2

(d) **Reaction 4**

$\text{HCl}$  (conc / xs) /  $\text{NaCl}$

*Allow any identified soluble chloride*

1



*Product 1, balanced equation 1*

2

[12]

41

- (a) Stop the formation of  $\text{MnO}_2$  / Ensures all  $\text{MnO}_4^-$  reacts to form  $\text{Mn}^{2+}$  / becomes colourless

1

- (b) Weak acid / Does not supply sufficient  $\text{H}^+$

1

- (c) It is self-indicating / Purple to colourless end-point or vice versa

*If colours mentioned they must be correct.*

1

[3]

42

- (a) In each of **P** and **Q** the oxidation state of Cr is +3 / both contain  $\text{Cr}^{3+}$

*If oxidation states are different lose M1 and M2*

1

In each of **P** and **Q** the electron configuration is the same /  $d^3$  /  $3d^3$

*Do not allow just same number of electrons*

1

Ligands are different

1

Different energies of (d) electrons / different split of (d) electron energy levels /  
different energy gap of (d) electrons / different (d) orbital energy

1



## EXAM PAPERS PRACTICE


Different wavelengths / frequencies / energies of light / colours (of light) are absorbed (by the d electrons)

*Reference to emission and / or uv light but not to visible loses M5 and M6*

1

Different wavelengths / frequencies / energies of light / colours (of light) are transmitted / reflected

1



*Allow  $\text{NH}_2\text{C}_2\text{H}_4\text{NH}_2$  and  $\text{CH}_2\text{NH}_2\text{CH}_2\text{NH}_2$*

*Allow partial substitution*

*Do not allow en or other formulae for M1 but can score M2*

1

4 particles form 7 particles / increase in number of particles

*Allow molecules, entities, ions, moles instead of particles*

*Do not allow atoms*

*Can score M2 if numbers match candidates incorrect equation provided number of particles increases*

1

disorder / entropy increases /  $\Delta S$  positive

*Cannot score M3 if number of particles stated or in equation is the same or decreases*

1

$\Delta H$  is approx. zero / no net change in bond enthalpies

*Allow same number and type of bonds broken and formed*


1

$\Delta G$  is negative /  $\Delta G \leq 0$

*Mark M4 and M5 independently*

1

(c) (i)



*Correct displayed structure*

*Must show all three N-H bonds on each N*

*Ignore arrows and lone pairs, attempt to show shape*

*Ignore charges on atoms in structure for M1*

1



Bond angle  $90^\circ$

*Allow 87 to 93 degrees*

*Allow this angle for any complex with 4 ligands eg if  $\text{NH}_2$  or  $\text{Cl}$  used instead of  $\text{NH}_3$*


1

Charge of zero

*Award this mark if no charge shown on structure but if charges shown on ligands in M1 must state that overall charge = 0*

*Allow M3 only if cisplatin is correct OR if trans form OR if  $\text{NH}_3$  not displayed OR if  $\text{NH}_2$  used instead of  $\text{NH}_3$*

1



*If formula of cisplatin is incorrect, mark consequentially provided  $\text{H}_2\text{O}$  replaces  $\text{Cl}^-$  and charge on complex increases by one*

1

(iii) Use in small amounts / short bursts / target the application / monitor the patients


*Allow: Give patient time between doses*

1



*Allow multiples*

1



1

Acts as a catalyst / lowers the activation energy

1

Speeds up the (overall) reaction (between  $\text{SO}_2$  and oxygen)

1

[20]

43

(a) (i) Correctly plots all points ( $\pm$  one square) and draws straight line of best fit

*Lose this mark if the candidate's line is doubled or kinked.*

*Lose this mark if the line does not pass within one square of the origin, extending the line if necessary.*

1

Plotted points cover over half of grid

1

(ii)  $0.046 \pm 0.002$  ( $\text{mol dm}^{-3}$ )

1





0.088 to 0.096 ( $\text{mol dm}^{-3}$ )

Allow  $M1 \times 2$

Allow two marks for correct answer.

Answer must be to at least two significant figures.

1

(iii) Total volume =  $(100 \times 0.1) / 0.04 = 250 \text{ (cm}^3\text{)}$

Allow any correct alternative method of working.

1

Therefore add  $150 \text{ cm}^3$

Correct answer without working scores M2 only.

1

(b) Iron needed for haemoglobin / for red blood cells / to carry oxygen around the body

Accept well-water may contain eg  $\text{Ca}^{2+}$  ions / dissolved minerals  
that are good for bones / teeth etc.

1

[7]

44

(a) moles of  $\text{Cr}_2\text{O}_7^{2-}$  per titration =  $21.3 \times 0.0150 / 1000 = \underline{3.195 \times 10^{-4}}$

1

$(\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 6\text{Fe}^{3+}) \text{ Cr}_2\text{O}_7^{2-}:\text{Fe}^{2+} = 1:6$

If 1:6 ratio incorrect cannot score M2 or M3

1

moles of  $\text{Fe}^{2+} = 6 \times 3.195 \times 10^{-4} = 1.917 \times 10^{-3}$

Process mark for  $M1 \times 6$  (also score M2)

1

original moles in  $250 \text{ cm}^3 = 1.917 \times 10^{-3} \times 10 = 1.917 \times 10^{-2}$

Process mark for  $M3 \times 10$

1

mass of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O} = 1.917 \times 10^{-2} \times 277.9 = 5.33 \text{ (g)}$

Mark for answer to  $M4 \times 277.9$

(allow 5.30 to 5.40)

Answer **must** be to at least 3 sig figs

Note that an answer of 0.888 scores M1, M4 and M5 (ratio 1:1  
used)

1

(b) (Impurity is a) reducing agent / reacts with dichromate / impurity is a version of  $\text{FeSO}_4$   
with fewer than 7 waters (not fully hydrated)

Allow a reducing agent or compound that that converts  $\text{Fe}^{3+}$  into  
 $\text{Fe}^{2+}$

1



Such that for a given mass, the impurity would react with more dichromate than a similar mass of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

OR for equal masses of the impurity and  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , the impurity would react with more dichromate.

*Must compare mass of impurity with mass of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$*

1

[7]

45

- (a) Manganate would oxidise / react with  $\text{Cl}^-$

1

Because  $E^\ominus$  for  $\text{MnO}_4^-$  is more positive than that for  $\text{Cl}^2 / 1.51 - 1.36 = +0.15$  (V)

*Must refer to data from the table for M2.*

1

- (b) Moles of  $\text{H}^+ = 25 \times 0.0200 \times 8 / 1000 = 4.00 \times 10^{-3}$

1

Moles of  $\text{H}_2\text{SO}_4 = 2.00 \times 10^{-3}$  ( $4.00 \times 10^{-3} / 2$ )

*Allow consequential marking on incorrect moles of  $\text{H}^+$*

1

Volume  $\text{H}_2\text{SO}_4 = 4.00$  ( $\text{cm}^3$ ) ( $2.00 \times 10^{-3} \times 1000 / 0.500$ )

*Allow consequential marking on incorrect moles of  $\text{H}_2\text{SO}_4$*

*Accept  $4 \text{ cm}^3$ .*

*$8 \text{ cm}^3$  scores 2 marks.*

*Do not penalise precision.*

*Correct answer without working scores M3 only.*

1

- (c) (i)  $\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightarrow \text{MnO}_2 + 2\text{H}_2\text{O}$

*Allow multiples, including fractions.*

*Ignore state symbols.*

1

- (ii) Can't see end point due to brown colour

1

Larger titre (than expected)

*Allow the idea that with two reactions can't make use of titre in calculations.*

*Do not allow 'an inaccurate result' without qualification.*

1

- (d) Solution (very) dilute / lots of water

1

[9]

46

- (a) (i) Flask with side arm

1

- (c)  $180^\circ / 180 / 90$

*Allow any angle between 85 and 95*

*Do not allow 120 or any other incorrect angle*

*Ignore units eg  $^\circ\text{C}$*

1

- (d) (i)  $3 : 5 / 5 \text{ FeC}_2\text{O}_4 \text{ reacts with } 3 \text{ MnO}_4^-$

*Can be equation showing correct ratio*

1

- (ii) **M1** Moles of  $\text{MnO}_4^-$  per titration =  $22.35 \times 0.0193 / 1000 = 4.31 \times 10^{-4}$

Method marks for each of the next steps (no arithmetic error allowed for M2):

*Allow  $4.3 \times 10^{-4}$  (2 sig figs)*

*Allow other ratios as follows:*

*eg from given ratio of 7/3*

1

**M2** moles of  $\text{FeC}_2\text{O}_4 = \text{ratio from (d)(i) used correctly} \times 4.31 \times 10^{-4}$

$$\text{M2} = 7/3 \times 4.31 \times 10^{-4} = 1.006 \times 10^{-3}$$

1

**M3** moles of  $\text{FeC}_2\text{O}_4$  in  $250 \text{ cm}^3 = \text{M2 ans} \times 10$

$$\text{M3} = 1.006 \times 10^{-3} \times 10 = 1.006 \times 10^{-2}$$

1

**M4** Mass of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = \text{M3 ans} \times 179.8$

$$\text{M4} = 1.006 \times 10^{-2} \times 179.8 = 1.81 \text{ g}$$

1

**M5** % of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = (\text{M4 ans} / 1.381) \times 100$

$$\text{M5} = 1.81 \times 100 / 1.381 = 131 \% (130 \text{ to } 132)$$

1



Buchner funnel and horizontal filter paper

*Allow Hirsch funnel and horizontal filter paper.*

*Do not allow standard Y-shaped funnel.*

*If there is not a clear air-tight seal (labelled or drawn) between the funnel and the flask maximum 1 mark.*

1

- (ii)  $M_r \text{KMnO}_4 = 158(.0)$

1

Mass =  $0.225 \times 158 / 3 = 11.9 \text{ (g)}$

*Lose M2 if no working shown.*

*Allow consequential mark on an incorrect  $M_r$  for  $\text{KMnO}_4$*

1

Precision mark: three significant figures

*Allow if mass incorrect.*

1

- (iii) (Unpleasant) taste

*Ignore smell.*

1

- (b) Difficult to see meniscus / line on graduated flask

*Do not allow reference to over filling.*

1

[7]

47

- (a) To reduce any  $\text{Fe}^{3+}$  ions to  $\text{Fe}^{2+}$  ions

*Allow 'to ensure that all of the iron present is in the form of  $\text{Fe}^{2+}$  ions' or 'to ensure that no  $\text{Fe}^{3+}$  ions are present'.*

1

- (b) Zinc would react with  $\text{MnO}_4^- / \text{Fe}^{3+}$  produced in titration

*Do not allow 'would increase titre value'.*

*Do not allow 'zinc would react' without further qualification.*

1

[2]

48

- (a) Co-ordinate / dative / dative covalent / dative co-ordinate

*Do not allow covalent alone*

1

- (b) (lone) pair of electrons on oxygen/O

*If co-ordination to  $\text{O}^{2-}$ ,  $\text{CE}=0$*

1

forms co-ordinate bond with Fe / donates electron pair to Fe

*'Pair of electrons on O donated to Fe' scores M1 and M2*

1



- (c)  $180^\circ / 180 / 90$

*Allow any angle between 85 and 95*

*Do not allow 120 or any other incorrect angle*

*Ignore units eg  $^\circ\text{C}$*

1

- (d) (i)  $3 : 5 / 5 \text{ FeC}_2\text{O}_4$  reacts with  $3 \text{ MnO}_4^-$

*Can be equation showing correct ratio*

1

- (ii) **M1** Moles of  $\text{MnO}_4^-$  per titration =  $22.35 \times 0.0193/1000 = 4.31 \times 10^{-4}$

Method marks for each of the next steps (no arithmetic error allowed for M2):

*Allow  $4.3 \times 10^{-4}$  (2 sig figs)*

*Allow other ratios as follows:*

*eg from given ratio of 7/3*

1

**M2** moles of  $\text{FeC}_2\text{O}_4 =$  ratio from (d)(i) used correctly  $\times 4.31 \times 10^{-4}$

$$\text{M2} = 7/3 \times 4.31 \times 10^{-4} = 1.006 \times 10^{-3}$$

1

**M3** moles of  $\text{FeC}_2\text{O}_4$  in  $250 \text{ cm}^3 = \text{M2 ans} \times 10$

$$\text{M3} = 1.006 \times 10^{-3} \times 10 = 1.006 \times 10^{-2}$$

1

**M4** Mass of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = \text{M3 ans} \times 179.8$

$$\text{M4} = 1.006 \times 10^{-2} \times 179.8 = 1.81 \text{ g}$$

1

**M5** % of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = (\text{M4 ans}/1.381) \times 100$

$$\text{M5} = 1.81 \times 100/1.381 = 131 \% (130 \text{ to } 132)$$

1



(OR for M4 max moles of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 1.381/179.8 (= 7.68 \times 10^{-3})$

for M5 % of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = (\text{M3 ans/above M4ans}) \times 100$

eg using correct ratio 5/3:

Moles of  $\text{FeC}_2\text{O}_4 = 5/3 \times 4.31 \times 10^{-4} = 7.19 \times 10^{-4}$

Moles of  $\text{FeC}_2\text{O}_4$  in  $250 \text{ cm}^3 = 7.19 \times 10^{-4} \times 10 = 7.19 \times 10^{-3}$

Mass of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 7.19 \times 10^{-3} \times 179.8 = 1.29 \text{ g}$

% of  $\text{FeC}_2\text{O}_4 \cdot 2\text{H}_2\text{O} = 1.29 \times 100/1.381 = 93.4$  (allow 92.4 to 94.4)

Note correct answer ( 92.4 to 94.4) scores 5 marks

*Allow consequentially on candidate's ratio*

eg **M2**  $= 5/2 \times 4.31 \times 10^{-4} = 1.078 \times 10^{-3}$

**M3**  $= 1.0078 \times 10^{-3} \times 10 = 1.078 \times 10^{-2}$

**M4**  $= 1.078 \times 10^{-2} \times 179.8 = 1.94 \text{ g}$

**M5**  $= 1.94 \times 100/1.381 = 140 \%$  (139 to 141)

*Other ratios give the following final % values*

1:1 gives 56.1% (55.6 to 56.6)

5:1 gives 281% (278 to 284)

5:4 gives 70.2% (69.2 to 71.2)

[10]

49

(a)

*For reactions 1 to 3 must show complex ions as reactants and products*

*Take care to look for possible identification on flow chart*

### Reaction 1

ammonia solution

1

**W** is  $[\text{Co}(\text{NH}_3)_6]^{2+}$

1

$[\text{Co}(\text{H}_2\text{O})_6]^{2+} + 6\text{NH}_3 \rightarrow [\text{Co}(\text{NH}_3)_6]^{2+} + 6\text{H}_2\text{O}$

*Correct equation scores all 3 marks*

1

### Reaction 2



*Allow oxygen, Do not allow air*

$\text{H}_2\text{O}_2$

1

**X** is  $[\text{Co}(\text{NH}_3)_6]^{3+}$

1



Correct equations score all 3 marks

1

**Reaction 3**


HCl

Do not allow  $\text{Cl}^-$  but mark on


1

Y is  $[\text{CoCl}_4]^{2-}$

1



Correct equation scores previous mark



This equation scores all three marks

1

**Reaction 4**


$\text{Na}_2\text{CO}_3$  Or  $\text{NaOH}/\text{NH}_3$

Do not allow  $\text{CaCO}_3$  as a reagent but mark on


1

Z is  $\text{CoCO}_3$   $\text{Co}(\text{OH})_2/\text{Co}(\text{H}_2\text{O})_4(\text{OH})_2$


1



Allow waters to stay co-ordinated to Co. This mark also previous mark



1



Allow multiples

1


The activation energy is lower (for the catalysed route)

Or  $\text{Co}^{3+}$  attracts  $\text{SO}_3^{2-}$  /  $\text{Co}^{2+}$  attracts  $\text{SO}_3^{2-}$  / oppositely charged ions attract

1



1



Allow these equations in either order

1

[16]

50

- (a) Orange dichromate

Allow max 2 for three correct colours not identified to species but in correct order

1

Changes to purple / green / ruby / red-violet / violet Chromium(III)

(Note green complex can be  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]^{2+}$  etc)



Do not allow green with another colour

1

That changes further to blue Chromium(II)

Allow max 1 for two correct colours not identified but in correct order

1



Ignore any further reduction of  $\text{Cr}^{2+}$

1



Ignore additional steps e.g. formation of  $\text{CrO}_4^{2-}$

1


- (b) Green precipitate

1

(Dissolves to form a) green solution


Solution can be implied if 'dissolves' stated

1



Penalise  $\text{Cr}(\text{OH})_3$  once only

1



Allow  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+} + 6\text{OH}^- \rightarrow [\text{Cr}(\text{OH})_6]^{3-} + 6\text{H}_2\text{O}$

Allow formation of  $[\text{Cr}(\text{H}_2\text{O})_2(\text{OH})_4]^-$  and  $[\text{Cr}(\text{H}_2\text{O})(\text{OH})_5]^{2-}$  in balanced equations

Ignore state symbols, mark independently

1





- (c) (ligand) substitution / replacement / exchange

*Allow nucleophilic substitution*

1

The energy levels/gaps of the d electrons are different (for each complex)*Ignore any reference to emission of light*

1

So a different wavelength/frequency/colour/energy of light is absorbed (when d electrons are excited)

OR light is absorbed and a different wavelength/frequency/colour/energy (of light) is transmitted/reflected

1

- (d)
- $E_{O_2} (/ H_2O) > E_{Cr^{3+}} (/ Cr^{2+}) / e.m.f = 1.67 V$

*Allow  $E_{(cell)} = 1.67$* 

1

So  $Cr^{2+}$  ions are oxidised by oxygen/air*Allow any equation of the form:*

1

With  $[Cr(H_2O)_6]^{2+}$  get  $CrCO_3$ *If named must be chromium(II) carbonate*

1

with  $[Cr(H_2O)_6]^{3+}$  get  $Cr(H_2O)_3(OH)_3 / Cr(OH)_3$ *Allow 0 to 3 waters in the complex*

1

and  $CO_2$ *Can score M3, M4, M5 in equations even if unbalanced*

1

 $Cr(III)$  differs from  $Cr(II)$  because it is acidic / forms  $H^+$  ions

1

because  $Cr^{3+}$  ion polarises water*Ignore charge/size ratio and mass/charge*

1

[19]

- (a) Variable / many oxidation states

1



*Equations can be in either order*

*Allow multiples*

1



1

- (c) (i) In a different phase / state from reactants

1

- (ii) Impurities poison / deactivate the catalyst / block the active sites

*Allow (adsorbs onto catalyst AND reduces surface area)*

1

- (d) (i) The catalyst is a reaction product

1

- (ii)  $Mn^{2+} / Mn^{3+}$  ion(s)

1



*Equations can be in either order*

1



1

[9]

52

- (a) Negative ions repel one another

1


- (b) Positive ions attract negative ions in catalysed process

*Allow activation energy decreases.*

*Allow alternative route with lower  $E_a$*

*Ignore references to heterogenous catalysis.*

1



*Allow multiples including fractions.*

*Ignore state symbols.*

1



*Allow multiples including fractions.*

*Ignore state symbols.*

*Allow the correct equation involving  $I_3^-$*



1

[4]



53

- (a) A ligand is an electron pair / lone pair donor

*Allow uses lone / electron pair to form a co-ordinate bond*

1

A bidentate ligand donates two electron pairs (to a transition metal ion) from different atoms / two atoms (on the same molecule / ion)

*QoL*

1

- (b)  $\text{CoCl}_4^{2-}$  diagram


1

Tetrahedral shape

1

$109^\circ 28'$

1



*Four chlorines attached to Co with net 2- charge correct*

*Charge can be placed anywhere, eg on separate formula*

*Penalise excess charges*

*Allow  $109^\circ$  to  $109.5^\circ$*

$[\text{Co}(\text{NH}_3)_6]^{2+}$  diagram


1

Octahedral shape

1

$90^\circ$

1



*Six ammonia /  $\text{NH}_3$  molecules attached to Co with 2+ charge correct*

*Allow  $180^\circ$  if shown clearly on diagram*

*$\text{CE} = 0$  if wrong complex but mark on if only charge is incorrect*

- (c) In different complexes the d orbitals / d electrons (of the cobalt) will have different energies / d orbital splitting will be different

1




Light / energy is absorbed causing an electron to be excited

1

Different frequency / wavelength / colour of light will be absorbed / transmitted / reflected

1

- (d) 1 mol of  $\text{H}_2\text{O}_2$  oxidises 2 mol of  $\text{Co}^{2+}$



1

$$M_r \text{ CoSO}_4 \cdot 7\text{H}_2\text{O} = 281$$

If  $M_r$  wrong, max 3 for M1, M4, M5

1

$$\text{Moles Co}^{2+} = 9.87 / 281 = 0.03512$$

1

$$\text{Moles H}_2\text{O}_2 = 0.03512 / 2 = 0.01756$$

M4 is method mark for (M3) / 2 (also scores M1)

1

$$\text{Volume H}_2\text{O}_2 = (\text{moles} \times 1000) / \text{concentration}$$

$$= 0.01756 \times 1000 / 5.00$$

$$= 3.51 \text{ cm}^3 / (3.51 \times 10^{-3} \text{ dm}^3)$$

Units essential for answer

M5 is method mark for (M4)  $\times 1000 / 5$

Allow 3.4 to 3.6  $\text{cm}^3$

If no 2:1 ratio or ratio incorrect Max 3 for M2, M3 & M5

Note: Answer of 7  $\text{cm}^3$  scores 3 for M2, M3, M5 (and any other wrong ratio max 3)

Answer of 16.8  $\text{cm}^3$  scores 3 for M1, M4, M5 (and any other wrong  $M_r$  max 3)

Answer of 33.5  $\text{cm}^3$  scores 1 for M5 only (so wrong  $M_r$  AND wrong ratio max 1)

1

[16]

54

- (a)  $\Delta E = h\nu$

Allow =  $hf$

1

$$\nu = \Delta E / h = 2.84 \times 10^{-19} / 6.63 \times 10^{-34} = 4.28 \times 10^{14} \text{ s}^{-1} / \text{Hz}$$

Allow  $4.3 \times 10^{14} \text{ s}^{-1} / \text{Hz}$

Answer must be in the range:

$$4.28 - 4.30 \times 10^{14}$$

1



- (b) (One colour of) light is absorbed (to excite the electron)

*If light emitted, CE = 0*

1

The remaining colour / frequency / wavelength / energy is transmitted (through the solution)

*Allow light reflected is the colour that we see.*

1

- (c) Bigger

1

Blue light would be absorbed

**OR** light that has greater energy than red light would be absorbed

**OR** higher frequency (of light absorbed / blue light) leads to higher  $\Delta E$

*Can only score M2 if M1 is correct.*

1

- (d) Any **three** from:

- (Identity of the) metal
- Charge (on the metal) / oxidation state / charge on complex
- (Identity of the) ligands
- Co-ordination number / number of ligands
- Shape

3 max

[9]

55

- (a) Cobalt has variable oxidation states


*Allow exists as Co(II) and Co(III)*

1

(It can act as an intermediate that) lowers the activation energy

*Allow (alternative route with) lower  $E_a$*

1



*Allow multiples; allow molecular formulae*

*Allow equations with  $\text{H}_3\text{O}^+$*

1



1

- (b) (i)  $[\text{Co}(\text{H}_2\text{O})_6]^{2+} + 3\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2 \rightarrow [\text{Co}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_3]^{2+} + 6\text{H}_2\text{O}$

*Do not allow en in equation, allow  $\text{C}_2\text{H}_5\text{N}_2$*

1

The number of particles increases / changes from 4 to 7

*Can score M2 and M3 even if equation incorrect or missing  
provided number of particles increases*

1



So the entropy change is positive / disorder increases / entropy increases

1

- (ii) Minimum for **M1** is 3 bidentate ligands bonded to Co

*Ignore all charges for M1 and M3 but penalise charges on any ligand in M2*

1

Ligands need not have any atoms shown but diagram must show 6 bonds from ligands to Co, 2 from each ligand

Minimum for **M2** is one ligand identified as  $\text{H}_2\text{N}-----\text{NH}_2$

*Allow linkage as  $-\text{C}-\text{C}-$  or just a line.*

1

Minimum for **M3** is one bidentate ligand showing two arrows from separate nitrogens to cobalt

1

- (c) Moles of cobalt =  $(50 \times 0.203) / 1000 = \underline{0.01015}$  mol

*Allow 0.0101 to 0.0102*

1

Moles of AgCl =  $4.22/143.4 = 0.0294$

*Allow 0.029*

*If not AgCl (eg  $\text{AgCl}_2$  or  $\text{AgNO}_3$ ), lose this mark and can only score **M1**, **M4** and **M5***

1

Ratio =  $\text{Cl}^-$  to Co = 2.9 : 1

*Do not allow 3 : 1 if this is the only answer but if 2.9:1 seen somewhere in answer credit this as **M3***

1

$[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  (square brackets not essential)

1

Difference due to incomplete oxidation in the preparation

*Allow incomplete reaction.*

*Allow formation  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$  etc.*

*Some chloride ions act as ligands / replace  $\text{NH}_3$  in complex.*

*Do not allow 'impure sample' or reference to practical deficiencies*

1

[15]

- (a) Stoppered flask or similar with side arm

*Allow gas outlet through stopper.*

1



Calibrated container for collection eg gas syringe

*Allow collection over water, but must use calibrated vessel for collection.*

*Lose 1 mark if apparatus is not gas tight.*

1

- (b) Plot a graph of 'volume (of gas)' against 'time'

1

Determine the slope (gradient) at the beginning

1

- (c) Repeat with same volume **or** concentration of hydrogen peroxide and at the same temperature

*Ignore references to results.*

*Do not allow 'keep everything the same' or words to that effect.*

*Must mention volume or concentration and temperature.*

1

Add cobalt(II) chloride to one experiment

1

[6]

57

- (a) (i) Two rings only around nitrogen or sulfur

*Lose this mark if more than 2 atoms are ringed.*

*Do not allow two atoms at the same end of the ion.*

1

- (ii) 275.8

*Accept this answer only. Do not allow 276*

1

- (iii) Carboxylate /  $\text{COO}^-$

*Allow salt of carboxylic acid or just carboxylic acid.*

1

- (b)  $(32.1 / 102.1) = 31.4\%$

*Do not penalise precision but do not allow 1 significant figure.*

1

- (c) Zineb is mixed with a solvent / water

*Max=2 if M1 missed*

1

Use of column / paper / TLC

*Lose M1 and M2 for GLC*

1

Appropriate collection of the ETU fraction

**OR** Appropriate method of detecting ETU

*Allow ETU is an early fraction in a column or collecting a range of samples over time, lowest retention time / travels furthest on paper or TLC (allow 1 mark for having the longest retention time in GLC).*

1



Method of identification of ETU (by comparison with standard using chromatography)

*If method completely inappropriate, only M1 is accessible*

1

[8]

58

- (a) Water in the gaseous state from the precipitate absorbed by drying agent

OR

Water vapour from the precipitate absorbed by drying agent

*Allow 'water vapour reacts with drying agent'.*

*Do not allow 'absorb water' without qualification.*

1

- (b) (Blue to) pink / pink colour observed

1

[2]

59

- (a)  $\text{FeSO}_4 + \text{Na}_2\text{C}_2\text{O}_4 \rightarrow \text{FeC}_2\text{O}_4 + \text{Na}_2\text{SO}_4$

*Allow multiples, including fractions.*

*Allow  $\text{Fe}^{2+} + \text{C}_2\text{O}_4^{2-} \rightarrow \text{FeC}_2\text{O}_4$*

*Allow correct equation which includes water of crystallisation.*

1

- (b)  $M_r \text{FeSO}_4 \cdot 7\text{H}_2\text{O} = 277.9$

*Allow if shown clearly in the calculation.*

*Allow 278*

1

$$\text{Moles} = 6.95 / 277.9 = 2.5(0) \times 10^{-2}$$

*Do not penalise precision but must be to a minimum of two significant figures.*

*Allow correct calculation using incorrect  $M_r$ .*

*Correct answer without working scores this mark only.*

1

- (c)  $3(.00) \times 10^{-2}$

1

- (d) Theoretical mass =  $2.50 \times 10^{-2} \times 179.8 = 4.50\text{g}$   
as long as  $2.50 \times 10^{-2}$  is the smaller of parts (b) and (c) (**M1**)

*Allow consequential answer from parts (b) and (c).*

*Allow theoretical mass = (smaller of parts (b) and (c))  $\times$  179.8*

*If larger of parts (b) and (c) used, lose **M1** but can score **M2**.*

*Allow answers based on moles of reactant and product.*

1

$$\text{Yield} = 3.31 \times 100 / 4.50 = 73.6\% \text{ (M2)}$$

*Award this mark only if answer given to 3 significant figures.*

*Correct answer without working scores this mark only, provided answer given to 3 significant figures.*

1